

**POSTER PRESENTATION**

**Open Access**

# Combination chemoimmunotherapy with monthly cycles of 5-fluorouracil + interferon-alfa-2b in previously-treated advanced cancer

Walter Quan Jr\*, Vivek Khemka, Laura S Martell, Marci J Pierog

From Society for Immunotherapy of Cancer 28th Annual Meeting  
National Harbor, MD, USA. 8-10 November 2013

The in vitro antineoplastic activity of the combination of 5-fluorouracil (5-FU) and interferon-alfa (IFN) has been attributed to multiple mechanisms including enhanced thymidylate synthase inhibition (Elias) and increased susceptibility to interferon-stimulated natural killer cells (Imai). Our group has previously examined weekly dosing schedules of 5-FU and IFN (Quan). Monthly bolus 5-FU + interferon-alfa-2b has not undergone formal phase II testing. In our current study, 18 patients with metastatic cancer who had experienced disease progression on at least 2 prior lines of systemic therapy, have been treated with 5-FU 500 mg/M2 intravenously over 3-5 minutes followed immediately by IFN-2b 3 million IU subcutaneously daily for 3 days monthly on an outpatient basis. Patient characteristics: 12 males/6 females, median age-57 (range: 35-69), median ECOG performance status-1; most common tumor types: colon (6), adenocarcinoma of the lung (4), rectum (3), pancreas (3); common metastatic sites: liver (11), lymph nodes (10), lungs (9). Median number of prior chemotherapy regimens: 3 (range: 2-8). Most common toxicities: rigors, nausea/emesis, fever, and fatigue. Median number of cycles received: 2 (1-11). Sixteen patients are evaluable for response (2 are too early). Two ongoing partial responses have been seen: One patient with adenocarcinoma of the lung (lung + lymph node metastases) at 8.9+ months and one with colon (liver and lymph node metastases at 1.25+ months). Thymidylate synthase levels are being analyzed. Monthly bolus 5-FU followed by subcutaneous IFN-2b is well-tolerated in previously-treated metastatic cancer and shows some evidence of activity in adenocarcinoma of the lung and colon cancer.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P85

**Cite this article as:** Quan et al.: Combination chemoimmunotherapy with monthly cycles of 5-fluorouracil + interferon-alfa-2b in previously-treated advanced cancer. *Journal for ImmunoTherapy of Cancer* 2013 1 (Suppl 1):P85.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



Western Regional Medical Center, Goodyear, AZ, USA



© 2013 Quan et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.