

INVITED SPEAKER PRESENTATION

Open Access

# S77. Proffered paper: In vitro induced response patterns of antileukemic T-cells – characterization by combination of functional assays, spectratyping and next generation sequencing

S Reuther<sup>1\*</sup>, P Krell<sup>2</sup>, FR Schuster<sup>2</sup>, C Grabrucker<sup>1</sup>, A Liepert<sup>1</sup>, T Kroell<sup>1</sup>, HJ Kolb<sup>1</sup>, A Borkhardt<sup>2</sup>, R Buhmann<sup>1</sup>, H Schmetzer<sup>1</sup>

From 1st Immunotherapy of Cancer Conference (ITOC1)  
Munich, Germany. 12-14 March 2014

## Background

T-cell receptor (TCR) diversity is characterised by somatic alterations in the complementary determining region 3 (CDR3) of the human TCR  $\beta$ -chain. Complemented with the TCR alpha-chain, TCR diversity can hypothetically result in up to  $10^{18}$  different TCR molecules.

Myeloid leukaemic cells can be induced to differentiate into leukaemia-derived dendritic cells (DC<sub>leu</sub>) regaining the stimulatory capacity of professional DCs while presenting the whole leukaemic antigen repertoire.

Our aim was to identify TCR  $\beta$ -chain-rearrangements in T-cells stimulated with leukaemic blasts and DC<sub>leu</sub> in 3 patients with AML and furthermore to detect, amplify or monitor T-cell clones with defined  $\beta$ -profiles in correlation with antileukaemic function, in vitro and in vivo.

## Material and methods

HLA matched or HLA haplo-identical (allogeneic) donor- or autologous T-cells were repeatedly stimulated, either with leukaemic blasts or the corresponding DC<sub>leu</sub> from 3 different AML-patients. Cytotoxicity assay was carried out for measuring the lytic activity of effector T-cells, spectratyping was performed to identify the restriction of the TCR  $\beta$ -repertoire in unstimulated and stimulated T-cells and Sanger sequencing to analyse the  $\beta$ -chain sequence information including the CDR3 regions. Additionally next generation sequencing (NGS) was established to

analyse the accurate TCR sequence information of thousands of TCR  $\beta$ -chains with high coverage.

## Results

No significant differences in T-cell proliferation were observed. The T-cell mediated cytolytic response patterns showed blast lysis (n=1) and blast proliferation (n=2).

Spectratyping revealed a remarkable TCR  $\beta$ -restriction of the CD4<sup>+</sup>- or CD8<sup>+</sup>-TCR repertoire of blast- or DC/DC<sub>leu</sub>-stimulated T-cells, independently of blast or DC/DC<sub>leu</sub> used as stimulators. Although in absolute terms, DC/DC<sub>leu</sub> stimulation induced the highest grade of restriction in the CD8<sup>+</sup> T-cell subset, the CD4<sup>+</sup> T-cells seemed to be relatively more affected.

In vitro stimulation with DC/DC<sub>leu</sub> resulted in an identical TCR ( $\beta$ -chain restriction pattern) as identified in vivo in a patient sample 3 months after allogeneic stem cell transplantation (SCT).

## Conclusion

A combined strategy using spectratyping and NGS with functional tests may provide useful information about the specificity and efficacy of the intra-individual variable induced T-cell response.

Spectratyping allows the identification of a restricted  $\beta$ -repertoire by Gaussian-like distribution, NGS allows sequencing of TCR repertoires with high coverage, novel software allows the analysis of the exact length and sequence composition (the combination of the  $\beta$ - and J $\beta$ -genes) of the  $\beta$ -chains, especially of the CDR3, and the exclusion of non-functional transcripts.

<sup>1</sup>University of Munich, Med. III Hematopoietic Transplantations, Munich, Germany

Full list of author information is available at the end of the article

The identification of defined V $\beta$ -T-cell clones may lead to selection procedures generating Graft-versus-Leukaemia reaction- but not Graft-versus-Host disease-mediating T-cells for adoptive immunotherapy after SCT.

#### Authors' details

<sup>1</sup>University of Munich, Med. III Hematopoetic Transplantations, Munich, Germany. <sup>2</sup>University of Duesseldorf, Department of Pediatric Oncology Hematology and Immunology, Duesseldorf, Germany.

Published: 12 March 2014

doi:10.1186/2051-1426-2-S2-I15

**Cite this article as:** Reuther *et al.*: S77. Proffered paper: In vitro induced response patterns of antileukemic T-cells – characterization by combination of functional assays, spectratyping and next generation sequencing. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 2):115.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

