

POSTER PRESENTATION

Open Access

Elucidating the role of Neuropilin-1 in intra-tumoral regulatory T cell stability


Abigail E Overacre*, Maria Chikina, Greg M Delgoffe, Dario AA Vignali

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Regulatory T cells (T_{regs}) play an integral role in the adaptive immune system through suppression of self-reactive immune responses in order to prevent autoimmunity and maintain homeostasis. However, they are deleterious in cancer through suppression of the anti-tumor immune response. In fact, we show that deletion of 50% of T_{regs} results in normal tumor growth. Therefore, it is advantageous to understand the role of T_{regs} in the tumor microenvironment in order to create targeted cancer therapies. Our lab has shown that the Neuropilin-1 (Nrp1) pathway is required for T_{reg} stability in the tumor microenvironment, but is dispensable for maintaining immune homeostasis in the periphery, identifying it as a prime therapeutic target.

In order to further understand the role of Nrp1-deficient T_{regs} intratumorally, we constructed a competitive

environment by utilizing *Foxp3*, which is located on the X chromosome, and as a result of X-inactivation, female *Foxp3^{Cre-YFP}* heterozygous mice are cellular heterozygotes. We generated *Nrp1^{L/L}Foxp3^{Cre-YFP/+}* heterozygous mice comprised of 50% WT T_{regs} and 50% Nrp1-deficient T_{regs} . Surprisingly, when given B16 melanoma, heterozygous mice **phenocopy** *Nrp1^{L/L}Foxp3^{Cre-YFP}* homozygous mice (Figure 1A). This suggests that **Nrp1-deficient T_{regs} are playing an active role in shifting the anti-tumor immune response by destabilizing surrounding WT T_{regs} as determined by DNA methylation status** (Figure 1B). Neither WT nor Nrp1-deficient T_{regs} in the tumor from *Nrp1^{L/L}Foxp3^{Cre-YFP/+}* mice can suppress in a standard microsuppression assay *ex vivo*, unlike WT T_{regs} from *Foxp3^{Cre-YFP}* mice. Through various co-culture experiments, we revealed that destabilization of WT T_{regs}



University of Pittsburgh, Pittsburgh, PA, USA



© 2015 Overacre et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

is possibly due to a soluble factor derived from Nrp1-deficient T_{regs} . Our data revealed that Nrp1-deficient T_{regs} produce large amounts of $IFN\gamma$ in the tumor microenvironment. Indeed, **when treated with $IFN\gamma$, WT T_{regs} lose suppressive capacity.** In order to uncover potential novel pathways leading to this phenotype, we are performing global transcript studies using RNASeq. Overall, we have shown that Nrp1 is required for intratumoral T_{reg} stability, and in its absence, there is an alteration in the tumor microenvironment, leading to an **enhanced anti-tumor immune response.** These studies uncover a novel potential target for future cancer immunotherapies that preserves peripheral immune health.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P274

Cite this article as: Overacre *et al.*: Elucidating the role of Neuropilin-1 in intra-tumoral regulatory T cell stability. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P274.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

