

POSTER PRESENTATION

Open Access

Metabolic insufficiency underlies intratumoral cytotoxic T cell dysfunction

Tushar Gupta, Nicole E Scharping, Rebecca S Moreci, Greg M Delgoffe*

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

T cells have the remarkable ability to recognize and remove abnormal cells with precision, a feature that is very desirable for the treatment of cancer. However, while T cells specific for tumor antigens are primed and can infiltrate tumors, they are quickly rendered dysfunctional, through both cell intrinsic and cell extrinsic mechanisms. One way that tumors cripple T cell function is through the generation of an immunosuppressive microenvironment that is chronically inflamed, hypoxic, and nutrient poor. T cell activation and subsequent generation of effector function is bioenergetically demanding, requiring large amounts of metabolic intermediates to effectively proliferate, produce cytokines, and lyse target cells. We hypothesized that T cell dysfunction in cancer is due, in part, to metabolic insufficiency caused by chronic activation in metabolically dearth conditions. Using single-cell metabolic assays and extracellular flux analysis, we show that CD8⁺ cytotoxic T cells that infiltrate tumors demonstrate a progressive loss of mitochondrial function and mass, concomitant with upregulation of markers that correlate with T cell exhaustion. This mitochondrial dysfunction occurs independently of coinhibitory molecule signaling and specifically in the tumor microenvironment. This results in a failure to generate an adequate pool of ATP and in inability to effectively translate effector gene transcripts. This in stark contrast to T cells responding to an acute viral infection, where activated effector T cells demonstrate increased mitochondrial mass and ATP reserve. Further, artificial induction of mitochondrial dysfunction in T cells results in upregulation of coinhibitory molecules and an 'exhausted-like' phenotype, suggesting that metabolic insufficiency underlies the dysfunctional phenotype in cancer. Taken together, our data support a model in which tumor-infiltrating T cells have metabolic needs

that cannot be met, resulting in failed effector function and tumor growth. Our studies also suggest that modulation or reprogramming of the altered metabolism of intratumoral T cells represents a potential strategy to reinvigorate dysfunctional T cells for the immunotherapeutic treatment of cancer.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P399

Cite this article as: Gupta et al.: Metabolic insufficiency underlies intratumoral cytotoxic T cell dysfunction. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P399.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



University of Pittsburgh, Pittsburgh, PA, USA



© 2015 Gupta et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.