

POSTER PRESENTATION

Open Access

# Epitope-optimization creates highly immunogenic alpha fetoprotein antigen to break immune tolerance and potently activates CD8 T cells to prevents autochthonous hepatocellular carcinoma

Yuan Hong<sup>1</sup>, Yibing Peng<sup>1</sup>, Sheng Z Guo<sup>3</sup>, Junfeng Pang<sup>1</sup>, Jose Guevara-patino<sup>2</sup>, David H Munn<sup>1</sup>, Nahid Mivechi<sup>1</sup>, David Bartlett<sup>3</sup>, Yukai He<sup>1\*</sup>

From Society for Immunotherapy of Cancer 28th Annual Meeting  
National Harbor, MD, USA. 8-10 November 2013

In this study, we investigated whether mouse alpha fetoprotein (mAFP), the shared self/tumor antigen of hepatocellular carcinoma (HCC), could be rationally engineered to create effective vaccine to break tolerance and potently activate CD8 T cells to prevent clinically-relevant carcinogen-induced autochthonous HCC. We found that the computer-guided epitope-optimization created optimized opt-mAFP and that immunization with lentivector (lv) expressing opt-mAFP, but not wt-mAFP, potently activated CD8 cells specific for three novel H-2b restricted CD8 epitopes, which cross-recognized wt-mAFP epitopes naturally processed and presented by wt-mAFP+ tumor cells. Immunization with opt-mAFP-lv, but not wt-mAFP-lv, completely protected mice from wt-mAFP+ tumor challenge and effectively prevented carcinogen-induced autochthonous HCC. Prime-boost with opt-mAFP-lv and vaccinia vector opt-mAFP-vv significantly increased the wt-mAFP-specific CD8 T cells that were highly responsive to emerging HCC tumor cells in the liver, enhancing prevention of autochthonous HCC. Our data demonstrate that epitope-optimization creates immunogenic opt-mAFP that is able to break tolerance and activate potent CD8 responses, which can cross-recognize wt-mAFP peptides, but also recognize and kill mAFP+ tumor cells. Our study provides a practical roadmap to develop effective human vaccines that should have a better chance of success than the current human HCC vaccines based on native wt-AFP.

#### Authors' details

<sup>1</sup>Cancer Center, Georgia Regents University, Augusta, GA, USA. <sup>2</sup>Surgery, Loyola University, Chicago, IL, USA. <sup>3</sup>Surgery, University of Pittsburgh, Pittsburgh, PA, USA.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P216

**Cite this article as:** Hong et al.: Epitope-optimization creates highly immunogenic alpha fetoprotein antigen to break immune tolerance and potently activates CD8 T cells to prevents autochthonous hepatocellular carcinoma. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P216.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Cancer Center, Georgia Regents University, Augusta, GA, USA  
Full list of author information is available at the end of the article