

POSTER PRESENTATION

Open Access

# Quantitative single-cell characterization of CAR+ T cell effector functions

Ivan Liadi<sup>1</sup>, Harjeet Singh<sup>3</sup>, Gabrielle Romain<sup>1</sup>, Nicolas Rey-Villamizar<sup>2</sup>, Amin Merouane<sup>2</sup>, Badrinath Roysam<sup>2</sup>, Laurence J Cooper<sup>3</sup>, Navin Varadarajan<sup>1\*</sup>

From Society for Immunotherapy of Cancer 28th Annual Meeting  
National Harbor, MD, USA. 8-10 November 2013

Adoptive cell therapy (ACT) utilizing chimeric antigen receptor (CAR) T cells rendered specific for CD19 have demonstrated significant anti-tumor effects in patients with CD19+ chronic lymphocytic leukemia (CLL). In spite of the clinical promise of ACT in achieving complete responses, their efficacy remains unpredictable and new approaches are needed to address a priori define the therapeutic potential of T-cell based therapies. In our current work, we characterize the in vitro functionality of CD19-specific (CD19RCD28) CAR+ T cells propagated using artificial antigen presenting cells expressing membrane bound IL-21, by employing a novel methodology single-cell nanowell screening that determines their cytotoxic ability and cytokine secretion capability at single-cell resolution. We show that CAR+ T cells exert specific cytotoxicity against NALM6 cells ( $31 \pm 8\%$ ) when co-incubated at a 1:1 ratio in nanowell containers. Furthermore, single CAR+ T cells were capable of engaging and killing multiple targets;  $17 \pm 8\%$  of T cells killed two target cells and  $9 \pm 3\%$  killed three target cells within the 6 hour window of observation. In parallel, microengraving was used to determine the cytokine secretion profile of these same cells. Hierarchical clustering of the two functions indicated that interferon-gamma ( $IFN\gamma$ ) secretion is not correlated to cytotoxicity or the ability of T cells to kill multiple target cells. Simultaneously, monitoring apoptosis on CAR+ T cells allowed us to quantify their activation-induced cell death (AICD). CAR+ T cells that secreted  $IFN\gamma$  upon target ligation did not undergo AICD whereas T cells that engaged in repeated killing showed an increased propensity to

undergo AICD ( $p = 0.04$ ). Dynamic time-lapse imaging of the interactions between CAR+ T cells and tumor cells indicated that the majority of CAR+ T cells have high basal motility, form long-lived interactions with tumor cells (50 - 100 min) that lead to motility arrest and subsequent tumor-cell apoptosis. However, contact lifetimes or overall contact duration were not reliable predictors of subsequent tumor-cell apoptosis. Finally, kinetics of serial killing suggest that motile CAR+ T cells that form short-lived contacts exhibit rapid killing with very little motility arrest in vitro. In summary, our SNS based methodology allows the deep functional characterization of clinical grade CAR+ T cells and can be used to: (1) determine in vitro functions of CAR+ T cells that correlate with clinical efficacy and (2) inform CAR design to maximize effector functionality while minimizing AICD.

#### Authors' details

<sup>1</sup>Chemical & Biomolecular Engineering, University of Houston, Houston, TX, USA. <sup>2</sup>Dept of Electrical Engineering, University of Houston, Houston, TX, USA. <sup>3</sup>Pediatrics, University of Texas M.D. Anderson Cancer Center, Houston, TX, USA.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P36

Cite this article as: Liadi et al.: Quantitative single-cell characterization of CAR+ T cell effector functions. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P36.

<sup>1</sup>Chemical & Biomolecular Engineering, University of Houston, Houston, TX, USA

Full list of author information is available at the end of the article