

POSTER PRESENTATION

Open Access

Safety and disease response to MEDI-551, an anti-CD19 antibody, in chronic lymphocytic leukemia patients previously treated with rituximab

Andres Forero^{1*}, Mehdi Hamadani², Thomas Kipps³, Michelle Fanale⁴, Antonio Cuneo⁵, Jaime Perez de Oteyza⁶, Douglas Gladstone⁷, Marc Andre⁸, Naresh Bellam¹, Trishna Goswami⁹, Ramy Ibrahim⁹, Amy Schneider¹⁰, Meina Liang¹⁰, Steven Eck⁹, Nairouz Elgeioushi⁹, Ronald Herbst⁹, Bruce D Cheson¹¹

From Society for Immunotherapy of Cancer 28th Annual Meeting National Harbor, MD, USA. 8-10 November 2013

Background

The expression of CD19 on chronic lymphocytic leukemia (CLL) cells offers a novel therapy for relapsed CLL patients (pts) previously treated with rituximab. MEDI-551 is an affinity-optimized anti-CD19 Ab with enhanced Ab-dependent cellular cytotoxicity (ADCC) effector function.

Methods

Response and toxicity of single-agent MEDI-551 in multiply relapsed CLL pts with prior rituximab therapy was assessed in a phase 1/2 (ph 1/2), open-label, dose-escalation and expansion study. Combination therapy was assessed in an ongoing phase 2 (ph 2) study comparing MEDI-551 or rituximab+bendamustine in relapsed/refractory CLL pts. For the ph 1/2 study, B cell depletion was assessed with flow cytometry and BAFF biomarker analysis; response was assessed using the 2008 IWG criteria.

Results

In the ph 1/2 study, 26 CLL pts received ≥ 1 dose of MEDI-551. In the ph 2 study, 44 pts received study drug as of 20Mar2013. Loss of CD19 detection due to depletion and/or occupancy with MEDI-551 was rapid and apparent after cycle 1. B cell depletion occurred 1 day after dose 1 and was associated with increased serum BAFF concentrations. In the ph 1/2 study, of 21 MEDI-551-treated CLL pts evaluable for response, 5 achieved partial remission

and 13 had stable disease. Commonly reported adverse events (AEs) in MEDI-551 pts were infusion-related reactions (IRRs; 62%), nausea (23%), pyrexia (23%), and neutropenia (23%) in the 26 ph 1/2 pts; in the 29 ph 2 pts, they were nausea (62%), IRRs (31%), pyrexia (28%), chills (28%), and fatigue (28%). 11 pts had \geq grade 3 AEs in the ph 1/2 study and 16 in the ph 2. Common treatment-related AEs: IRRs (58%) and nausea (12%) in the ph 1/2; nausea (52%), IRRs (28%), chills (24%), and fatigue (24%) in the ph 2. Three treatment-unrelated AEs of general health deterioration (ph 1/2), subarachnoid hemorrhage (ph 1/2), and sepsis (ph 2), resulted in death.

Conclusions

MEDI-551 as a single agent demonstrated B-cell depletion, increased serum BAFF levels, clinical activity, and an acceptable risk-benefit profile in relapsed/refractory CLL pts. Preliminary results of the ongoing ph 2 study of MEDI-551+bendamustine demonstrated an acceptable safety profile.

Authors' details

¹University of Alabama at Birmingham, Birmingham, AL, USA. ²West Virginia University, Morgantown, WV, USA. ³Moore's UCSD Cancer Center, San Diego, CA, USA. ⁴University of Texas, Houston, TX, USA. ⁵Azienda Ospedaliera Universitaria Arcispedale Sant'Anna, Ferrara, Italy. ⁶Centro Integral Oncologico Clara Campal, Madrid, Spain. ⁷Johns Hopkins University, Baltimore, MD, USA. ⁸CHU Mont-Godinne, Yvoir, Belgium. ⁹MedImmune, Gaithersburg, MD, USA. ¹⁰MedImmune, Gaithersburg, CA, USA. ¹¹Georgetown University Hospital, Washington DC, MD, USA.

¹University of Alabama at Birmingham, Birmingham, AL, USA
Full list of author information is available at the end of the article

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P43

Cite this article as: Forero et al.: Safety and disease response to MEDI-551, an anti-CD19 antibody, in chronic lymphocytic leukemia patients previously treated with rituximab. *Journal for ImmunoTherapy of Cancer* 2013 1(Suppl 1):P43.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

