

POSTER PRESENTATION

Open Access

Combination therapies augment the anti-tumor activity of agonist CD27 mAb in human CD27 transgenic mouse models

Li-Zhen He¹, Lawrence J Thomas², James Testa¹, Jeffrey Weidlick¹, Crystal Sisson¹, Russell Hammond², Laura Vitale¹, Henry Marsh², Tibor Keler^{1*}

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

CDX-1127 is a fully human antibody to CD27, a TNF receptor superfamily member expressed on the majority of T cells and subsets of NK cells and B cells. We have previously characterized the co-stimulatory activities of CDX-1127 with human T cell cultures and a human CD27 transgenic mouse model (hCD27-Tg). Similar to the findings originally shown by M.J. Glennie and colleagues using an agonist anti-mouse CD27 mAb, CDX-1127 has potent antitumor activity as monotherapy in several syngeneic tumor models in hCD27-Tg mice. In the current studies, we sought to enhance the anti-tumor efficacy of CDX-1127 in challenging tumor settings by combination with clinically relevant therapies. Specifically, we focused on therapies that could decrease or control tumor growth while providing a source of antigen to drive anti-tumor immunity (e.g. chemotherapy or targeted therapy) and immune modifiers that may allow the CD27 driven T cell response to overcome self-regulation (e.g. checkpoint inhibitors or immune activators). In the EG7 delayed treatment model (average tumor size of ~ 50 mm³ when treatment initiated), the combination of CDX-1127 with cyclophosphamide significantly improved survival (>70% survival) compared to either agent alone (<30 % survival). Notably, we found that the combination therapy was associated with increases in the ratio of effector to regulatory T cells in the tumors compared to either single agent group. Additional combination studies with various agents are ongoing, and initial studies with CDX-1127 combined with anti-CTLA-4 mAb has shown superior anti-tumor activity (median survival 36.5 days in combination versus

20 days with either single agent). These studies, along with the good safety profile of CDX-1127 reported in a Phase 1 clinical trial, supports the design of future combination studies in patients with cancer.

Authors' details

¹Celldex Therapeutics, Inc., Phillipsburg, NJ, USA. ²Celldex Therapeutics, Inc., Needham, NJ, USA.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P76

Cite this article as: He et al.: Combination therapies augment the anti-tumor activity of agonist CD27 mAb in human CD27 transgenic mouse models. *Journal for ImmunoTherapy of Cancer* 2013 1(Suppl 1):P76.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Celldex Therapeutics, Inc., Phillipsburg, NJ, USA
Full list of author information is available at the end of the article