

Chromatin remodeling (SWI/SNF) complexes, cancer, and response to immunotherapy

Nithya Krishnamurthy ^{1,2}, Shumei Kato,^{2,3} Scott Lippman,^{1,3} Razelle Kurzrock⁴

To cite: Krishnamurthy N, Kato S, Lippman S, *et al.* Chromatin remodeling (SWI/SNF) complexes, cancer, and response to immunotherapy. *Journal for ImmunoTherapy of Cancer* 2022;**10**:e004669. doi:10.1136/jitc-2022-004669

Accepted 01 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Icahn School of Medicine at Mount Sinai, New York, New York, USA

²Yale University, New Haven, Connecticut, USA

³Department of Medicine, Division of Hematology/Oncology, and Center for Personalized Cancer Therapy, University of California, Moores Cancer Center, La Jolla, California, USA

⁴Worldwide Innovative Network for Personalized Cancer Therapy, San Diego, California, USA

Correspondence to

Nithya Krishnamurthy;
nithya.krishnamurthy@icahn.
mssm.edu

ABSTRACT

Chromatin regulation involves four subfamilies composed of ATP-dependent multifunctional protein complexes that remodel the way DNA is packaged. The SWI/SNF chromatin remodeling complex subfamily mediates nucleosome reorganization and hence activation/repression of critical genes. The SWI/SNF complex is composed of the BRG-/BRM-associated factor and Polybromo-associated BAF complexes, which in turn have multiple subunits. Significantly, ~20% of malignancies harbor alterations in >1 of these subunits, making the genes encoding SWI/SNF family members among the most vulnerable to genomic aberrations in cancer. ARID1A is the largest subunit of the SWI/SNF complex and is altered in ~40%–50% of ovarian clear cell cancers and ~15%–30% of cholangiocarcinomas, in addition to a variety of other malignancies. Importantly, outcome was improved after immune checkpoint blockade (ICB) in patients with *ARID1A*-altered versus wild-type tumors, and this result was independent of microsatellite instability or tumor mutational burden. Another subunit—PBRM1—is mutated in ~40% of clear cell renal cell carcinomas and ~12% of cholangiocarcinomas; there are contradictory reports regarding ICB responsiveness. Two other SWI/SNF subunits of interest are SMARCA4 and SMARCB1. SMARCA4 loss is the hallmark of small cell carcinoma of the ovary hypercalcemic type (and is found in a variety of other malignancies); *SMARCA4* germline alterations lead to rhabdoid tumor predisposition syndrome-2; *SMARCB1* germline alterations, rhabdoid tumor predisposition syndrome-1. Remarkable, although anecdotal, responses to ICB have been reported in both *SMARCA4*-aberrant and *SMARCB1*-aberrant advanced cancers. This review focuses on the role that SWI/SNF chromatin remodeling subunits play in carcinogenesis, the immune microenvironment, and in immunotherapy responsiveness.

INTRODUCTION

Chromatin remodeling elements refer to a group of proteins that remodel the way DNA architecture is packaged in order to permit access of condensed genomic DNA to the transcription machinery and thereby regulate gene expression. Chromatin itself is a complex of DNA and protein; its primary function is to package long DNA molecules into more compact structures.¹ A primary

protein component of chromatin is histones. Histones bind to DNA and function as ‘anchors’ around which the strands are wound. There are several levels of chromatin organization, and one of the important ones relate to nucleosomes. A nucleosome is the basic repeating unit of eukaryotic chromatin. A solitary nucleosome is made up of about 150 base pairs of DNA sequence blanketed around a core of histone proteins. DNA must be compacted into nucleosomes to fit into the nucleus and each human cell contains about 30 million nucleosomes.²

Although humans have four chromatin remodeler protein families—SWI/SNF (SWI/SNF chromatin remodeling complex), CHD ((Chromodomain-Helicase-DNA binding), Imitation SWI/SNF 1 and INO80 (inositol requiring 80)^{3–6}—the best studied of these families of chromatin remodeler proteins are the SWI/SNF complexes. SWI/SNF is a subfamily of ATP-dependent chromatin remodeling complexes that associate to remodel the way DNA is packaged. This complex possesses a DNA-stimulated ATPase activity that can destabilize histone-DNA interactions in nucleosomes in an ATP-dependent manner. The SWI/SNF complex mediates nucleosome reorganization, allowing genes to be activated or repressed.^{3 6}

The SWI/SNF complex is composed of polymorphic BRG-associated/BRM-associated factor (BAF) and Polybromo-associated BAF (PBAF) complexes. BAF is made up of SMARCA2 or 4, ACTL6A, SMARCC1, SMARCC2, SMARCC1, DFP1/2/3, ARID1A/B, SMARCD1/2/3 and SMARCB1 subunits; PBAF is made up of SMARCA2 or 4, ACTL6A, SMARCC1, SMARCC2, SMARCC1, DFP1/2/3, SMARCD1/2/3 and SMARCB1, ARID2, BRD7, and PBRM1 subunits (figure 1). Importantly, ~20% of malignancies have an aberration in one of the genes encoding these subunits.⁷ Furthermore, abnormalities in SWI/SNF chromatin

SWI/SNF Complex- Components and Functions




Figure 1 SWI/SNF complex is an evolutionarily conserved ATP-dependent complex that contains multiple subunits. It has a key role in chromatin remodeling and regulation of transcription by recruitment of transcription factors, coactivators and repressors and histone modifiers. BRG1-associated or BRM1-associated factors (BAF) and Polybromo-associated BAF (PBAF) are two subclasses of this complex and differ in a few subunits as shown. These complexes consist of one of the two mutually exclusive catalytic ATPase subunits: SMARCA2 (Brahma or BRM) or SMARCA4 (BRG1), and other subunits such as SMARCB1, SMARCC1, and SMARCC2. PBAF complexes can be distinguished from BAF complexes because the former contains PBRM1 and ARID2 but lack ARID1A/B. SWI/SNF complexes regulate chromatin access by controlling the processes of histone dimer ejection, nucleosome ejection, and repositioning of nucleosomes by sliding. ARID1A binds DNA and may regulate the chromatin remodeling activity of the SWI/SNF complex through recruitment and binding of transcriptional factors. ARID subunits help with binding of the ATPase subcomplex. PBRM1 is essential for the stability of the SWI/SNF chromatin remodeling complex SWI/SNF-B (PBAF). ACTL6A, an actin domain, SMARCC1, and DFP1/2/3 are accessory subunits common to both BAF and PBAF and are rarely mutated in cancers. SWI/SNF, SWItch/Sucrose Non-Fermentable.

remodeling complex subunits have shown promise as markers for ICB responsiveness (table 1, figure 1).^{7–48}

The development of ICB for cancer therapy, with several molecules now approved, signals a dynamic change in the field of immuno-oncology. Ipilimumab, which is anti-cytotoxic T-lymphocyte antigen-4 (CTLA-4), was the first approved immune checkpoint inhibitor for treating patients with advanced melanoma.⁴⁹ Pembrolizumab, nivolumab, atezolizumab, avelumab, durvalumab, and cemiplimab, all of which are anti-programmed death 1 (anti-PD1) or anti-programmed death ligand 1 (anti PDL1) antibodies, have been subsequently approved.^{50–51} The overall response rates across tumors on checkpoint blockade therapy is, however, only in the order of 15%–20%, making it imperative to develop reliable and biologically based biomarkers to predict response.⁵² Some established biomarkers include deficient mismatch repair/microsatellite instability-high (dMMR/MSI-H)⁵³ and high tumor mutational burden (TMB) (≥ 10 mutations/megabase),^{54,55} but there are also reports of new markers such as major histocompatibility

presentation of neoantigens and immune response gene signature panels.^{56–59} Additionally, genomic markers such as *PDL1* amplification,⁶⁰ chromosome 9p21.3 loss,⁶¹ and novel markers such as chromatin remodeling gene aberrations—*ARID1A*, *PBRM1*, *SMARCA4*, and *SMARCB1*^{14 21 37 46}—may also predict ICB responsiveness. Much of this data has just recently emerged and some of the clinical linkages between chromatin remodeling genes and immunotherapy remain anecdotal or a matter of debate. Even so, these linkages merit further understanding and exploration because of their high potential impact in the patient care setting. Here, we provide an overview of the emerging data on the valuable role of SWI/SNF chromatin remodeling genes in cancer and in immune responsiveness. We also balance the discussion with commentary on the pitfalls of using some of the chromatin remodeling gene aberrations such as *PBRM1* loss as a marker for immunotherapy response, as the linkage may be with other aspects of tumor proliferation like angiogenesis.

Table 1 Examples of genes controlling SWI/SNF chromatin remodeling and immune checkpoint blockade (ICB) responsiveness

Name of gene	% Patients with cancer and alterations in the gene	Chromatin remodeling complex functions	Response to immunotherapy	Comment
<i>ARID1A</i>	~3%–7% of all cancers ^{1,7} Highest rates of mutation are seen in clear cell cancers of the ovary (~50% of patients), endometroid tumors (~40%), gastric cancers (~30%) ^{7–9}	<ul style="list-style-type: none"> ▶ Largest subunit in SWI/SNF chromatin-remodeling complex^{6,10} ▶ DNA binding subunit 	<p>Preclinical studies-treatment with anti-PD-L1 antibody show reduced tumor burden and increased survival in <i>ARID1A</i> loss mice ovarian tumors versus <i>ARID1A</i> wild type.¹¹</p> <p><i>ARID1A</i> aberrations resulted in limited chromatin accessibility to interferon (IFN) responsive genes causing impaired IFN expression, poor T cell response and reduced tumor immune response.¹²</p> <p>Pan-cancer patients with <i>ARID1A</i> alterations had significantly prolonged OS on ICB¹³</p> <p><i>ARID1A</i> alterations were a positive predictor for longer PFS after checkpoint blockade (HR (95% CI), 0.61 (0.39 to 0.94), p=0.02).¹⁴</p> <p>PFS benefit after ICB was not dependent on MSI or TMB.¹⁴</p> <p>Loss of <i>ARID1A</i> associated with: High PDL1 in gastric cancers¹⁵ and EBV¹⁶ (and both EBV and PDL1 predict for response in gastric cancer¹⁷)</p>	<p><i>ARID1A</i> is implicated in interactions with mismatch repair gene, MSH2, possibly compromising its function.¹⁸</p>
<i>PBRM1</i>	~3% of diverse cancers ¹⁹ Highest rates of mutation seen in clear cell renal cell carcinoma tumor (seen in 40% of patients), cholangiocarcinomas (12%). ^{19–21}	<ul style="list-style-type: none"> ▶ Nucleosome-recognition subunit in the PBAF SWI/SNF chromatin-remodeling complex.¹⁹ 	<p>Mixed data showing both that <i>PBRM1</i> predicts response to ICB and does not predict response to ICB</p> <p>ICB response was associated with loss-of-function mutations in the <i>PBRM1</i> gene (p=0.012) in metastatic ccRCC, a cancer not associated with MSI or high TMB.²²</p> <p>OS was significantly better with <i>PBRM1</i> loss in 324 metastatic clear cell renal cell carcinomas on nivolumab (not reached vs 25 mos, p=0.05).²³</p> <p>However, in IMmotion in metastatic ccRCCs showed worse outcome for <i>PBRM1</i>-altered tumors in atezolizumab versus sunitinib group.²⁴</p> <p>A recent study with multivariate models of ccRCC patients treated with ICB (n=189), loss-of-function (LOF) mutations in <i>PBRM1</i> were not associated with OS (HR=1.24, p=0.47) or time to treatment failure (HR=0.85, p=0.44), Pan cancer across 11 solid tumors (n=2936), LOF mutations not associated with improved OS (HR=0.9, p=0.7).²⁵</p> <p>In an NSCLC retrospective analysis, <i>PBRM1</i> mutation predicted for worse prognosis on ICB regardless of high or low TMB (median OS of <i>PBRM1</i>-mutant versus wild-type patients was 6 vs 13 months)²⁶</p>	<p><i>PBRM1</i> is thought to confer resistance to T cell-induced apoptosis, and a <i>PBRM1</i> deletion in a B16F10 melanoma mouse model increases chances of response to anti-PD-1 and anti-CTLA4 agents.¹²</p> <p>Enhancement of immunostimulatory genes which play a role in hypoxia response and JAK-STAT signaling in <i>PBRM1</i>-mutant ccRCC cell lines.²⁷</p> <p><i>PBRM1</i> loss is consistent with a decreased immunogenic tumor microenvironment and instead upregulated angiogenesis.²⁸</p>
<i>SMARCA4</i>	About 5%–7% of all cancers ²⁹ Highest rates of mutation seen in SCCOHT (close to 100% patients) ³⁰ and NSCLC (10% of patients), ³¹ but also seen in undifferentiated endometrial sarcomas, SMARCA4-deficient thoracic sarcoma, gastric cancers, and a subset of multiple other tumor types ^{32–34}	<ul style="list-style-type: none"> ▶ <i>SMARCA4</i> (BRG1) is important ATPase subunit of the mammalian SWI/SNF chromatin remodeling complex. ▶ Uses the energy from ATP hydrolysis to disrupt nucleosomes at target regions.³⁵ 	<p><i>SMARCA4</i> is hallmark of SCCOHT; 4/4 patients with objective response include three with durable CRs.³⁶</p> <p><i>SMARCA4</i> mutations were more prevalent in responders to PDL1 blockade in Keynote 012 trial in HNSCC.³⁷</p> <p><i>SMARCA4</i>-mutated NSCLC was associated with improved outcomes with ICB in MSK-IMPACT study.³⁸</p> <p>Case reports show response to nivolumab or pembrolizumab in <i>SMARCA4</i>-mutated NSCLC.^{39,40}</p> <p><i>SMARCA4</i>-deficient thoracic sarcoma (<i>SMARCA4</i>-DTS) case reports show response to nivolumab in third line therapy and another report shows rapid response to pembrolizumab in PDL1 positive <i>SMARCA4</i>-DTS.^{41,42}</p>	<p><i>SMARCA4</i> alteration is sole mutation in rare SCCOHT which is characterized by high PDL-1 expression and T-cell infiltrate⁴³; these tumors respond anecdotally to checkpoint blockade⁴³</p>
<i>SMARCB1</i>	Loss of <i>SMARCB1</i> expression is seen in malignant rhabdoid tumors and epithelioid sarcomas. renal medullary carcinoma, cribriform neuroepithelial tumors, malignant mesotheliomas, synovial sarcomas, extracellular myxoid chondrosarcomas and familial schwannomatosis <i>SMARCB1</i> loss is at the protein and DNA level in most tumors ^{44,45}	<ul style="list-style-type: none"> ▶ <i>SMARCB1</i> (INI1) (part of SWI/SNF complex) tumor-suppressor gene is located at 22q11.2 	<p>Malignant rhabdoid tumors have low TMB across cancers, but case reports of 3 patients with <i>SMARCB1</i>-loss aggressive pediatric cancers demonstrate evidence of response to ICB.⁴⁶</p> <p>47% (14/30) of patients with <i>SMARCB1</i>-loss tumors had positive PD-L1 staining.^{45,46}</p> <p>Anecdotal reports show a pediatric renal medullary cancer with <i>SMARCB1</i> loss responding to atezolizumab.⁴⁷</p> <p>An adult patient with recurrent <i>SMARCB1</i>-loss renal medullary carcinoma had a complete response to nivolumab lasting greater than 9 months despite low TMB.⁴⁷</p> <p>An advanced refractory, <i>SMARCB1</i>-deficient epithelioid sarcoma accomplished a complete remission to combined ipilimumab and nivolumab⁴⁸</p>	

ARID1A, AT-rich Interactive Domain-containing protein; ccRCC, clear cell renal cell carcinoma; CR, complete remission; EBV, Epstein Barr virus; HCC, hepatocellular carcinoma; HNSCC, head and neck squamous cell cancer; Jak-STAT, Janus kinase-signal transducer and activator of transcription; MSI, microsatellite instability; MSK, Memorial Sloan Kettering; NSCLC, non-small cell lung cancer; OS, overall survival; *PBRM1*, polybromo 1; PD-L1, programmed death ligand; PFS, progression-free survival; SCCOHT, small cell cancer ovarian hypercalcemic type; *SMARCA4*, SWI/SNF Related, Matrix Associated, Actin Dependent Regulator Of Chromatin; *SMARCA4*-DTS, *SMARCA4*-deficient thoracic sarcomas; SWI/SNF, SWItching/Sucrose Non-Fermenting; TIM-3, T-cell immunoglobulin mucin-3; TMB, tumor mutational burden.

Mutations in chromatin-remodeling SWI/SNF complex genes and tumor immunity

Alterations in various subunits of the SWI/SNF chromatin remodelers are found in about 20% of human cancers.⁷ The literature suggests that the most frequently altered subunit gene is *ARID1A* (BAF250A), which is aberrant in up to 7% of cancers, with other subunits such as *PBRM1* and *SMARCA4* altered in ~3%–7% of all malignant neoplasms.^{7 18 29}

Importantly, tumor immunity may be affected by alterations in SWI/SNF chromatin remodeling genes in multiple ways: (1) loss of *PBRM1* (polybromo 1) and *ARID2* (AT-rich Interactive Domain-containing protein), which leads to increased expression of genes that play a role in IFN γ (interferon-gamma) signaling and thus could increase responses to immunotherapy,^{12 62} as high IFN γ activates Janus kinase (JAK)–signal transducer and activator of transcription (STAT), which turns on PD-L1 expression⁶³; (2) *SMARCB1*-mutant rhabdoid tumors show infiltration by subpopulations of clonally expanded T cells, suggesting a tumor-specific immune response⁶⁴; and (3) *ARID1A* may interact with MSH2, a MMR protein. *ARID1A* alterations are associated with improved outcome after checkpoint blockade; these responses appear to be independent of TMB and microsatellite stability status.^{11 14 65}

ARID1A

ARID1A is the largest subunit of the SWI/SNF complex. It is located on chromosome 1p and is involved in chromatin remodeling activity via a conserved DNA binding domain, which enables binding transcription factors as well as transcriptional coactivator/corepressor complexes.^{1 10} Germline mutations in *ARID1A* are associated with truncating mutations that cause a severe form of Coffin-Siris syndrome that is characterized by feeding difficulties, bowel obstruction, and early severe respiratory problems as well as congenital heart disease (table 2).⁶⁶

ARID1A alterations occur in 3%–7% of cancers overall^{1 7}; a variety of malignancies are affected, including but not limited to 46–50% of ovarian clear cell cancers,^{7–9} 15%–27% of cholangiocarcinomas,⁶⁷ endometrial carcinomas,⁶⁸ colorectal cancers,⁶⁹ 10%–35% of gastric cancers,⁹ and many others.

From a functional viewpoint, as a subunit of SWI/SNF chromatin remodeler, *ARID1A* facilitates target-specific binding of SWI/SNF complexes to chromatin, thereby modifying the accessibility of chromatin to nuclear factors. In malignancies, *ARID1A* possesses the features of a gatekeeper in that it regulates cell cycle progression, and of a caretaker in that, it prevents genomic instability.⁷⁰

Preclinical data in mice models suggest that *ARID1A* may be involved with MMR gene MMR and MSH2

Table 2 Examples of germline mutations in chromatin remodeling genes and their associated conditions

Gene	Name of syndrome	Features of syndrome	Examples of cancers in patients with germline mutations	Comment
<i>ARID1A</i>	Coffin-Siris syndrome ⁶⁶	<i>ARID1A</i> alterations were found in ~7% of Coffin-Siris syndrome ⁶⁶ Features of Coffin-Siris syndrome include developmental delays, hypoplastic digits, hirsutism, and microcephaly ⁶⁶	Cancers not described	Coffin-Siris syndrome is caused by mutations in the <i>ARID1A</i> , <i>ARID1B</i> , <i>SMARCA4</i> , <i>SMARCB1</i> or <i>SMARCE1</i> genes. ⁶⁶
<i>ARID1B</i>	Coffin-Siris syndrome	<i>ARID1B</i> mutations were the most common cause of Coffin-Siris syndrome (51%–75%)	Cancers not described	
<i>PBRM1</i>	Familial renal cell carcinoma	A case report showed a family with renal cell carcinoma with germline mutations of <i>PBRM1</i> ⁷⁵	Renal cell carcinoma	Germline <i>PBRM1</i> truncating mutation (p. Asp1333Glyfs) associated with renal cell carcinoma ⁷⁵
<i>SMARCB1</i>	Coffin Siris syndrome Rhabdoid tumor predisposition syndrome (RTPS1) ⁸³	Typically, infants and children present with cancers, while some affected patients present with benign schwannomas. ⁹⁶	Rhabdoid tumors, ^{83 96}	
<i>SMARCA4</i>	Rhabdoid tumor predisposition syndrome (RTPS2) ⁸³	Typically, infants and children present with cancers, and 11% of Coffin Siris ⁸² syndrome have <i>SMARCA4</i> germline mutations.	Rhabdoid tumors ⁸³ SCCOHT ³⁰	

ARID1A, AT-rich Interactive Domain-containing protein; PBRM1, polybromo 1; SCCOHT, small cell carcinoma of the ovary hypercalcemic type; SMARCA4, SWI/SNF Related, Matrix Associated, Actin Dependent Regulator of Chromatin; SWI/SNF, mating-type SWitching (SWI) and sucrose fermentation (Sucrose Non-Fermenting - SNF).

protein.^{11 65} In ovarian cancer cell lines of a syngeneic mice model, there are increased tumor-infiltrating lymphocytes, and higher TMB and PDL1 expression.^{70 71} Additionally, *ARIDIA* interacts on the carboxy-terminal with the Enhancer of Zeste 2 PRC2 subunit (EZH2) that functions as a catalytic subunit of the polycomb repressive complex 2 (PRC2). This interaction results in inhibition of the repressive function of EZH2 in IFN-responsive genes in human cancer cell lines.⁷¹ Furthermore, in gastric cancers, *ARIDIA* aberrations are associated with Epstein-Barr positivity and high PDL1 expression which also strengthens the responsiveness to ICB.^{15–17} *ARIDIA* is also vital for creating an open chromatin state on DNA damage and mediates the non-homologous end-joining (NHEJ) pathway.⁷² *ARIDIA* deficient cells cannot mount NHEJ repair and combination treatment with low-dose radiation and Olaparib, a PARP inhibitor, elicited tumor responses in ovarian cancer mice model.⁷² These preclinical observations are the basis of the ATARI clinical trial with an ATR (Ataxia telangiectasia and Rad3 related) inhibitor in combination with a PARP inhibitor in gynecological cancers.⁷³ Finally, alterations in *ARIDIA* activate the phosphatidylinositol 3-kinase PI3K/serine-threonine kinase AKT/mammalian target of rapamycin mTOR pathways.⁷⁴

In the clinical pan-cancer setting, MSI-H, as well as high TMB, were significantly more frequent in *ARIDIA*-altered versus *ARIDIA* wild-type tumors (20% vs 0.9%, $p < 0.001$; and 26% vs 8.4%, $p < 0.001$, respectively). Median PFS after checkpoint blockade immunotherapy was significantly longer in the patients with *ARIDIA*-altered tumors than in those with *ARIDIA* wild-type tumors (11 months vs 4 months, $p = 0.006$). Multivariate analysis showed that *ARIDIA* alterations predicted a better outcome after ICB and this result was not dependent on MSI or TMB.^{14 65}

PBRM1

PBRM1 is a nucleosome-recognition subunit of the PBAF SWI/SNF chromatin-remodeling complex. Polybromo-1 (*PBRM1*), found on chromosome 3p21, functions as a tumor suppressor gene and is most commonly mutated in clear cell cancer of the kidney (~40% of patients) and cholangiocarcinomas (~12% of patients).^{19–21} Recently, a germline frameshift mutation in *PBRM1* was identified as a predisposing factor for renal cell cancer (RCC).⁷⁵ Preclinical studies show that *PBRM1* itself may confer resistance to T cell-induced apoptosis and *PBRM1*-deficient murine melanomas had increased infiltration by cytotoxic T cells.¹² Deletion of *PBRM1* in a B16F10 melanoma mouse model increased susceptibility to anti-PD-1 and anti-CTLA4 agents.¹²

Some clinical studies have shown a response to ICB. For example, immunotherapy response was associated with loss-of-function mutations in the *PBRM1* gene ($p = 0.012$) in metastatic clear cell RCC (ccRCC),²² cancer not associated with MSI-H or high TMB, and overall survival (OS) was significantly better with *PBRM1* loss in patients with metastatic ccRCC on nivolumab versus those not receiving

nivolumab (not reached vs 25 months, $p = 0.05$).²³ Enhancement of immunostimulatory genes involved in hypoxia response and JAK-STAT signaling in *PBRM1*-mutant ccRCC lines may elucidate the immunotherapy response.^{27 28} However, contrary to the above, there are conflicting reports suggesting that *PBRM1* mutations correlate with a decreased immunogenic tumor micro-environment in human RCC lines.^{24 25} *PBRM1*-deficient mouse subcutaneous renal tumors show resistance to ICB.²⁸ Moreover, analysis of the IMmotion150 renal cell carcinoma study also suggests that *PBRM1* mutations correlate with attenuated immunotherapy responsiveness.^{24 28} In multivariate models of ccRCC patients treated with ICB ($n = 189$) at Memorial Sloan Kettering, loss-of-function mutations in *PBRM1* were not associated with longer OS (HR=1.24, $p = 0.47$) or time-to-treatment failure (HR=0.85, $p = 0.44$).²⁵

Similarly, in a non-small-cell lung cancer (NSCLC) retrospective analysis, *PBRM1* mutations predicted a worse prognosis on ICB regardless of high or low TMB (median survival of *PBRM1*-mutant versus wild-type patients was 6 versus 13 months).²⁶

There are notable limitations to using *PBRM1* as a biomarker for immunotherapy response as the positive responses are in the setting of antiangiogenic therapy and not in the front-line setting. As seen in the IMMOTION 150 study, *PBRM1* mutations are associated with high angiogenesis and renal cell carcinomas may benefit more from antiangiogenic therapy.^{76 77} Patients with *PBRM1* mutated tumors had better progression-free survival with a multi-receptor tyrosine kinase inhibitor sunitinib than with ICB, suggesting that *PBRM1* loss may be more useful as a marker for anti-angiogenesis therapy.^{78–80}

A recent study of *PBRM1* loss in diverse cancer types concluded that *PBRM1* loss portended worse outcomes on ICB in ccRCC, adenocarcinomas of the lung, cutaneous melanoma, and bladder cancers even with high TMB.⁸¹ Thus, immunotherapy must be used with caution for these tumors and large-scale trials are needed to resolve this conflict.

SMARCA4 and SMARCB1

The *SMARCA4* gene is situated on chromosome 19p and encodes the BRG1 protein. It belongs to the SWI/SNF chromatin remodeling complex and functions as an ATPase. There are two main categories of cancer-related *SMARCA4* alterations: class 1 mutations—truncating mutations, fusions, and homozygous deletion (loss of function); and class 2 mutations—missense mutations (dominant-negative/gain of function through loss of function can also occur). Protein loss generally occurs with class 1 mutations.³⁵

Overall, about 5%–7% of cancers have *SMARCA4* alterations.²⁹ Deficiency of *SMARCA4* has been implicated in oncogenesis in small cell carcinoma of the ovary hypercalcemic type (SCCOHT),³⁰ undifferentiated endometrial carcinomas, and uterine sarcomas,³³ the aggressive

SMARCA4-deficient thoracic sarcoma,³² and in some non-small-cell lung adenocarcinomas (NSCLCs).³¹

Germline mutations in *SMARCA4* have been detected in ~11% of those with Coffin-Siris syndrome (table 2). In particular, missense mutations with gain-of-function or dominant-negative effects are seen; some features of the syndrome are microphthalmia, intellectual disability, and lack of predisposition to cancers.⁸² Inactivating *SMARCA4* mutations are seen in SCCOHT (almost all of which have *SMARCA4* mutations, with about 40% having germline mutations).⁸² Germline mutations in *SMARCA4* are also associated with the rhabdoid tumor predisposition syndrome-2 (RTPS2) (table 2).⁸³ Most individuals diagnosed with *SMARCA4*-related RTPS inherited a pathogenic variant from an unaffected parent. The hallmark of RTPS2 is a notably increased predisposition to rhabdoid tumors which are rare but highly aggressive tumors arising in children under the age of four. Rhabdoid tumors can occur in almost any anatomic location but occur most frequently in the central nervous system, with >50% presenting in the cerebellum. Overall, ~35% of rhabdoid tumors are associated with germline SWI/SNF mutations.⁸⁴

SCCOHT is a rare but aggressive monogenic tumor characterized by a low TMB, but still, exhibits a highly immunogenic tumor microenvironment with high PDL1 expression and infiltration by T cells.^{36 43} Anecdotally, there are case reports of a sustained clinical response to anti-PDL1 therapy in patients with SCCOHT, suggesting that *SMARCA4* mutations, even in the setting of low mutational burden can lead to an immunogenic tumor environment.^{36 85}

Despite associations with aggressive disease, class 1 *SMARCA4* mutations in NSCLC were associated with response to ICB.^{38 39} *SMARCA4* genomic alterations are found in ~10% of all NSCLC.⁸⁶ Recently a case report showed a more than 14-month sustained response when treated with nivolumab in a *SMARCA4*-deficient NSCLC with a high tumor mutation burden.³⁹ Similarly, case reports of sustained tumor regression have been seen with nivolumab in *SMARCA4*-deficient thoracic sarcoma (*SMARCA4*-DTS) in third-line therapy after prior cytotoxic therapy.⁴¹ In another case report, there was a rapid response to pembrolizumab in a *SMARCA4*-deficient thoracic sarcoma overexpressing PDL1.⁴⁰ Taken together these observations suggest that *SMARCA4*-deficient tumors merit prospective evaluation in clinical trials for their ICB responsiveness.

SMARCB1 (SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily B member 1) or INI1 (integrase interactor 1) is located at chromosomal position 22q11.2 and functions as a core subunit protein in the SWI/SNF complex.^{44 46} Loss of *SMARCB1* expression/altered *SMARCB1* is predominantly seen in pediatric renal and extrarenal malignant rhabdoid tumors (almost all of which show *SMARCB1* alterations), epithelioid sarcomas, atypical central nervous system teratoid/rhabdoid tumors; renal medullary carcinoma,

synovial sarcomas malignant mesotheliomas, sinonasal carcinomas, and cribriform neuroepithelial tumors.^{44 45} In particular, loss of *SMARCB1* is present in a high number of epithelioid sarcomas.⁸⁷ Interestingly, loss of *SMARCB1* results in high expression of EZH2 leading to upregulation of several oncogenic pathways including Wnt/beta-catenin, Myc, and the Sonic Hedgehog pathways.^{88 89} Recent studies have shown promising results for EZH2 inhibitors in many cancer types including epithelioid sarcoma (which have *SMARCB1* aberrations).⁹⁰ Indeed, tazemetostat, an EZH2 inhibitor was granted accelerated approval in 2020 for use in metastatic and locally advanced epithelioid sarcomas.⁹¹

Additionally, recent studies point to other ways a *SMARCB1* deficiency can modulate tumor immunogenicity, that is, by selecting for de-repression of endogenous retroviral elements (ERV). ERV de-repression can cause the accumulation of double-stranded RNA in the cytoplasm, which can stimulate a cell's innate immune response through the IFN- α and IFN- λ pathways.⁹² DNA methyltransferase inhibitors have been shown to upregulate a cell's immune signaling through a type 1 interferon response which is accompanied by ERV expression in many cancers, including ovarian cancer. Furthermore, ERV expression, especially expression of ERV3-2, is associated with ICB response in clear cell renal carcinomas.^{93 94} Additionally, association with DNA *transposase*, a piggyBac transposable element derived 5 (PGBD5) gene, can cause genetic rearrangements in lethal rhabdoid tumors. There is evidence to suggest that PGBD5 expression can cause *SMARCB1* somatic inactivation in these tumors; this includes PGBD5-specific signals (PSS) sequences found in *SMARCB1* deficient rhabdoid tumors.⁹⁵

The archetype of a *SMARCB1*-deficient tumor is the malignant rhabdoid tumor, initially described in the kidney but also is seen in soft tissue, viscera, and the brain (where it is designated as an atypical teratoid rhabdoid tumor). These cancers are overwhelmingly diagnosed in the very young, and most have a fatal course. Pathologically, most but not all contain a population of 'rhabdoid' cells, typically cells with vast cytoplasm, eccentric vesicular nuclei, perinuclear spherical inclusions, and large inclusion-like nucleoli.⁹⁶ Germline mutations occur as part of the RTPS and are of two subclasses. RTPS1 is caused by alterations in *SMARCB1* and RTPS2 is caused by alterations in *SMARCA4*. Most cases are due to *SMARCB1*, but the features of RTPS1 and RTPS2 are clinically similar. In comparison to sporadic isolated rhabdoid tumors, the syndromic form is associated with an increased risk of developing multiple tumors at younger ages and schwannomas (benign nerve sheath tumors) that present primarily in adulthood.⁸³ RTPS is inherited in an autosomal dominant manner. Many individuals have the disorder as the result of a de novo germline *SMARCB1* pathogenic variant.^{83 96} Rhabdoid tumors are among the least mutated tumors with very low TMB but surprisingly have significant PDL1 expression and, in a

small case series, patients showed responses to anti-PDL1 therapies.^{46 92}

Renal medullary carcinoma is a rare but aggressive cancer characterized by balanced translocations that disrupt the tumor suppressor role of *SMARCB1*.⁹⁷ There are anecdotal reports of responses to anti-PDL1 therapy in renal medullary cancers in both pediatric and adult patients but there are also reports of a lack of response.^{47 98} In a mouse model of rhabdoid tumors, there was complete tumor regression in 67% to 80% of treated mice on anti-PDL1 therapy.⁶⁴ Additionally, there was clear evidence of tumor-infiltrating CD8⁺ T cells which had high expression of clinically targetable inhibitory immune-checkpoint receptors, including PD-1, LAG-3, and TIM-3.⁶⁴ Finally, an advanced refractor, *SMARCB1*-deficient epithelioid sarcoma had a complete remission after taking a combined anti-PD1 immune checkpoint inhibitor therapy and anti-CTLA4 agent.⁴⁸

CONCLUSIONS

The SWI/SNF chromatin-remodeling complex is vital for transcriptional activation of genes normally repressed by chromatin. The role that these complexes play in responses to commonly used immune checkpoint inhibitors is still being delineated, but recent data points to improved outcomes in patients with *ARID1A*-altered malignancies (across tumor types) even in the absence of traditionally used markers of immunotherapy response such as PDL1 and TMB.¹⁴ Some of the underlying mechanisms may include inhibition of the repressive function of EZH2 in IFN-responsive gene function,⁷¹ increased tumor-infiltrating lymphocytes,⁶⁴ and interaction with MMR genes such as MSH2.¹¹ While *ARID1A* loss has been posited as a reliable biomarker for ICB response,^{14 65} there are contradictory reports regarding PBRM1 loss, with some studies showing response and other pan-cancer trials showing a lack of response to ICB.^{25 26}

SMARCA4 alterations are seen in rare but aggressive cancers such as SCCOHT, and *SMARCA4*-deficient thoracic and uterine sarcomas. *SMARCA4* alterations are also seen in ~10% of NSCLC. Anecdotal studies suggest that some of these cancers may be responsive to ICB administration.^{36 37} The DART prospective clinical trial, which combines nivolumab and ipilimumab in rare tumors, also has a cohort that specifically addresses SCCOHT (NCT02834013). *SMARCB1*-deficient tumors have also been reported anecdotally to respond to ICB, including with compete for remissions in advanced diseases such as epithelioid sarcoma.⁴⁸

Germline mutations in *SMARCB1* and *SMARCA4* are responsible for the rhabdoid tumor predisposition syndromes (RTPS1 and RTPS2, respectively) and there are reports of response to ICB in pediatric tumors that arise in patients with these cancers.^{83 96}

A very recent study looked at nine patients with metastatic pancreatic cancer who harbored SWI/SNF alterations and found that 8/9 showed responsiveness to ICB

even though only three were microsatellite unstable, and pancreatic cancers are usually resistant to ICB.⁹⁹ These findings suggest a role for prospective clinical trials using SWI/SNF alterations as a predictive marker for ICB.

In conclusion, the SWI/SNF chromatin-remodeling complex is responsible for a multitude of functions that play an important role in gene transcription. The various subunits are mutated in a substantial subset of patients with malignancy (~20% of human cancers overall); germline mutations predisposing individuals to tumors also occur. The SWI/SNF chromatin remodeling complex has unique effects on the tumor immune microenvironment, translating to both preclinical and early clinical data suggesting responsiveness to commonly used immune checkpoint inhibitors across a variety of cancer types. Further prospective studies of immunotherapy in patients with SWI/SNF chromatin remodeling subunit gene aberrations and malignancies are urgently warranted.

Contributors NK and RK wrote, reviewed, and revised the article. SK and SL reviewed, edited, and revised the article. All authors read and approved the final article.

Funding Funded in part by National Cancer Institute grant P30 CA023100 and the Joan and Irwin Jacobs Fund philanthropic fund.

Competing interests SK serves as a consultant for Foundation Medicine, NeoGenomics and CureMatch. He receives speaker's fee from Roche and advisory board for Pfizer. He has research funding from ACT Genomics, Sysmex, Konica Minolta and OmniSeq. RK has the following disclosure information: Stock and Other Equity Interests (IDbyDNA, CureMatch, Inc., and Soluventis); Consulting or Advisory Role (Gaido, LOXO, X-Biotech, Actuate Therapeutics, Roche, NeoMed, Soluventis, Pfizer, and Merck); Speaker's fee (Roche); Research Funding (Incyte, Genentech, Merck Serono, Pfizer, Sequenom, Foundation Medicine, Guardant Health, Grifols, Konica Minolta, DeBiopharm, Boehringer Ingelheim, and OmniSeq [All institutional]); Board Member (CureMatch, Inc., and CureMetric, Inc.). SL is the Co-founder of io9, advisor for SymptoHealth, and on the scientific advisory board for Biological Dynamics and Human Longevity Inc. NK has no disclosures to report.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Nithya Krishnamurthy <http://orcid.org/0000-0001-9171-7506>

REFERENCES

- Clapier CR, Cairns BR. The biology of chromatin remodeling complexes. *Annu Rev Biochem* 2009;78:273–304.
- Annunziato A. DNA packaging: nucleosomes and chromatin. *Nature Education* 2008;1:26.
- Clapier CR, Iwasa J, Cairns BR, et al. Mechanisms of action and regulation of ATP-dependent chromatin-remodelling complexes. *Nat Rev Mol Cell Biol* 2017;18:407–22.
- Muchardt C, Yaniv M. The mammalian SWI/SNF complex and the control of cell growth. In: *Seminars in cell & developmental biology*. Academic Press, 1999: 10. 189–95.
- Gursoy-Yuzugullu O, House N, Price BD. Patching broken DNA: nucleosome dynamics and the repair of DNA breaks. *J Mol Biol* 2016;428:1846–60.
- Lorch Y, Maier-Davis B, Kornberg RD. Mechanism of chromatin remodeling. *Proc Natl Acad Sci U S A* 2010;107:3458–62.

- 7 Hodges C, Kirkland JG, Crabtree GR. The many roles of BAF (mSWI/SNF) and PBAF complexes in cancer. *Cold Spring Harb Perspect Med* 2016;6:a026930.
- 8 Jones S, Wang T-L, Shih I-M, et al. Frequent mutations of chromatin remodeling gene ARID1A in ovarian clear cell carcinoma. *Science* 2010;330:228–31.
- 9 Lin C, Song W, Bi X, et al. Recent advances in the ARID family: focusing on roles in human cancer. *Oncol Targets Ther* 2014;7:315.
- 10 Han L, Madan V, Mayakonda A, et al. Chromatin remodeling mediated by ARID1A is indispensable for normal hematopoiesis in mice. *Leukemia* 2019;33:2291–305.
- 11 Shen J, Ju Z, Zhao W, et al. ARID1A deficiency promotes mutability and potentiates therapeutic antitumor immunity unleashed by immune checkpoint blockade. *Nat Med* 2018;24:556–62.
- 12 Pan D, Kobayashi A, Jiang P, et al. A major chromatin regulator determines resistance of tumor cells to T cell-mediated killing. *Science* 2018;359:770–5.
- 13 Jiang T, Chen X, Su C, et al. Pan-cancer analysis of ARID1A alterations as biomarkers for immunotherapy outcomes. *J Cancer* 2020;11:776.
- 14 Okamura R, Kato S, Lee S, et al. ARID1A alterations function as a biomarker for longer progression-free survival after anti-PD-1/PD-L1 immunotherapy. *J Immunother Cancer* 2020;8.
- 15 Wang K, Kan J, Yuen ST, et al. Exome sequencing identifies frequent mutation of ARID1A in molecular subtypes of gastric cancer. *Nat Genet* 2011;43:1219.
- 16 Kim Y-B, Ahn JM, Bae WJ, et al. Functional loss of ARID1A is tightly associated with high PD-L1 expression in gastric cancer. *Int J Cancer* 2019;145:916–26.
- 17 Miliotis CN, Slack FJ. Multi-layered control of PD-L1 expression in Epstein-Barr virus-associated gastric cancer. *J Cancer Metastasis Treat* 2020;6. doi:10.20517/2394-4722.2020.12. [Epub ahead of print: 23 05 2020].
- 18 Allo G, Bernardini MQ, Wu R-C, et al. ARID1A loss correlates with mismatch repair deficiency and intact p53 expression in high-grade endometrial carcinomas. *Mod Pathol* 2014;27:255–61.
- 19 Savas S, Skardasi G. The SWI/SNF complex subunit genes: their functions, variations, and links to risk and survival outcomes in human cancers. *Crit Rev Oncol Hematol* 2018;123:114–31.
- 20 Varela I, Tarpey P, Raine K, et al. Exome sequencing identifies frequent mutation of the SWI/SNF complex gene PBRM1 in renal carcinoma. *Nature* 2011;469:539–42.
- 21 Misumi K, Hayashi A, Shibahara J, et al. Intrahepatic cholangiocarcinoma frequently shows loss of BAP1 and PBRM1 expression, and demonstrates specific clinicopathological and genetic characteristics with BAP1 loss. *Histopathology* 2017;70:766–74.
- 22 Miao D, Margolis CA, Gao W, et al. Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma. *Science* 2018;359:801–6.
- 23 Vano YA, Rioux-Leclercq N, Dalban C, et al. NIVOREN GETUG-AFU 26 translational study: association of PD-1, AXL, and PBRM1 with outcomes in patients (PTS) with metastatic clear cell renal cell carcinoma (mccRCC) treated with nivolumab (N). *J Clin Oncol* 2020;38:618.
- 24 McDermott DF, Huseni MA, Atkins MB, et al. Clinical activity and molecular correlates of response to atezolizumab alone or in combination with bevacizumab versus sunitinib in renal cell carcinoma. *Nat Med* 2018;24:749–57.
- 25 Hakimi AA, Attalla K, DiNatale RG, et al. A pan-cancer analysis of PBAF complex mutations and their association with immunotherapy response. *Nat Commun* 2020;11:1.
- 26 Zhou H, Liu J, Zhang Y, et al. PBRM1 mutation and preliminary response to immune checkpoint blockade treatment in non-small cell lung cancer. *NPJ Precis Oncol* 2020;4:1–4.
- 27 Vuong L, Kotecha RR, Voss MH, et al. Tumor microenvironment dynamics in clear-cell renal cell carcinoma. *Cancer Discov* 2019;9:1349–57.
- 28 Liu X-D, Kong W, Peterson CB, et al. PBRM1 loss defines a nonimmunogenic tumor phenotype associated with checkpoint inhibitor resistance in renal carcinoma. *Nat Commun* 2020;11:1–4.
- 29 Fernando TM, Piskol R, Bainer R, et al. Functional characterization of SMARCA4 variants identified by targeted exome-sequencing of 131,668 cancer patients. *Nat Commun* 2020;11:1–3.
- 30 Jelinic P, Mueller JJ, Olvera N, et al. Recurrent SMARCA4 mutations in small cell carcinoma of the ovary. *Nat Genet* 2014;46:424–6.
- 31 Medina PP, Romero OA, Kohno T, et al. Frequent BRG1/SMARCA4-inactivating mutations in human lung cancer cell lines. *Hum Mutat* 2008;29:617–22.
- 32 Perret R, Chalabreysse L, Watson S, et al. SMARCA4-deficient thoracic sarcomas. *Am J Surg Pathol* 2019;43:455–65.
- 33 Kolin DL, Quick CM, Dong F, et al. SMARCA4-deficient uterine sarcoma and undifferentiated endometrial carcinoma are distinct clinicopathologic entities. *Am J Surg Pathol* 2020;44:263–70.
- 34 Huang S-C, Ng K-F, Yeh T-S, et al. The clinicopathological and molecular analysis of gastric cancer with altered SMARCA4 expression. *Histopathology* 2020;77:250–61.
- 35 Tang L, Nogales E, Ciferri C. Structure and function of SWI/SNF chromatin remodeling complexes and mechanistic implications for transcription. *Prog Biophys Mol Biol* 2010;102:122–8.
- 36 Witkowski L, Goudie C, Ramos P, et al. The influence of clinical and genetic factors on patient outcome in small cell carcinoma of the ovary, hypercalcemic type. *Gynecol Oncol* 2016;141:454–60.
- 37 Hanna GJ, Lizotte P, Cavanaugh M, et al. Frameshift events predict anti-PD-1/L1 response in head and neck cancer. *JCI Insight*. 2018;3.
- 38 Schoenfeld AJ, Bandlamudi C, Lavery JA, et al. The Genomic Landscape of SMARCA4 Alterations and Associations with Outcomes in Patients with Lung Cancer. *Clin Cancer Res* 2020;26:5701–5708.
- 39 Naito T, Umemura S, Nakamura H, et al. Successful treatment with nivolumab for SMARCA4-deficient non-small cell lung carcinoma with a high tumor mutation burden: a case report. *Thorac Cancer* 2019;10:1285–8.
- 40 Henon C, Blay J-Y, Massard C, et al. Long lasting major response to pembrolizumab in a thoracic malignant rhabdoid-like SMARCA4-deficient tumor. *Ann Oncol* 2019;30:1401–3.
- 41 Iijima Y, Sakakibara R, Ishizuka M, et al. Notable response to nivolumab during the treatment of SMARCA4-deficient thoracic sarcoma: a case report. *Immunotherapy* 2020;12:563–9.
- 42 Takada K, Sugita S, Murase K, et al. Exceptionally rapid response to pembrolizumab in a SMARCA4-deficient thoracic sarcoma overexpressing PD-L1: a case report. *Thorac Cancer* 2019;10:2312–5.
- 43 Jelinic P, Ricca J, Van Oudenhove E, et al. Immune-active microenvironment in small cell carcinoma of the ovary, hypercalcemic type: rationale for immune checkpoint blockade. *J Natl Cancer Inst* 2018;110:787–90.
- 44 Kohashi K, Oda Y. Oncogenic roles of SMARCB1/INI1 and its deficient tumors. *Cancer Sci* 2017;108:547–52.
- 45 Agaimy A, Hartmann A, Antonescu CR, et al. SMARCB1 (INI-1)-deficient Sinonasal Carcinoma: A Series of 39 Cases Expanding the Morphologic and Clinicopathologic Spectrum of a Recently Described Entity. *Am J Surg Pathol* 2017;41:458.
- 46 Forrest SJ, Al-Ibraheemi A, Doan D, et al. Genomic and immunologic characterization of INI1-deficient pediatric cancers. *Clin Cancer Res* 2020;26:2882–2890.
- 47 Beckermann KE, Jolly PC, Kim JY, et al. Clinical and immunologic correlates of response to PD-1 blockade in a patient with metastatic renal medullary carcinoma. *J Immunother Cancer* 2017;5:1–5.
- 48 Pecora A, Halpern S, Weber M, et al. Rapid and complete response to combination anti-CTLA-4 and anti-PD-1 checkpoint inhibitor therapy in a patient with stage IV refractory end-stage epithelioid sarcoma: a case report. *J Immunother* 2020;43:286–90.
- 49 Hargadon KM, Johnson CE, Williams CJ. Immune checkpoint blockade therapy for cancer: an overview of FDA-approved immune checkpoint inhibitors. *Int Immunopharmacol* 2018;62:29–39.
- 50 Vaddepally RK, Kharel P, Pandey R, et al. Review of indications of FDA-approved immune checkpoint inhibitors per NCCN guidelines with the level of evidence. *Cancers* 2020;12:738.
- 51 Migden MR, Rischin D, Schmults CD, et al. PD-1 blockade with cemiplimab in advanced cutaneous squamous-cell carcinoma. *N Engl J Med* 2018;379:341–51.
- 52 Meng X, Huang Z, Teng F, et al. Predictive biomarkers in PD-1/PD-L1 checkpoint blockade immunotherapy. *Cancer Treat Rev* 2015;41:868–76.
- 53 Le DT, Uram JN, Wang H, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509–20.
- 54 Goodman AM, Sokol ES, Frampton GM, et al. Microsatellite-Stable tumors with high mutational burden benefit from immunotherapy. *Cancer Immunol Res* 2019;7:1570–3.
- 55 Jardim DL, Goodman A, de Melo Gagliato D, et al. The challenges of tumor mutational burden as an immunotherapy biomarker. *Cancer Cell* 2021;39:154–73.
- 56 Goodman AM, Castro A, Pyke RM, et al. MHC-I genotype and tumor mutational burden predict response to immunotherapy. *Genome Med* 2020;12:1–3.
- 57 Goodman AM, Kato S, Chattopadhyay R, et al. Phenotypic and genomic determinants of immunotherapy response associated with squamousness. *Cancer Immunol Res* 2019;7:866–73.
- 58 Boichard A, Pham TV, Yeerna H, et al. APOBEC-related mutagenesis and neo-peptide hydrophobicity: implications for response to immunotherapy. *Oncimmunology* 2019;8:1550341.

- 59 Pham TV, Boichard A, Goodman A, *et al.* Role of ultraviolet mutational signature versus tumor mutation burden in predicting response to immunotherapy. *Mol Oncol* 2020;14:1680–94.
- 60 Goodman AM, Piccioni D, Kato S, *et al.* Prevalence of PDL1 amplification and preliminary response to immune checkpoint blockade in solid tumors. *JAMA Oncol* 2018;4:1237–44.
- 61 Han G, Yang G, Hao D, *et al.* 9p21 loss confers a cold tumor immune microenvironment and primary resistance to immune checkpoint therapy. *Nature Communications* 2021;12:1–9.
- 62 Karachaliou N, Gonzalez-Cao M, Crespo G, *et al.* Interferon gamma, an important marker of response to immune checkpoint blockade in non-small cell lung cancer and melanoma patients. *Ther Adv Med Oncol* 2018;10:1758834017749748.
- 63 Villarino AV, Kanno Y, O'Shea JJ. Mechanisms and consequences of JAK-STAT signaling in the immune system. *Nat Immunol* 2017;18:374.
- 64 Leruste A, Tosello J, Ramos RN, *et al.* Clonally expanded T cells reveal immunogenicity of rhabdoid tumors. *Cancer Cell* 2019;36:597–612.
- 65 Mullen J, Kato S, Sicklick JK, *et al.* Targeting ARID1A mutations in cancer. *Cancer Treat Rev* 2021;100:102287.
- 66 Tsurusaki Y, Okamoto N, Ohashi H, *et al.* Mutations affecting components of the SWI/SNF complex cause Coffin-Siris syndrome. *Nat Genet* 2012;44:376–8.
- 67 Zou S, Li J, Zhou H, *et al.* Mutational landscape of intrahepatic cholangiocarcinoma. *Nat Commun* 2014;5:1.
- 68 Wiegand KC, Lee AF, Al-Agha OM, *et al.* Loss of BAF250a (ARID1A) is frequent in high-grade endometrial carcinomas. *J Pathol* 2011;224:328–33.
- 69 Cancer Genome Atlas Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 2012;487:330.
- 70 Wu R-C, Wang T-L, Shih I-M. The emerging roles of ARID1A in tumor suppression. *Cancer Biol Ther* 2014;15:655–64.
- 71 Li J, Wang W, Zhang Y, *et al.* Epigenetic driver mutations in ARID1A shape cancer immune phenotype and immunotherapy. *J Clin Invest* 2020;130:2712–26.
- 72 Park Y, Chui MH, Suryo Rahmanto Y, *et al.* Loss of ARID1A in tumor cells renders selective vulnerability to combined ionizing radiation and PARP inhibitor therapy. *Clin Cancer Res* 2019;25:5584–94.
- 73 Banerjee S, Stewart J, Porta N, *et al.* ATARI trial: ATR inhibitor in combination with olaparib in gynecological cancers with ARID1A loss or no loss (ENGOT/GYN1/NCRI). *International Journal of Gynecologic Cancer* 2021;31:1471–5.
- 74 Samartzis EP, Gutsche K, Dedes KJ, *et al.* Loss of ARID1A expression sensitizes cancer cells to PI3K- and AKT-inhibition. *Oncotarget* 2014;5:5295.
- 75 Benusiglio PR, Couvé S, Gilbert-Dussardier B, *et al.* A germline mutation in PBRM1 predisposes to renal cell carcinoma. *J Med Genet* 2015;52:426–30.
- 76 Briggs LG, Cone EB, Lee RJ, *et al.* Prognostic and predictive biomarkers for metastatic renal cell carcinoma. *J Cancer Metastasis Treat* 2021;7:46.
- 77 Moreira M, Pobel C, Epailard N, *et al.* Resistance to cancer immunotherapy in metastatic renal cell carcinoma. *Cancer Drug Resist* 2020;3:454–71.
- 78 Braun DA, Ishii Y, Walsh AM, *et al.* Clinical validation of PBRM1 alterations as a marker of immune checkpoint inhibitor response in renal cell carcinoma. *JAMA Oncol* 2019;5:1631–3.
- 79 Carril-Ajuria L, Santos M, Roldán-Romero JM, *et al.* Prognostic and predictive value of PBRM1 in clear cell renal cell carcinoma. *Cancers* 2019;12:16.
- 80 Argentiero A, Solimando AG, Krebs M, *et al.* Anti-angiogenesis and immunotherapy: novel paradigms to envision tailored approaches in renal cell carcinoma. *J Clin Med* 2020;9:1594.
- 81 Yang Q, Shen R, Xu H, *et al.* Comprehensive analyses of PBRM1 in multiple cancer types and its association with clinical response to immunotherapy and immune infiltrates. *Ann Transl Med* 2021;9:465.
- 82 Errichiello E, Mustafa N, Vetro A, *et al.* SMARCA4 inactivating mutations cause concomitant Coffin-Siris syndrome, microphthalmia and small-cell carcinoma of the ovary hypercalcaemic type. *J Pathol* 2017;243:9–15.
- 83 Nemes K, Bens S, Bourdeaut F. *Rhabdoid Tumor Predisposition Syndrome*. *GeneReviews® [Internet]*. Seattle: University of Washington, 2017.
- 84 Bourdeaut F, Lequin D, Brugières L, *et al.* Frequent hSNF5/INI1 germline mutations in patients with rhabdoid tumor. *Clin Cancer Res* 2011;17:31–8.
- 85 Mardinian K, Adashek JJ, Botta GP, *et al.* SMARCA4: Implications of an Altered Chromatin-Remodeling Gene for Cancer Development and Therapy. *Mol Cancer Ther* 2021;20:2341–51.
- 86 Dagogo-Jack I, Schrock AB, Kem M. Clinicopathologic characteristics of BRG1-Deficient non-small cell lung cancer. *Journal of Thoracic Oncology* 2020;15:766–76.
- 87 Sullivan LM, Folpe AL, Pawel BR, *et al.* Epithelioid sarcoma is associated with a high percentage of SMARCB1 deletions. *Mod Pathol* 2013;26:385–92.
- 88 Hohmann AF, Vakoc CR. A rationale to target the SWI/SNF complex for cancer therapy. *Trends Genet* 2014;30:356–63.
- 89 Italiano A. Targeting epigenetics in sarcomas through EZH2 inhibition. *J Hematol Oncol* 2020;13:33.
- 90 Kang N, Eccleston M, Clermont P-L, *et al.* EZH2 inhibition: a promising strategy to prevent cancer immune editing. *Epigenomics* 2020;12:1457–76.
- 91 Hoy SM. Tazemetostat: first approval. *Drugs* 2020;80:513–21.
- 92 Ngo C, Postel-Vinay S. Immunotherapy for SMARCB1-Deficient sarcomas: current evidence and future developments. *Biomedicines* 2022;10:650.
- 93 Chiappinelli KB, Strissel PL, Desrichard A, *et al.* Inhibiting DNA methylation causes an interferon response in cancer via dsRNA including endogenous retroviruses. *Cell* 2015;162:974–86.
- 94 Solovyov A, Vabret N, Arora KS, *et al.* Global cancer transcriptome quantifies repeat element polarization between immunotherapy responsive and T cell suppressive classes. *Cell Rep* 2018;23:512–21.
- 95 Henssen AG, Koche R, Zhuang J, *et al.* PGBD5 promotes site-specific oncogenic mutations in human tumors. *Nat Genet* 2017;49:1005–14.
- 96 Pawel BR. SMARCB1-deficient tumors of childhood: a practical guide. *Pediatr Dev Pathol* 2018;21:6–28.
- 97 Calderaro J, Masliah-Planchon J, Richer W. Balanced translocations disrupting SMARCB1 are hallmark recurrent genetic alterations in renal medullary carcinomas. *Eur Urol* 2016;69. doi:10.1016/j.eururo.2015.09.027. [Epub ahead of print: 22 09 2021].
- 98 Sodji Q, Klein K, Sravan K, *et al.* Predictive role of PD-L1 expression in the response of renal medullary carcinoma to PD-1 inhibition. *J Immunother Cancer* 2017;5:62.
- 99 Botta GP, Kato S, Patel H, *et al.* SWI/SNF complex alterations as a biomarker of immunotherapy efficacy in pancreatic cancer. *JCI Insight* 2021;6. doi:10.1172/jci.insight.150453. [Epub ahead of print: 22 09 2021].