POSTER PRESENTATION

Open Access

An image analysis algorithm based on the hue saturation density transformation, an important tool for melanoma immunotherapy research

Jeffrey A Dzubay^{1*}, Alan Jerusalmi¹, Rajiv Jesudason¹, Johannes Zimmermann², Martin Baatz², Zipei Feng³, Sachin Puri³, Tarsem Moudgil³, Carlo Bifulco³, Bernard Fox³

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Introduction

Melanoma is the most serious form of skin cancer, resulting in nearly 10,000 deaths per year in the US alone. Approximately 76,000 people are diagnosed with melanoma each year, and that number has been growing steadily over the last 30 years. Tragically, melanoma affects younger as well as older people. In fact, it is one of the most common forms of cancer in people younger than 30 [1]. A significant challenge impeding melanoma research is that the very cells that cause the cancer, the melanocytes, can confound measurements made using traditional immunohistochemical (IHC) techniques. This is because the brown melanin in the cells creates a high background for bright-field IHC stains, which are often brown or red.

Results

A digital bright-field image is made up of separate channels (red, green and blue), with individual intensities determined by a three color CCD acquisition camera. Different colors perceived by eye are a mixture of these three channels, and thus not directly separated by traditional image analysis software. For example, the dark brown melanin filled regions of the image will appear in all three of the channels with different intensities. Definiens' stain isolation algorithm makes use of the Hue-Saturation-Density (HSD) transformation, which is an adaptation of the Hue-Saturation-Intensity model where the optical density for each individual channel is used as opposed to intensity values [2]. We used this powerful technique to successfully separate the red stained CD3 and CD8 positive immune cells from a background of brown melanocytes in TMA

core samples from melanoma patients. Each cell type was isolated into a separate channel on the basis of hue. The resulting image is functionally like a fluorescent image, with monochromatic grey scale image for each color (Figure 1 and 2). This greatly simplified and increased the accuracy of the automated detection and quantification of the distinct cell types. We demonstrated rapid and accurate quantification of the immune response that would be virtually impossible using manual techniques.

Figure 1 Original high resolution RGB scan of a classically stained TMA core.

¹Definiens, Inc., Carlsbad, CA, USA

Full list of author information is available at the end of the article

Figure 2 Pseudo-fluorescent image resulting from the stain isolation algorithm using H.

Authors' details

¹Definiens, Inc., Carlsbad, CA, USA. ²Definiens AG, Munich, Germany. ³Earle A. Chiles Research Institute, Providence Cancer Center, Portland, USA.

Published: 6 November 2014

References

1. American Cancer Society: *Cancer Facts & Figures 2014*. Atlanta, Ga: American Cancer Society; 2014.
2. van der Laak J, Pahlplatz C, Hanselaar A, Wilde P: Hue-Saturation-Density (HSD) Model for Stain Recognition in Digital Images From Transmitted Light Microscopy. *Cytometry* 2000, **39**:275-284.

doi:10.1186/2051-1426-2-S3-P134

Cite this article as: Dzubay et al.: An image analysis algorithm based on the hue saturation density transformation, an important tool for melanoma immunotherapy research. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P134.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit