POSTER PRESENTATION

Open Access

Epigenetic regulation of CD1d-mediated antigen presentation in B cell lymphoma

Irina Tiper*, Tonya J Webb

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Tumors frequently alter antigen processing and presentation by major histocompatibility complex (MHC) proteins in order to evade recognition by the immune system. CD1d, a non-polymorphic MHC class I-like molecule, presents lipid antigens to Natural killer T (NKT) cells, which have potent anti-tumor effector functions. NKT cells are able to directly lyse malignant cells and induce anti-tumor responses by modulating other immune cells. Many hematologic malignancies express CD1d and the co-stimulatory proteins needed to induce anti-tumor responses by NKT cells, yet most tumors are poorly immunogenic. Here we sought to test the hypothesis that B cell lymphomas use epigenetic mechanisms to dysregulate CD1d-mediated antigen processing and presentation leading to a functional impairment in the ability of NKT cells to recognize tumors. To assess the functional outcomes of epigenetic modulation, murine and human B cell lymphomas were pretreated with HDAC inhibitors and then we assessed their ability to process and present antigen. Pretreatment with Trichostatin A (TSA) resulted in a dose-dependent increase in CD1d-mediated NKT cell activation by lymphoma cells without altering CD1d or co-stimulatory molecule cell surface expression. Similarly, pretreatment with TSA enhanced MHC class II mediated antigen presentation to CD4+ T cells. In contrast, treatment with the more selective HDACi, MC1568, resulted in an increase in CD1d-mediated NKT cell activation, but did not enhance antigen presentation by HLA-DR4. These data indicate that HDACi differentially modulate CD1d and MHC class II-mediated antigen presentation and suggests a role for multiple HDACs in regulating antigen processing and presentation. Overall, our studies demonstrate the efficacy of HDACi in restoring NKT cell mediated anti-tumor responses and may provide the basis for an NKT cell-based immunotherapeutic strategy that not only enhances

the immune response, but also increases the immunogenicity of the tumor itself.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P177

Cite this article as: Tiper and Webb: Epigenetic regulation of CD1d-mediated antigen presentation in B cell lymphoma. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P177.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

University of Maryland School of Medicine, Baltimore, MD, USA

© 2014 Tiper and Webb; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.