POSTER PRESENTATION

Open Access

Cross-talk between CD8+ T cells and natural killers: the role of mitochondrial Aa2+ transport

Roman Uzhachenko^{1*}, Shawn Goodwin¹, Menaka Thounaojam¹, William Hofmeister², Anil Shanker³

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Unraveling complex interactions between immune cells is a key to the development of new strategies for immunotherapy. In the present study, we investigated functional outcome of bidirectional interaction between activated CD8+T and naive natural killer (NK) cells. We found that phorbol 12-myristate 13-acetate (PMA)/Ionomycin (Io)-stimulated CD8+T cells form multiple intercellular contacts with naive NK lymphocytes. Co-culture of activated T cells with naive NK cells results in the selective down-regulation of CD25 molecule in T cells while elevating CD25 and CD69 expression on naive NK cells. Further, CD8+T and NK cells cross-regulate mitochondrial homeostasis including calcium transport. This effect is dependent on both cytokines and intercellular contacts, and partially involves natural killer group 2 member D (NKG2D) receptor activation. Data also suggest that activated CD8+T cells might directly transfer mitochondria and activation molecules such as CD25 and CD69 to naive NK cells. Alterations in phosphorylation status of multiple signaling proteins during CD8+T/NK interaction suggest a functional remodeling whereby NK cells shift activated CD8+T cells towards T central-memory (TCM) phenotype and activated CD8+T lymphocytes alter naive state of NK cells towards effector/regulatory phenotype. Inhibition of mitochondrial Ca²⁺ uptake (mCU) or Na⁺/Ca²⁺ exchanger (mNCE) with Ru360 and CGP37157 respectively mimicked observed alterations in CD8+ and NK cells upon their interaction. These data suggest a potential role of mitochondrial Ca²⁺ homeostasis in the acquisition of mixed activation/regulatory phenotype by NK cells during T cell-NK cell interaction. We believe that understanding the mechanisms of CD8+-NK interplay will help to develop new approaches for cellular immunotherapy.

Authors' details

¹Meharry Medical College, Nashville, TN, USA. ²University of TN Space Institute, Nashville, TN, USA. ³Meharry Medical College School of Medicine / Vanderbilt-Ingram Cancer Center, Nashville, TN, USA.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P189

Cite this article as: Uzhachenko et al.: Cross-talk between CD8+ T cells and natural killers: the role of mitochondrial Aa2+ transport. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P189.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

¹Meharry Medical College, Nashville, TN, USA
Full list of author information is available at the end of the article