

POSTER PRESENTATION

Open Access

TLR2/6 agonists and IFN-gamma treatment induces favorable immune cell recruiting signatures from melanoma associated with STAT1 and IL-32 signaling

Ileana S Mauldin^{1*}, Craig L Slingsluff¹, Ena Wang²

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Intralesional therapies offer promise to modulate immune signatures within melanoma and other cancers, either as monotherapy or as a component of combination immune therapy. We have shown that the TLR2/6 agonists (MALP-2 and FSL-1) and IFN γ induce human melanoma cells to synergistically produce T-cell attracting chemokine CXCL10. Despite the promise of inducing CXCL10 in the tumor microenvironment, IFN γ may induce negative immune regulatory processes, and TLR agonists have the potential to induce anti-apoptotic signaling in tumor cells. We hypothesized that synergy of IFN γ and TLR2/6 depends on STAT1 signaling, does not protect melanoma cells from apoptosis, and induces a more favorable immune signature than IFN γ alone.

To assess global effects of TLR2/6 agonist and IFN γ on melanoma, gene expression profiling of 4 melanoma cell lines was performed. They revealed that IFN γ treatment alone induced genes for CXCL9, and CXCL10 immune cell recruiting chemokines but also induced genes for negative immune regulators IDO and PD-L1 when compared to untreated cells. Comparison of TLR2/6+IFN γ stimulated melanoma cells to IFN γ stimulation alone showed induction of CXCL10, CXCL11, and C3 expression. Furthermore, genes encoding Treg-recruiting chemokines were not induced with TLR2/6 agonists+IFN γ ; instead, profiles that promote Th1 and Th17 cells were observed. Gene profiling also demonstrated that IL-32 and STAT1 were induced by TLR2/6 agonists+IFN γ treatment when compared to TLR2/6 agonists or IFN γ treatment alone; we hypothesize that they may be the mechanistic mediators of

the synergy between TLR2/6 agonists+IFN γ , since they have been shown to mediate this synergy for other TLR agonists.

Proliferation assays show that treatment with TLR2/6 agonists+IFN γ does not promote melanoma cell proliferation. Furthermore, viability assays demonstrate that TLR2/6 agonist+IFN γ treatment of melanoma does not hinder apoptosis. To address the specificity of TLR2/6 agonists we knocked down genes encoding for TLRs 1, 2 and 6 using siRNA. We find that TLR2/6 agonists may signal through TLR1, 2, and 6 suggesting that these putative TLR 2/6 agonists may also signal through TLR1/2 to mediate CXCL10 production in a wider range of melanoma tumors.

Collectively, our data suggest that TLR2/6 agonists induce favorable gene signatures which may promote immune cell infiltration of melanoma. Gene array analysis reveals that IL-32 and STAT1 may mediate the synergistic CXCL10 production observed from TLR2/6 agonist+IFN γ stimulated melanoma cells.

Authors' details

¹Department of Surgery, University of Virginia, Charlottesville, VA, USA.

²Transfusion Medicine, NIH, Bethesda, MD, USA.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P225

Cite this article as: Mauldin et al.: TLR2/6 agonists and IFN-gamma treatment induces favorable immune cell recruiting signatures from melanoma associated with STAT1 and IL-32 signaling. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P225.

¹Department of Surgery, University of Virginia, Charlottesville, VA, USA
Full list of author information is available at the end of the article