

POSTER PRESENTATION

Open Access

Defining the effects of age and gender on clinical immune response to cancer vaccination

Adriana Ramirez*, Nolan Wages, Mark E Smolkin, Craig L Slingluff

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Objectives

Cancer vaccines have promise as monotherapy or as part of combination immunotherapy regimens. Age and gender implications on immune response to cancer vaccinations have not previously been well described. There is uncertainty about including elderly patients as study participants in vaccine trials because of the decline in immune response with increasing age [1,2]. Compelling evidence has also demonstrated gender differences in immune response associated with injury and vaccination for infectious diseases [3,4]. We hypothesized younger age and female gender may be predictive of higher rates of immune response to a multi-peptide cancer vaccine. Data collected from three clinical trials evaluating T cell response to a polyvalent melanoma peptide vaccine provide an opportunity to assess the association of immune response with age and gender.

Methods

Patients with resected stage IIB-IV melanoma were enrolled in three clinical trials: Mel43, Mel44, and Mel48, in which they received 6 vaccinations with 12 Class I MHC restricted peptides from melanocytic differentiation antigens and cancer testis antigens with each of various adjuvants. T cell responses were detected by direct IFN-gamma ELISpot assay. Clinical data, including age and gender, were collected. The cumulative incidence in immune response over time was plotted for each variable. Chi-squared analysis was performed on the number of immune responders by age and gender.

Results

A total of 327 patients were evaluated for immune response measurable by Week 7. Immune responses were detected in 111/224 (49.5%) males and 49/103 (47.6%) females, $p = 0.74$. Immune responses were

detected in 130/249 (53%) in patients less than 64 years old, 30/78 (38.5%) older patients ($p = 0.026$ by cumulative incidence over time; $p = 0.034$ by Chi-squared analysis). Age greater than 64 accounted for a 14.5 % decrease in immune response.

Conclusion

While our study shows a measurable decline in immune response with increasing age, a significant percentage of elderly patients do develop immune responses to vaccination. Meanwhile, gender was not shown to affect immune response despite evidence suggesting an immunoprotective association with female gender [3]. This information should be considered in patient selection for cancer vaccines. In particular, elderly patients should be made aware of this difference prior to undergoing vaccination but should not be excluded on the basis of age.

Acknowledgements

Funding for Mel43(NCT00089193), Mel44(NCT00118274), and Mel48 (NCT00705640) was provided by R21 CA103528, R01 CA118386, R01 CA57653, and the UVA cancer center support grant (NIH/NCI P30 CA44579)

Published: 6 November 2014

References

1. Pawelec G, Goldeck D, Derhovanessian E: Inflammation, ageing and chronic disease. *Curr Opin Immunol* 2014, **21**:23-28.
2. Knöferl M, Angele M, Diodato M: Female sex hormones regulate macrophage function after trauma-hemorrhage and prevent increased death rate from subsequent sepsis. *Annals of Surgery* 2002, **235**:105-112.
3. Klein S: Sex influences immune responses to viruses, and efficacy of prophylaxis and treatments for viral diseases. *Bioessays* 2012, **34**:1050-1059.
4. Hewagama A, Patel D, Yarlagadda S: Stronger inflammatory/cytotoxic T-cell response in women identified by microarray analysis. *Genes and immunity* 2009, **10**:509-516.

doi:10.1186/2051-1426-2-S3-P59

Cite this article as: Ramirez et al.: Defining the effects of age and gender on clinical immune response to cancer vaccination. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P59.

University of Virginia, Charlottesville, United States



© 2014 Ramirez et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.