ORAL PRESENTATION

Open Access

Virus-specific CD8⁺ T cells infiltrate melanoma lesions and retain function despite high PD-1 expression

Dan A Erkes*, Toktam Mohgbeli, Christopher M Snyder

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015)

National Harbor, MD, USA. 4-8 November 2015

It is well known that CD8⁺ T cell infiltration of tumors is correlated with positive patient prognosis. Because of this, many viral vaccine vectors have been created to induce expansion and infiltration of tumor-specific CD8⁺ T cells. Cytomegalovirus (CMV) is a herpes virus that induces a persistent/latent infection for the life of the host. It is able to drive large populations of viral-specific T cells that are fully functional and infiltrate most tissues. We have established that acute infection with a recombinant murine CMV virus expressing an altered gp100 antigen (MCMV-gp100) caused expansion and tumor infiltration of gp100-specific CD8⁺ T cells, leading to a therapeutic delay in the growth of B16 melanomas in mice. Coincidentally, we observed that at least 20% of the CD8⁺ tumor infiltrating lymphocytes (TIL) were MCMV-specific. These virus-specific T cells infiltrated tumors independently of viral replication. Unexpectedly, T cell infiltration of tumors occurred even in latently infected animals, long after viral replication had been controlled. Together, these data suggest that virus-specific CD8⁺ T cells can become TIL, even long after the primary infection. We were curious about differences between tumor and virus-specific CD8⁺ TIL. As expected, after acute MCMV-gp100 infection, gp100-specific CD8⁺ TIL were PD-1^{hi} and markedly dysfunctional. Virus-specific CD8⁺ T cells were also PD-1^{hi} in the tumor after acute MCMV-gp100 infection, but these cells remained fully functional. PD-1 is an activation marker expressed on recently stimulated CD8⁺ T cells and both gp100-specific and virus-specific T cells expressed PD-1 after acute MCMV-gp100 infection. In the circulation of tumor-bearing mice or in the absence

of a tumor, both gp100-specific and virus-specific CD8⁺ T cells lost PD-1 expression over time. However, PD-1^{hi} cells that migrated into tumors retained expression of PD-1 regardless of specificity. Strikingly, introducing B16 tumors into MCMV-gp100 latently infected mice after PD-1 expression was lost, resulted in PD-1 up-regulation on and subsequent dysfunction of gp100-specific T cells that entered the tumor, but not virus-specific T cells found in the same tumors. Together, these data suggest that the tumor environment sustains PD-1 expression on recently activated CD8⁺ TIL independently of antigen in the tumor, but that the presence of antigen in the tumor drives subsequent CD8⁺ TIL dysfunction.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-O6

Cite this article as: Erkes et al.: Virus-specific CD8⁺ T cells infiltrate melanoma lesions and retain function despite high PD-1 expression. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):O6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

Thomas Jefferson University, Philadelphia, PA, USA

© 2015 Erkes et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.