

POSTER PRESENTATION

Open Access

Co-expression networks identify distinct immune infiltrates in hepatocellular carcinoma

Derek Y Chiang*, Alan Huang

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Background

The vast majority of hepatocellular carcinoma (HCC) arise in the context of chronic inflammation, especially with hepatitis B or hepatitis C viral infection. Several studies have identified prognostic immune biomarkers in HCC tumors and peritumoral regions. Recently, a Phase 1 trial of a PD-1 inhibitor has demonstrated efficacy in HCC. In order to characterize the diversity of immune microenvironments in HCC, we investigated co-expression networks of immune lineage-specific genes.

Results

We conducted a meta-analysis of gene expression data from over 500 HCC tumors and matched adjacent liver specimens. PD-L1 and PD-L2 had higher RNA levels in adjacent liver, compared with tumors. Tumoral expression of PD-L1 and PD-L2 were correlated with macrophage lineage genes. We identified 3 major co-expression network modules that corresponded with different immune cell sub-types: (1) An infiltrating T cell module was enriched for TCR activation, recruitment chemokines and elevated immune checkpoints. (2) A hepatic stellate cell module was associated with extracellular matrix remodeling, epithelial-to-mesenchymal transition and TGF-beta signaling. (3) A macrophage module had elevated macrophage lineage genes. By integrating these co-expression modules with HCC molecular sub-classes, the infiltrating T cell signature was enriched in the Hoshida S1 subclass¹ and Chiang interferon subclass², and was less prevalent among HCC with *CTNGB1* mutations.

Conclusion

Transcriptomic analyses revealed immune cell types and potential regulators in HCC. The joint profiling of

infiltrating immune sub-types and genetic alterations may guide the selection of combination therapies.

Published: 4 November 2015

References

1. Hoshida Y, et al: *Cancer Research* 2009, **69**:7385-7392.
2. Chiang DY, et al: *Cancer Research* 2008, **68**:6779-6788.

doi:10.1186/2051-1426-3-S2-P213

Cite this article as: Chiang and Huang: Co-expression networks identify distinct immune infiltrates in hepatocellular carcinoma. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P213.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Novartis Institutes for Biomedical Research, Cambridge, MA, USA



© 2015 Chiang and Huang This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.