

POSTER PRESENTATION

Open Access

# Inducible expression of antigen processing and antigen presentation molecules and cancer testis antigens in human prostate cancer

Erika Heninger\*, Timothy Krueger, Brian Johnson, Jamie Sperger, David Jarrard, Joshua Lang

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Defects in immune surveillance have been correlated with tumorigenesis and poor clinical outcomes in cancer patients. Given recent advances in vaccine therapies and checkpoint inhibition, therapeutic targeting of defective antigen processing may play a critical role in improving the broad utility of immunotherapies. Epigenetic alterations have been found to alter expression of genes involved in antigen processing and presentation. We evaluated the ability of epigenetic modifying agents to increase expression of antigen presentation machinery (APM) molecules *ex vivo* drug culture using primary human prostate tumor tissue biopsies. Prostate tissue biopsies were collected from patients with advanced, localized prostate cancer undergoing radical prostatectomy. These tumor biopsies sectioned into 200µM slices and cultured with the hypomethylating agent decitabine (5AZA) and/or histidine-deacetylase inhibitor panobinostat (LBH589). After 72 hours of treatment, tissue slices were harvested and subjected to qRT-PCR analysis. 5AZA alone or in combination with LBH589 have increased expression of MHC Class I molecules (HLA-ABC, B2M) and APM elements including TAP2, Tapasin and LMP7. Increased expression of MHC class I molecules was identified in human prostate cancer treated with 5AZA and the combination of 5AZA and LBH589. Significant inter-patient heterogeneity was also observed, suggesting other mechanisms by which prostate cancer downregulates MHC class I expression. We further identify increased expression of Cancer-Testis Antigens in these samples, indicating multiple mechanisms by which epigenetic modifying agents may improve immune recognition of human prostate cancer. These include NY-ESO, SSX2 and NYSAR35. These results

suggest novel therapeutic strategies that can be employed with other immune based therapies for men with localized and advanced prostate cancer.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P224

**Cite this article as:** Heninger et al.: Inducible expression of antigen processing and antigen presentation molecules and cancer testis antigens in human prostate cancer. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P224.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



University of Wisconsin, Madison, WI, USA



© 2015 Heninger et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.