POSTER PRESENTATION

Open Access

Human cancer cells respond to cytosolic nucleic acids via enhanced production of interferon- β and apoptosis

Shan Zhu^{1*}, Yuan Qiao¹, Jing Wu¹, Guoxia Zang¹, Yong-Jun Liu², Jingtao Chen³

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

The innate immune system utilizes pattern-recognition receptors (PRRs) to detect the invasion of pathogens and initiate host antimicrobial responses. Recently, it was reported that human cancer cells also expressed PRRs, and responded to cytosolic nucleic acids via the production of type I interferon and apoptosis in susceptible cells. However, the definite nucleic acids senses in different cancer cells remains unclear. In this study, we investigated the effects of nucleic acids in eight types of cancer cells, including pancreatic cancer, glioma, breast cancer, lung cancer, hepatoma, gastric cancer, colorectal cancer, and cervical cancer. We found that IFN- β secretion was increased after poly(I:C) and poly(dA:dT) stimulation. In order to understand the mechanism, we established the role of TLR3 and RIG-I/MDA-5 signaling in pancreatic cancer cell line PANC-1. poly(I:C) and poly(dA:dT) upregulated the expression of PKR, TLR3, RIG-I, MDA5 and LGP2. Knockdown experiments confirmed PKR, TLR3, TRIF, RIG-I, LGP2, and IPS-1 involved in response to cytosolic poly(I:C) and poly(dA:dT). In addition, poly(I:C) and poly(dA:dT) induced apoptosis via caspase-8 and caspase-9 in pancreatic cancer cells. In summary, cytosolic poly(I:C) and poly(dA:dT) induce IFN- β secretion via TLR3-TRIF and RIG-I/LGP2-IPS-1 signaling pathway and apoptosis in a caspase-dependent manner at cancer cells. This study may provide a novel way for drug and vaccine design on cancer immunotherapy.

Authors' details

¹Institute of Translational Medicine, the First Hospital, Jilin University, Changchun, China. ²MedImmune, Gaithersburg, Changchun, China. ³Institute

of Translational Medicine, the First Hospital, Jilin University, Changchun, Jilin, China.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P250

Cite this article as: Zhu et al.: Human cancer cells respond to cytosolic nucleic acids via enhanced production of interferon- β and apoptosis. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P250.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

¹Institute of Translational Medicine, the First Hospital, Jilin University, Changchun, China

Full list of author information is available at the end of the article

© 2015 Zhu et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.