POSTER PRESENTATION

Open Access

Identification and isolation of high avidity tumor-specific CD8 T cells by experimentally assessing pMHC-TCR binding parameters with soluble pMHC complexes

Julien Schmidt^{1*}, Michael Hebeisen², Philippe Guillaume³, Morgane Magnin¹, Nathalie Rufer², Immanuel Luescher¹

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Adoptive transfer of high avidity tumor-specific CD8 T cells is a promising treatment for late stage cancers. Because most tumor antigens are self-antigens, tumor-specific T cells typically express low affinity T cell antigen receptors (TCR) due to thymic negative selection, which accounts mostly for their poor tumoricidal activity. To identify rare, most tumor-reactive T cells we developed a robust and accurate screening assay. The functional avidity of T cell responses is related to pMHC-TCR binding parameters, with an assumed optimal range of pMHC-TCR affinity for the most potent interaction. We recently described the development of NTAmers[1], reversible pMHC multimers built on switchable Ni²⁺-NTA-His-tag complexes, that allow accurate measurement of monomeric pMHC-TCR dissociation kinetics on living tumor-specific CD8 T cells by flow cytometry. We found a good correlation between off-rates and cellular responses on cloned CD8 T cells from cancer patients[2]. Moreover we developed pMHC dimers allowing measurement of pMHC-TCR association and dissociation kinetics on cancer-specific T cells. Combined measurement of pMHC-TCR on and off rates allowed accurate identification and isolation of functional high avidity tumor-specific CD8 T cells. In vivo experiments confirmed that high avidity CD8 T cells isolated from a cancer patient conferred superior tumor control. This novel technology has further allowed us to describe the functional involvement of the CD8 co-receptor in this interaction. Using CD8-binding deficient pMHC complexes, we observed that CD8

increases on CD8 T cells the off-rate by 3-4 fold but has no discernable effect on the on-rate.

In conclusion, NTAmers and pMHC dimers enable precise measurement of pMHC-TCR binding parameters under physiological conditions and subsequent isolation of live functionally active tumor-specific CD8 T cells. Translation of these techniques to the single cell level for high-throughput screening and isolation of rare and highly functional tumor-specific CD8 T cells directly from cancer patients is in progress.

Authors' details

¹Ludwig Center for Cancer Research, Epalinges, Switzerland. ²Department of Oncology, Epalinges, Switzerland. ³TCMtrix Ltd, Epalinges, Switzerland.

Published: 4 November 2015

References

- Schmidt J, et al: Reversible major histocompatibility complex I-peptide multimers containing Ni(2+)-nitrilotriacetic acid peptides and histidine tags improve analysis and sorting of CD8(+) T cells. *J Biol Chem* 2011, **286**(48):41723-35.
- Hebeisen M, et al: Identification of Rare High-Avidity, Tumor-Reactive CD8+ T Cells by Monomeric TCR-Ligand Off-Rates Measurements on Living Cells. *Cancer Res* 2015, **75**(10):1983-91.

doi:10.1186/2051-1426-3-S2-P263

Cite this article as: Schmidt et al: Identification and isolation of high avidity tumor-specific CD8 T cells by experimentally assessing pMHC-TCR binding parameters with soluble pMHC complexes. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P263.

¹Ludwig Center for Cancer Research, Epalinges, Switzerland
Full list of author information is available at the end of the article