POSTER PRESENTATION

Open Access

CXCR5⁺CD8⁺ T cells are localized in B cell follicles and germinal centers and exhibit regulatory and anti-tumor function

Fuliang Chu*, Sattva S Neelapu

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Germinal centers (GC) are specialized anatomic sites where B cells undergo somatic hypermutation, clonal expansion, and affinity-based selection. Stringent regulation of GC reactions is critical for homeostatic B cell development, as well as T-dependent humoral immunity against self and foreign antigens. Follicular helper T cells (T_{fh}, CXCR5^{hi}PD1^{hi}Bcl-6⁺CD4⁺) that reside in GC serve as specialized B helper T cells and provide survival and selection signals to GC B cells. Despite recent understandings of T_{fh} function in autoimmunity, immunodeficiency, and B-cell lymphoma, mechanisms that regulate T_{fh} development and function remain limited.

In the current study, we determined the presence of CD45RA⁻CXCR5⁺CD8⁺ T cells in human peripheral blood, tonsillar tissues, and follicular lymphoma (FL) tumor samples by FACS and confocal microscopy. We found that CXCR5⁺CD8⁺ T cells were present in high numbers and localized to GCs and T cell zones in the tonsillar tissues and FL, but present in low numbers in the peripheral blood. Surface marker analysis suggested that these cells displayed an activated status. Importantly, FL tumor samples had significantly more CXCR5⁺CD8⁺ T cells than tonsillar tissues. Further analysis using intracellular cytokine staining showed that CXCR5⁺CD8⁺ T cells produced higher levels of IFN- γ and TNF- α compared to CXCR5⁻CD8⁺ subset.

Next, we performed T-B cell co-culture experiments to evaluate whether CXCR5⁺CD8⁺ T cells could regulate T_{fh} function in humans. FACS-sorted T_{fh} cells were co-cultured with naïve or memory B cells in the presence or absence of CXCR5⁺CD8⁺ T cells. These experiments indicated that CXCR5⁺CD8⁺ T cells suppressed T_{fh}

function, as demonstrated by reduced differentiation of naïve and/or memory B cells into plasmablasts cells (CD19^{int}CD38⁺). To test CXCR5⁺CD8⁺ T cell function in vivo, we employed the EG7 lymphoma model in TCR^{-/-} mice. Briefly, ovalbumin-primed OT1-specific CXCR5⁺CD8⁺ T cells were adoptively transferred into EG7 tumor-bearing TCR^{-/-} mice, and tumor size and survival rate were determined.

In summary, our data suggest that CXCR5⁺CD8⁺ T cells may have multiple roles, as they not only inhibited T_{fh} function, but also exhibited strong anti-tumor immunity.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P321

Cite this article as: Chu and Neelapu: CXCR5⁺CD8⁺ T cells are localized in B cell follicles and germinal centers and exhibit regulatory and anti-tumor function. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P321.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

The University of Texas MD Anderson Cancer Center, Houston, TX, USA

© 2015 Chu and Neelapu This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.