

POSTER PRESENTATION

Open Access

# Mcl-1 expression influences CD8+ anti-tumor immunity.

Lukas W Pfannenstiel\*, Brian R Gastman

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

The recent clinical successes of immunotherapies such as anti-PD-1 and anti-CTLA-4 antibody blockade have re-invigorated the idea that the immune system has the potential to be a powerful tool to destroy tumors. However despite ample evidence that these therapies are able to enhance the efficacy of anti-tumor CD8+ T cell responses, the association of pro-survival molecules such as Mcl-1 with this process remains poorly understood. We have previously shown that short-term *in vitro* co-culture of human T cells with human-derived tumor cell lines of various origins can induce the gain of senescence-like features CD8+ T cells, particularly the loss of CD27/CD28 expression and the gain of a potent suppressive function in *in vitro* suppression assays. In subsequent studies, we found that IL-7 could protect T cells from the development of dysfunction/suppression, and that this process is highly dependent on the expression of the pro-survival protein Mcl-1. In the current study we sought to determine whether the beneficial effects of IL-7 could also be shown *in vivo* in a mouse model of head and neck squamous cell carcinoma (HNSCC). We show that the use of exogenous IL-7 treatment results in a delay in tumor growth, and that the combination of IL-7 with other immunotherapies, particularly anti-PD-1 antibody blockade, results in a synergistic response which is better than either therapy alone. Further, when combined with an adoptive transfer of TCR transgenic T cells, IL-7 treatment results in enhanced delay of tumor growth and greater numbers of tumor-specific T cells in the tumor microenvironment. Similar to our previous *in vitro* studies, we further find that expression of Mcl-1 can have a significant effect on anti-tumor CD8+ T cells responses. Knock-down of Mcl-1 expression significantly abrogates the function of tumor-specific, adoptive-transferred T cells while the

enhancement of Mcl-1 expression also enhances CD8+ function and significantly delays tumor growth. These data indicate that one benefit of existing immunotherapy strategies like IL-7 could be in the enhancement of expression of pro-survival factors like Mcl-1, and those future treatment strategies which specifically target their expression could improve anti-tumor immune responses.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P323

Cite this article as: Pfannenstiel and Gastman: Mcl-1 expression influences CD8+ anti-tumor immunity.. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P323.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



Cleveland Clinic, Cleveland, OH, USA



© 2015 Pfannenstiel and Gastman This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.