POSTER PRESENTATION

Open Access

Lung cancer recruitment of CD14⁺ cells results in an immunosuppressive phenotype and improved tumor survival under stress conditions

Erin Schenk*, Allan Dietz

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

We have previously shown that a higher burden of CD14⁺HLA-DR^{lo/neg} cells in the peripheral blood of cancer patients is associated with systemic immune suppression and a poor outcome. We have also reported that CD14⁺ cells isolated from healthy donors are converted to the immunosuppressive CD14⁺HLA-DR^{lo/neg} phenotype when cocultured with tumor cell lines or primary tumor cultures. Therefore, one potential mechanism of CD14⁺HLA-DR^{lo/neg} generation in cancer patients is the recruitment of CD14⁺ cells to the tumor microenvironment. In this study, we tested the ability of multiple human lung cancer cell lines to recruit CD14⁺ cells, convert normal donor CD14⁺ cells to CD14⁺HLA-DR^{lo/neg}, and the impact on tumor survival under stress conditions in the presence of CD14⁺HLA-DR^{lo/neg} cells. Monocyte recruitment by cell line supernatants was studied using freshly isolated CD14⁺ cells and measured in real time with live cell imaging. Monocytes from 10/10 unique healthy donors migrated towards H23 supernatant in a dose dependent fashion. Supernatants from other human lung cancer cell lines including H889, H1155, H1581, and H1975 also recruited healthy donor monocytes 6 to 9 fold greater than culture media alone. The CD14⁺ phenotype resulting from interaction with lung cancer cell lines was studied in a 2 day coculture system. Coculture of H1581 with monocytes from 27 of 29 unique healthy donors resulted in a downregulation of monocyte cell surface expression of HLA-DR by 48.7% ± 22.1%, a phenotype known to mediate both local and systemic immune suppression. CD14⁺ HLA-DR downregulation after coculture was observed with H23 (17/18 unique healthy donors, 49.7% ± 22.1% HLA-DR reduction) and H889 (5/6 unique healthy donors, 47.3% ± 16.1%

HLA-DR reduction). Importantly, CD14⁺ cells promoted lung tumor cell line survival under stress conditions. After 6 days of low serum growth conditions, the survival and proliferation of H23 and H1581 was approximately 2 fold greater when cocultured with CD14⁺ cells than without monocytes. H889 continuously exposed to cisplatin in culture had significantly improved survival when co-cultured with monocytes for 2 days prior to cisplatin exposure. In conclusion, we have demonstrated multiple lung cancer cell lines recruit and transform monocytes to the HLA-DR^{lo/neg} phenotype. These data suggest monocyte cancer cross-talk results in improved tumor survival when exposed to the likely *in vivo* conditions of a nutrient poor environment or chemotherapy independent of local immune suppression and angiogenesis.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P416

Cite this article as: Schenk and Dietz: Lung cancer recruitment of CD14⁺ cells results in an immunosuppressive phenotype and improved tumor survival under stress conditions. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P416.

Mayo Clinic, Rochester, MN, USA

© 2015 Schenk and Dietz This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.