RESEARCH ARTICLE

Open Access

Tumor-released autophagosomes induces CD4⁺ T cell-mediated immunosuppression via a TLR2–IL-6 cascade

Yong-Qiang Chen^{1†}, Peng-Cheng Li^{1†}, Ning Pan^{1†}, Rong Gao¹, Zhi-Fa Wen¹, Tian-Yu Zhang¹, Fang Huang¹, Fang-Yuan Wu², Xi-Long Ou³, Jin-Ping Zhang⁵, Xue-Jun Zhu^{1,7}, Hong-Ming Hu^{1,4}, Kang Chen^{6,7}, Yun-Lang Cai^{2*} and Li-Xin Wang^{1*}

Abstract

Background: CD4⁺ T cells are critical effectors of anti-tumor immunity, but how tumor cells influence CD4⁺ T cell effector function is not fully understood. Tumor cell-released autophagosomes (TRAPs) are being recognized as critical modulators of host anti-tumor immunity during tumor progression. Here, we explored the mechanistic aspects of TRAPs in the modulation of CD4⁺ T cells in the tumor microenvironment.

Methods: TRAPs isolated from tumor cell lines and pleural effusions or ascites of cancer patients were incubated with CD4⁺ T cells to examine the function and mechanism of TRAPs in CD4⁺ T cell differentiation and function. TRAPs-elicited CD4⁺ T cells were tested for their suppression of effector T cell function, induction of regulatory B cells, and promotion of tumorigenesis and metastasis in a mouse model.

Results: Heat shock protein 90α (HSP90α) on the surface of TRAPs from malignant effusions of cancer patients and tumor cell lines stimulated CD4⁺ T cell production of IL-6 via a TLR2–MyD88–NF-κB signal cascade. TRAPs-induced autocrine IL-6 further promoted CD4⁺ T cells secretion of IL-10 and IL-21 via STAT3. Notably, TRAPs-elicited CD4⁺ T cells inhibited CD4⁺ and CD8⁺ effector T cell function in an IL-6- and IL-10-dependent manner and induced IL-10-producing regulatory B cells (Bregs) via IL-6, IL-10 and IL-21, thereby promoting tumor growth and metastasis. Consistently, inhibition of tumor autophagosome formation or IL-6 secretion by CD4⁺ T cells markedly retarded tumor growth. Furthermore, B cell or CD4⁺ T cell depletion impeded tumor growth by increasing effector T cell function.

Conclusions: HSP90α on the surface of TRAPs programs the immunosuppressive functions of CD4⁺ T cells to promote tumor growth and metastasis. TRAPs or their membrane-bound HSP90α represent important therapeutic targets to reverse cancer-associated immunosuppression and improve immunotherapy.

Keywords: Extracellular vesicles (EVs), Tumor-released autophagosome (TRAP), CD4⁺ T cell, Regulatory B cell, IL-6, Heat shock protein 90α (HSP90α)

* Correspondence: ylseu63@163.com; lxwang@seu.edu.cn

[†]Yong-Qiang Chen, Peng-Cheng Li and Ning Pan contributed equally to this work.

²Department of Obstetrics and Gynecology, Zhongda Hospital, Medical School of Southeast University, 87 Dingjiaqiao Road, Nanjing 210009, China

¹Department of Microbiology and Immunology, Medical School of Southeast University, 87 Dingjiaqiao Road, Nanjing 210009, China

Full list of author information is available at the end of the article

© The Author(s). 2019 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

CD4⁺ T cells play a critical role in modulating both innate and adaptive anti-tumor immune responses. Research over the past two decades has revealed that CD4⁺ effector T cells, especially IFN- γ -producing T helper 1 (Th1) cells, can exhibit anti-tumor activity [1]. However, other subtypes of tumor-infiltrating CD4⁺ T cells may play a pro-tumorigenic role in the tumor microenvironments via the secretion of inflammatory or regulatory cytokines, such as interleukin (IL)-6, IL-10, IL-17, IL-21, and transforming growth factor (TGF)- β , as the abundance of such CD4⁺ T cells has been associated with a poor clinical outcome of various types of cancer [1–4]. It has also become clear that many tumor-derived molecules or extracellular vesicles likely influence the differentiation of CD4⁺ T cells [5, 6]. However, the precise mechanisms underlying CD4⁺ T cell differentiation and functions in the tumor microenvironment are not completely understood.

Extracellular vesicles (EVs) have emerged as a new mode of intercellular communication by functioning as the carriers of bioactive molecules to influence the extracellular environment and the immune system [6–8]. Recent evidences indicate that secretory autophagy, in contrast to canonical autophagy, is an alternative non-degradative mechanism for cellular trafficking and unconventional secretion of proteins and small molecules [9], such as IL-1 β [10], high mobility group box 1 (HMGB1) [11], adenosine triphosphate (ATP) [12], TGF- β [13], and lysozyme [14]. More importantly, secretory autophagosomes carrying cytoplasmic cargoes, including tumor-specific antigens or viruses, fail to fuse with lysosomes and instead are released into the extracellular environment by the cells under stress [15, 16].

We have previously found extracellular secretory autophagosomes from the supernatant of tumor cells or malignant effusions and ascites of cancer patients [17, 18], and have termed such tumor-released autophagosomes TRAPs. We confirmed that TRAPs can be taken up by phagocytes such as neutrophils and macrophages, as well as B cells, and endow them with immunosuppressive activities [18–20]. These observations highlight that TRAPs are part of an elaborate network of tumor-derived vesicles that can reroute the immune response towards a cancer-promoting direction and should be targeted to improve cancer therapy. However, the mechanistic aspects of TRAPs in the modulation of immune cell function, especially the key anti-tumor effector cell, CD4⁺ T cell, in the tumor microenvironment and during tumor progression are unclear.

Here, we demonstrate that TRAPs could educate CD4⁺ T cells to produce IL-6 that functions in an auto-crine manner to promote the production of IL-10 and IL-21. TRAPs-elicited CD4⁺ T cells (T_{TRAP}) directly

inhibit the anti-tumor IFN- γ response of CD4⁺ T and CD8⁺ T cells and also induce IL-10⁺ Bregs, which creates a favorable environment to facilitate tumor growth and metastasis. Mechanistic studies revealed that membrane-bound HSP90 α on intact TRAPs is crucial for inducing IL-6 production in CD4⁺ T cells via a TLR2–MyD88–NF- κ B signal cascade. Moreover, auto-crine IL-6 further stimulates CD4⁺ T cells to produce IL-10 and IL-21 via STAT3. Our study unveils novel cellular and molecular mechanisms of tumor-derived extracellular vesicles in regulating CD4⁺ effector T cell function and pinpoint TRAPs as a therapeutic target for cancer immunotherapy.

Materials and methods

Human subjects

Malignant pleural effusions and ascites were collected from cancer patients pathologically diagnosed with multiple cancer types. The clinicopathological characteristics of the enrolled patients are presented in Additional file 1: Table S1. The study was approved by the Ethics Committee for Human Studies of Southeast University (protocol 2016ZDKYSB112).

Mice

C57BL/6 female mice were purchased from the Comparative Medicine Center of Yangzhou University. *Tlr4*^{−/−}, *Tlr2*^{−/−}, *Myd88*^{−/−} and OT-I mice were purchased from the Nanjing Biomedical Research Institute of Nanjing University (Nanjing, China). *Il6*^{−/−} mice were gifts from Dr. Jinping Zhang (Institutes of Biology and Medical Sciences, Soochow University, Suzhou, China). Mice were maintained in the barrier facility at Southeast University. All animal experiments were approved by the Institutional Animal Care and Use Committee of Southeast University.

Cell culture

The murine hepatic carcinoma line Hepa1–6, melanoma line B16F10, Lewis lung carcinoma line LLC, lymphoma line EL4, and the human melanoma line A375, hepatic carcinoma line HepG2 and breast carcinoma line MDA-MB-231 were cultured in complete RPMI-1640 medium with 10% FBS (Gibco), 100 U/ml penicillin and 0.1 mg/ml streptomycin at 37 °C in a 5% CO₂ incubator. *Becn1* knockdown (*Becn1* KD) and negative control B16F10 cells (*Becn1* NC) were established by using lentivirus expressing *Becn1*-targeting (5′-GCGGGAGUAUAGUGAGUUUTT-3′) and scrambled (5′-TTCTCCGAA CGTGTACGTA-3′) shRNA (Hanbio Biotechnology, Shanghai, China), respectively.

Chemicals

The inhibitors PD98059, SP600125, SB203580, LY294002, BAY11-7082, and Stattic were purchased from MCE (Shanghai, China). Recombinant murine IL-2 and IL-12 were purchased from PeproTech (Rocky Hill, USA). CFSE were purchased from Invitrogen/Thermo Fisher Scientific. IL-6, IL-10 and IL-21 neutralizing antibodies were purchased from R&D Systems. Lymphocyte separation media were purchased from MultiSciences (Hangzhou, China). All other reagents were obtained from Sigma-Aldrich (St. Louis, MO).

TRAPs purification and characterization

Tumor cells were seeded in a T175 flask in complete RPMI-1640 culture medium supplemented with 10% heat-inactivated FBS (Gibco), 100 U/ml penicillin, and 0.1 mg/ml streptomycin and incubated for 3–4 days at 37 °C, 5% CO₂ until 100% confluency was reached. Tumor cell culture supernatants were collected for TRAPs isolation as described previously [18, 20]. Briefly, supernatants were centrifuged at 2000 rpm for 10 min to remove whole cells and debris. The supernatants were further centrifuged at 12,000 g for 30 min to harvest the TRAPs-containing pellet. The TRAPs-containing pellet was washed three times with PBS and isolated with magnetic beads (Miltenyi Biotec) combined with LC3b antibody (Cell Signaling Technology) for TRAPs. The purity of TRAPs was analyzed by flow cytometry and western blot. The size of TRAPs was determined by dynamic light scattering using a Malvern Instrument.

Primary cell isolation

Mouse splenic B cells (Invitrogen, 11422D), CD4⁺ T cells (Invitrogen, 11415D), CD8⁺ T cells (Invitrogen, 11417D) and human peripheral blood CD4⁺ T cells (Miltenyi Biotec, 130-045-101) were purified by magnetic-activated cell sorting (MACS) following the manufacturer's instructions. After the MACS, the purity of T and B cells were > 95% as assessed by flow cytometry.

Flow cytometry

Purified CD4⁺ T or CD8⁺ T cells were cultured in a 24-well plate pre-coated with 2 µg/ml anti-CD3 (BD Biosciences, 550,275) and 2 µg/ml anti-CD28 mAb (BD Biosciences, 553,294) in the presence of 50 U/ml IL-2 (PeproTech), purified TRAPs and 30% culture supernatants from CD4⁺ T cells or B cells. In some cases, culture supernatants from CD4⁺ T cells or B cells were pretreated with neutralizing mAbs against IL-6, IL-10, or IL-21 for 1 h at 4 °C and subsequently exposed to T cells or B cells. Three days later, IFN-γ⁺ CD4⁺ T, IFN-γ⁺ CD8⁺ T or IL-10⁺ B cells were evaluated by flow cytometry. For intracellular staining, the cells were stimulated with the ovalbumin (OVA) protein or anti-CD3 and

anti-CD28 mAbs at 37 °C for 24 or 72 h. Leukocyte activation cocktail and GolgiPlug (BD Biosciences) were added to the culture 5 h prior to flow cytometric analysis. Subsequently, the cells were stained with antibodies specific to the various surface molecules, fixed and permeabilized with a Fixation/Permeabilization Kit (BD Biosciences), and finally stained with antibodies against the various intracellular molecules. To detect Bcl-6 and Foxp3, the cells were fixed and permeabilized using a Transcription Factor Buffer Set (BD Biosciences). Data were acquired using a FACS Calibur analyzer (BD Biosciences) and analyzed by FlowJo. The gates were set according to the staining by isotype-matched control antibodies of the respective cells. The fluorochrome-conjugated Abs used are listed in Additional file 1: Table S2.

Quantitative real-time PCR

Total RNA from CD4⁺ T cells was isolated with TRIzol reagent (Invitrogen) and reverse-transcribed using 5 × PrimeScriptRT Master Mix (Takara), following the manufacturer's instructions. The specific primers used to amplify the genes are listed in Additional file 1: Table S3. The PCR was performed in triplicate using Fast Start Universal SYBR Green Master (ROX) (Roche Life Science) in a StepOne Real-Time PCR System (Thermo Fisher Scientific). GAPDH was used as an internal standard.

Elisa

Cytokines in the sera or cell culture supernatants were quantified using ELISA kits according to the manufacturer's protocol. ELISA sets were purchased from eBioscience (IL-6 and IL-10) and R&D Systems (IL-21).

Western blot

The proteins samples were extracted from CD4⁺ T cells with RIPA lysis buffer. They were separated and transferred as previously described [21]. The membranes were blocked with 5% BSA in TBST for 1 h and separately incubated with the primary antibodies overnight at 4 °C. After washing with TBST buffer, the membranes were incubated with horseradish peroxidase-conjugated secondary antibodies for 1 h. The results were visualized by enhanced chemiluminescence according to the manufacturer's protocol. The primary antibodies used are listed in Additional file 1: Table S4.

Animal models

Wild type C57BL/6 mice or *Il6*^{-/-} C57BL/6 mice were subcutaneously inoculated with B16F10, B16F10 *Becn1* NC or B16F10 *Becn1* KD cells (2 × 10⁵ cells/mouse). Tumor growth was measured using a caliper. On day 21, draining lymph nodes (dLN), spleens or tumor tissues

were harvested from tumor-free or tumor-bearing mice. The frequencies of IL-10⁺ CD4⁺ T cells, IL-21⁺ CD4⁺ T cells, or IL-10⁺ B cells were evaluated by flow cytometry after ex vivo stimulation with the leukocyte activation cocktail and GolgiPlug (BD Biosciences) for 5 h. In the subcutaneous tumor model, B16F10 tumor cells (2×10^5 cells/mouse) and CD4⁺ T cells treated with TRAPs, or B cells treated with the indicated culture conditions (2×10^6 cells/mouse) were subcutaneously injected into the right flank of C57BL/6 mice. Subcutaneous tumor growth was monitored and measured using vernier calipers. In the tumor metastasis model, B16F10 tumor cells (5×10^5 cells/mouse) were intravenously injected into C57BL/6 mice and TRAPs-treated or untreated CD4⁺ T cells or B cells (5×10^6 cells/mouse) treated with the indicated culture conditions were injected every other day for 3 times. Three weeks later, mice were sacrificed, and the tumor nodules in the lungs were examined. To evaluate the role of CD4⁺ T cells and B cells treated with the indicated culture conditions in OVA-loaded DC-mediated specific immune response, C57BL/6 mice were adoptively transferred with OT-I splenocytes (1×10^7 cells/mouse) on day 0 and vaccinated with OVA-loaded DCs (1×10^6 cells/mouse) on days 1, 4, and 7. After intravenous administration of CD4⁺ T cells and B cells on days 2, 5, and 8, mice from each group were sacrificed on day 14 and the frequency and number of CD8⁺Vβ5.1⁺ T cells were evaluated by flow cytometry. The frequency of IFN-γ⁺ CD4⁺ and CD8⁺ T cells in the spleens was determined by intracellular cytokine staining after ex vivo stimulation with the OVA protein for 24 h.

T and B cell depletion

C57BL/6 mice ($n = 5$ /group) were inoculated subcutaneously in the flank with 1×10^6 *Becn1*-NC or *Becn1*-KD B16F10 cells. On day 9, the tumor-bearing mice were subsequently depleted of either CD4⁺ T cells, CD8⁺ T cells or CD20⁺ B cells by intravenous administration of 250 μg/mouse of anti-mouse CD4 (clone GK1.5, BioX-Cell) or anti-mouse CD8 (clone 2.43, BioXCell) twice weekly throughout the course of tumor growth, or 250 μg of anti-mouse CD20 (clone SA271G2, BioLegend), respectively. Control mice were treated similarly but with isotype-matched control antibodies. Depletion was confirmed by staining of peripheral blood cells with anti-mouse CD4 (RM4-5, BD Pharmingen), anti-mouse CD8 (clone 53-6.7, BioLegend), or anti-mouse CD19 (clone 6D5, BioLegend).

Statistical analysis

Data were derived from at least 3 independent experiments and analyzed using GraphPad Prism 5.0 software.

Multiple group comparisons were performed by one-way ANOVA and the Tukey-Kramer multiple test. Comparisons between 2 groups were performed using unpaired Student's t-test or Mann-Whitney U test. $P < 0.05$ was considered significant.

Results

TRAPs induce CD4⁺ T cells to produce IL-6, IL-10, and IL-21

To determine whether TRAPs impact CD4⁺ T cell function, we first isolated TRAPs from the culture supernatants of mouse B16F10 melanoma cells [18, 19]. The TRAPs preparation specifically contained the mature autophagosome marker LC-3II (Fig. 1a, b) and exhibited an average size of 436.3 nm, which was distinct from isolated exosomes that had an average size of 85.6 nm (Fig. 1c) and expressed the exosome markers CD63 and TSG101 (Fig. 1d). Treatment of mouse splenic CD4⁺ T cells with TRAPs during activation by anti-CD3 and anti-CD28 resulted in the induction of the transcripts encoding *Il6*, *Il21*, *Il10*, and *Il17*, but not *Il1b*, *Il2*, *Il4*, *Il9*, *Tnf*, *Ifng*, *Foxp3* or *Tgfb1* (Additional file 2: Figure S1a). Consistently, the frequency of IL-6⁺, IL-10⁺ or IL-21⁺ CD4⁺ T cells and the secretion of IL-6, IL-10 or IL-21 by CD4⁺ T cells were increased by TRAPs treatment (Fig. 1e, f). TRAPs-induced IL-21⁺ CD4⁺ T cells expressed elevated levels of the follicular helper T cell (T_{fh})-associated molecules CXCR5 and Bcl-6 (Additional file 2: Figure S1b, c). In contrast, TRAPs reduced the frequency of IFN-γ⁺ CD4⁺ T cells (Fig. 1e) and suppressed IL-12-mediated induction of IFN-γ⁺ Th1 cells (Additional file 2: Figure S1d). Depletion of TRAPs from the culture media via ultracentrifugation (Additional file 2: Figure S1e) resulted in a significant reduction of IL-6, IL-10 and IL-21 production by CD4⁺ T cells (Fig. 1g). Intriguingly, we also found that LC3B⁺ EVs (TRAPs) purified from B16F10 culture supernatant were more potent than LC3B⁻ EVs and exosomes in upregulating IL-6 expression, suggesting that LC3B⁺ EVs (TRAPs) are the dominant large EVs that instruct CD4⁺ T cells (Additional file 2: Figure S2a-d). In order to ascertain the role of TRAPs in inducing IL-6, IL-10 and IL-21 production by CD4⁺ T cells in vivo, normal saline (NS) or TRAPs were administered intravenously (i.v.) into C57BL/6 mice every other day for 3 times. The frequencies of IL-6⁺, IL-10⁺ and IL-21⁺ CD4⁺ T cells in the inguinal lymph node and spleen increased markedly after TRAPs administration (Fig. 1h). Consistently, in B16F10 tumor-bearing mice, the frequency of IL-6⁺, IL-21⁺ and IL-10⁺ CD4⁺ T cells in the draining lymph node and spleen were also increased (Fig. 1i). Taken together, these results show that TRAPs could modulate CD4⁺ T cell differentiation by inducing IL-6, IL-10, and IL-21 expression and suppressing their IFN-γ production.

TRAPs-induced IL-6, IL-10, and IL-21 production requires TLR2–MyD88 signaling

We then investigated the mechanism by which TRAPs induce IL-6, IL-10, and IL-21 in CD4⁺ T cells. Within the time frame of the induction of these cytokines, TRAPs adhered to the surface of CD4⁺ T cells in a dose-dependent manner without being internalized (Fig. 2a, b), suggesting the involvement of surface molecules on

TRAPs that interact with receptors on CD4⁺ T cells. TRAPs are enriched with various danger-associated molecular patterns (DAMPs) capable of stimulating pattern recognition receptors (PRRs) [17, 18]. CD4⁺ T cells expressed appreciable levels of TLR2 and TLR4 (Additional file 2: Figure S3a). We therefore examined whether TLR2 or TLR4 was involved in sensing TRAPs by CD4⁺ T cells. While TRAPs-induced IL-6, IL-10 and

IL-21 secretion by CD4⁺ T cells was independent of TLR4, *Tlr2*^{-/-} and *Myd88*^{-/-} CD4⁺ T cells were completely defective in producing these cytokines in response to TRAPs (Fig. 2c). Consistently, TLR2 on the surface of CD4⁺ T cells was in direct contact with TRAPs (Fig. 2d). In agreement with the above finding, *Tlr2*^{-/-} mice bearing B16F10 tumors had a significant reduction of IL-21⁺ and IL-10⁺ CD4⁺ T cells in the tumor tissue compared to WT tumor-bearing mice (Fig. 2e, f). Collectively, these results show that TRAPs induce CD4⁺ T cells to produce IL-6, IL-10, and IL-21 in a TLR2- and MyD88-dependent manner.

TRAPs-elicited IL-6 production by CD4⁺ T cells depends on NF-κB/p38/Akt signaling

We further sought to determine the signals downstream of TLR2 in the induction of IL-6, IL-10, and IL-21 by TRAPs. TRAPs treatment of WT CD4⁺ T cells resulted in the phosphorylation of NF-κB, Akt, p38 and STAT3, but not ERK1/2 or JNK1/2 (Fig. 3a), whereas TRAPs failed to induce NF-κB, Akt, p38 and STAT3 phosphorylation in *Tlr2*^{-/-} or *Myd88*^{-/-} CD4⁺ T cells when compared to CD4⁺ T cells from WT or *Tlr4*^{-/-} mice (Additional file 2: Figure S3b). Pretreatment of CD4⁺ T

cells with an inhibitor of NF-κB, Akt or p38 attenuated TRAPs-induced secretion of IL-6, IL-10 and IL-21, whereas the inhibition of JNK1/2 or ERK1/2 had no effect (Fig. 3b). Of note, pretreatment of CD4⁺ T cells with a STAT3 inhibitor diminished the production of IL-10 and IL-21, but not IL-6, in a dose-dependent manner (Fig. 3b, c), indicating that NF-κB, Akt, and p38 activation was needed for TRAPs-induced IL-6, IL-10 and IL-21 production but STAT3 activation was only required for IL-10 and IL-21 production.

The induction of IL-10 and IL-21 depends on autocrine IL-6 signaling

The IL-6-STAT3 pathway plays a crucial role in Th cell differentiation [22]. Upon IL-6 neutralization with a blocking antibody, the induction of IL-21 and IL-10 mRNA and proteins by TRAPs was completely abolished, with a concomitant decline of STAT3 phosphorylation (Fig. 3d, Additional file 2: Figure S4a). Consistently, TRAPs failed to induce IL-10 and IL-21 expression or STAT3 phosphorylation in *Il6*^{-/-} CD4⁺ T cells (Fig. 3e, Additional file 2: Figure S4b). Moreover, following i.v. administration of TRAPs, the frequencies of IL-10⁺ and IL-21⁺ CD4⁺ T cells in the

Fig. 3 TRAPs promote CD4⁺ T cell expression of IL-6 in an NF-κB/p38/Akt-dependent manner and induce IL-10 and IL-21 via IL-6-STAT3 signaling. **a** Western blot analyses of the phosphorylation of JNK, ERK, p38, Akt, IKKα/β, IκBα, p65 and STAT3 in CD4⁺ T cells treated with TRAPs (3 μg/ml) for the indicated time. **b** CD4⁺ T cells pre-treated with the indicated inhibitors for 1 h and then co-cultured with TRAPs (3 μg/ml) for 72 h. IL-6, IL-10 and IL-21 levels in the supernatants were determined by ELISA. **c** Western blot analyses of the phosphorylation of STAT3 in CD4⁺ T cells pre-treated with the STAT3 inhibitor Stattic at the indicated concentrations (0.5, 1 or 2 μM) for 1 h, and then co-cultured with TRAPs (3 μg/ml) for 2 h. ELISA of IL-10 and IL-21 secretion by CD4⁺ T cells treated as above for 72 h. **d** Western blot analyses of the phosphorylation of STAT3 in CD4⁺ T cells treated with anti-IL-6 neutralizing antibody (1 μg/ml) and TRAPs (3 μg/ml) for 2 h. ELISA of IL-10 and IL-21 secretion by CD4⁺ T cells treated as above for 72 h. **e** Western blot analyses of the phosphorylation of STAT3 in WT or *Il6*^{-/-} CD4⁺ T cells treated with TRAPs (3 μg/ml) for 2 h and ELISA of IL-10 and IL-21 secretion by WT or *Il6*^{-/-} CD4⁺ T cells for 72 h. **f, g** Flow cytometric and statistical analyses of the percentage of IL-21⁺ CD4⁺ T cells (f) or IL-10⁺ CD4⁺ T cells (g) in the iLN and spleens of WT or *Il6*^{-/-} C57BL/6 mice (*n* = 6) 7 days after i.v. administration of normal saline (NS) or TRAPs (30 μg/mouse) every other day for 3 times. Data (mean ± SEM) represent 3 independent experiments. *, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001; ns, not significant by one-way ANOVA with the Tukey-Kramer multiple test, 2-tailed unpaired t-test or Mann-Whitney U test

inguinal lymph node and spleen were much lower in *Il6*^{-/-} mice than in WT mice (Fig. 3f, g). Collectively, these results support a TRAPs-initiated regulatory cascade of CD4⁺ T cell differentiation involving TLR2-NF-κB/p38/Akt-dependent induction of auto-crine IL-6 which then promotes IL-10 and IL-21 expression via STAT3.

Hsp90α is a TRAPs surface ligand that induces IL-6 in CD4⁺ T cells

To identify the molecular components in TRAPs that are responsible for stimulating CD4⁺ T cell production of IL-6, we first subjected TRAPs to proteinase K digestion or sonication. These treatments impaired the ability of TRAPs to induce IL-6 from CD4⁺ T cells (Fig. 4a, b),

indicating that proteins on the surface, but not the soluble contents, of TRAPs are largely responsible for IL-6 induction in CD4⁺ T cells. In addition, TRAPs from the hepatic carcinoma Hepa1–6, lung cancer LLC or lymphoma EL4 cells also potently enhanced IL-6 secretion in

CD4⁺ T cells (Additional file 2: Figure S5a). Several ligands of TLR2, including HMGB1, Hsp60, Hsp70, and Hsp90α [18, 23], were enriched in and present on the surface of TRAPs (Fig. 4c, Additional file 2: Figure S5b). Blocking of Hsp90α, but not HMGB1, Hsp60 or Hsp70,

on the surface of TRAPs partially diminished TRAPs-induced IL-6 secretion by CD4⁺ T cells, indicating that other molecules on TRAPs may also play a role (Fig. 4d). Accordingly, an anti-Hsp90α antibody dose-dependently inhibited TRAPs binding to CD4⁺ T cells (Fig. 4e, f), reduced TRAPs-induced IL-6 secretion (Fig. 4g), and suppressed the activation of NF-κB, Akt and p38 (Additional file 2: Figure S5c). Remarkably, compared to intact TRAPs, tumor cell lysates containing an equal amount of total protein but much more Hsp90α, or sonicated TRAPs containing an equal amount of Hsp90α, or proteinase K-treated TRAPs were much less effective in inducing IL-6 secretion from CD4⁺ T cells (Fig. 4h). Taken together, these results show that membrane-bound Hsp90α on intact TRAPs effectively induces IL-6 expression from CD4⁺ T cells.

To further determine whether human TRAPs (hTRAPs) could induce human CD4⁺ T cells to produce IL-6, we collected hTRAPs from the culture media of 3 human tumor cell lines, A375, MDA-MB-231 and HepG2, and from the malignant effusions or ascites of 8 cancer patients (Additional file 1: Table S1). Western blotting analysis revealed that LC3-II was expressed at high levels in the collected hTRAPs and Hsp90α was detected in most of hTRAPs (Additional file 2: Figure S5d). RT-PCR analysis and ELISA showed that hTRAPs from cancer patients and tumor cell lines efficiently induced human peripheral blood CD4⁺ T cells to express *IL6* transcript and secrete IL-6 (Fig. 4i, Additional file 2: Figure S5e). Similar to mouse TRAPs, hTRAPs-induced IL-6 transcription and secretion by human CD4⁺ T were almost completely abolished by pretreatment of hTRAPs with an anti-hsp90α blocking antibody (Fig. 4j, Additional file 2: Figure S5f). Altogether, these results indicate that induction of CD4⁺ T cells IL-6 expression by HSP90α on the surface of TRAPs is a common characteristic in humans and mice.

TRAPs-elicited CD4⁺ T cells (T_{TRAP}) suppress effector T cells and promote tumorigenesis

To characterize the function of TRAPs-elicited CD4⁺ T cells (T_{TRAP}), we activated CD4⁺ and CD8⁺ T cells with anti-CD3 and anti-CD28 in culture supernatants harvested from T_{TRAP} or control CD4⁺ T cells. T_{TRAP} supernatants (SN/T_{TRAP}) strongly suppressed the secretion of IFN-γ by activated CD4⁺ and CD8⁺ T cells (Fig. 5a). Pretreatment of SN/T_{TRAP} with a neutralizing antibody against IL-6 or IL-10, but not IL-21, abolished its suppressive effect on IFN-γ production by activated CD4⁺ and CD8⁺ T cells (Fig. 5b). We then transferred control CD4⁺ T cells or T_{TRAP} into C57BL/6 mice that had received OVA-specific Vβ5.1⁺CD8⁺ OT-I T cells and vaccinated with OVA-loaded dendritic cells (DC_{OVA}). DC_{OVA} vaccination induced the expansion of Vβ5.1⁺CD8⁺ OT-I T cells in the host, which was suppressed by the adoptive

transfer of T_{TRAP} but not control CD4⁺ T cells (Fig. 5c). Moreover, the transfer of T_{TRAP} but not control CD4⁺ T cells led to a decrease of IFN-γ⁺ CD8⁺ and CD4⁺ T cells induced by DC_{OVA} vaccination (Fig. 5d). Therefore, T_{TRAP} could suppress T cell IFN-γ response in vivo.

To see whether T_{TRAP} have a tumor-promoting effect in vivo, we subcutaneously (s.c.) inoculated B16F10 melanoma cells into C57BL/6 mice with or without co-administration of control CD4⁺ T cells or T_{TRAP}. Co-administration of B16F10 cells with T_{TRAP} enhanced tumor growth as compared to inoculation of B16F10 cells alone or co-administration with control CD4⁺ T cells (Fig. 5e). When B16F10 melanoma cells were inoculated i.v. together with T_{TRAP} T_{TRAP} promoted tumor metastasis to the lung (Fig. 5f). Collectively, these results show that T_{TRAP} could promote tumor growth and metastasis in vivo.

T_{TRAP} enhance regulatory B cell function via IL-6, IL-10, and IL-21

To better define the immunosuppressive capacity of T_{TRAP} we examined the impact of T_{TRAP} on regulatory B cell (Breg) differentiation. In accordance with our earlier findings [18], TRAPs induced B cell differentiation into IL-10-producing Bregs (Fig. 6a). Moreover, co-culture of B cells and CD4⁺ T cells in the presence of TRAPs led to a marked increase in Bregs differentiation (Fig. 6a). Consistently, SN/T_{TRAP} could directly promote IL-10⁺ Bregs differentiation and IL-10 secretion (Additional file 2: Figure S6a, b). Next, adoptive transfer of T_{TRAP} but not control CD4⁺ T cells, also significantly increased the frequency and number of IL-10⁺ Bregs in vivo (Fig. 6b). Therefore, TRAPs can promote IL-10⁺ Breg differentiation directly by activating on B cells and indirectly by conditioning CD4⁺ T cells.

We then investigated the mechanism by which T_{TRAP} promote IL-10⁺ Bregs differentiation. In agreement with the above results, culturing B cells in SN/T_{TRAP} together with TRAPs resulted in a synergistic increase the frequencies of IL-10⁺ Bregs and IL-10 secretion as compared to TRAPs or SN/T_{TRAP} alone, whereas the supernatant of control CD4⁺ T cells did not have this effect (Fig. 6c). Neutralizing IL-6, IL-10 or IL-21 partially abolished the effect of SN/T_{TRAP} in promoting IL-10 production of TRAPs-induced B cells (Fig. 6c). These data indicate that secreted cytokines, including IL-6, IL-10, and IL-21, from T_{TRAP} were involved in promoting Bregs differentiation.

Subsequently, the potential regulatory effect of B cells pretreated by TRAPs and SN/T_{TRAP} (B_{TRAP} + SN/T_{TRAP}) on the antitumor effector function of T cells was assessed. IFN-γ production by activated CD4⁺ and CD8⁺ T cells was strongly suppressed when these cells were cultured in the supernatants from B_{TRAP} + SN/T_{TRAP} (SN/

Fig. 5 IL-6/IL-10 from T_{TRAP} is responsible for tumor growth and metastasis. **a** Flow cytometric and statistical analyses of the percentage of IFN- γ ⁺CD4⁺ and IFN- γ ⁺CD8⁺ T cells treated with the supernatants from T_{TRAP} (SN/T_{TRAP}) or control CD4⁺ T cells for 3 d in the presence of anti-CD3/CD28. **b** SN/T_{TRAP} was pretreated with anti-IL-6, IL-10, IL-21 neutralizing antibodies respectively, and then co-cultured with the purified CD4⁺ T cells and CD8⁺ T cells in the presence of anti-CD3/CD28 for 3 d. The percentages of IFN- γ ⁺CD4⁺ T cells and CD8⁺ T cells were evaluated by flow cytometry. **c** C57BL/6 mice were adoptively transferred i.v. with OT-I spleen cells and then vaccinated s.c. with OVA-loaded DC on day 1, 4 and 7, following injection i.v. with T_{TRAP} or control CD4⁺ T cells on day 2, 5 and 8. On the 15th day, the frequencies and the number of V β 5.1⁺CD8⁺ T cells in spleen were analyzed by flow cytometry. **d** C57BL/6 mice were vaccinated with OVA-loaded DC and following adoptively transferred with T_{TRAP} or CD4⁺ T cells. On the 15th day, the splenocytes were re-stimulated with OVA-protein for 24 h, and the frequencies of IFN- γ ⁺ T cells were determined by flow cytometry. **e** B16F10 tumor cells were mixed with T_{TRAP} or control CD4⁺ T cells and injected s.c. into C57BL/6 mice ($n = 6$ per group). The growth of the tumor was monitored. **f** B16F10 tumor cells were intravenously injected into C57BL/6 mice ($n = 4$ to 6 per group) to establish a lung metastasis model. Subsequently, T_{TRAP} or control CD4⁺ T cells were adoptively transferred i.v. 3 times with 1 d of interval. Three weeks later, the tumor nodules in the lungs were examined. Data (mean $\pm$ SEM) represent 3 independent experiments. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ns, not significant, by 1-way ANOVA with the Tukey-Kramer multiple test, 2-tailed unpaired t-test or Mann-Whitney U test

B_{TRAP} + SN/T_{TRAP}), and the suppressive activity of the SN/B_{TRAP} + SN/T_{TRAP} on IFN- γ production by T cell was largely abolished using an anti-IL-10 neutralizing antibody (Fig. 6d). To further investigate the suppressive effects of B_{TRAP} + SN/T_{TRAP} on effector T cell response in vivo, C57BL/6 mice, with or without adoptive transfer of OT-I cells were vaccinated with DC_{OVA} and

subsequently were adoptively transferred with B_{TRAP} + SN/T_{TRAP} or B_{TRAP}. DC_{OVA} vaccination induced the expansion of V β 5.1⁺CD8⁺ OT-I T cells in the recipient mice. Adoptive transfer of B_{TRAP} inhibited the expansion of OT-I T cells, and the transfer of B_{TRAP} + SN/T_{TRAP} resulted in a more pronounced and almost complete inhibition of the expansion of OT-I T cells (Fig. 6e).

Moreover, adoptive transfer of B_{TRAP} + SN/T_{TRAP} decreased the numbers of IFN-γ⁺ CD8⁺ and CD4⁺ T cells induced by DC_{OVA} vaccination (Fig. 6f) and promoted the growth of B16F10 melanoma cells and their metastasis to the lung (Fig. 6g, h). Taken together, these results suggest that IL-6, IL-10, and IL-21 from T_{TRAP} augment the differentiation and immunosuppressive function of TRAPs-induced B cells to facilitate tumor growth and metastasis.

Inhibition of autophagosomes formation or IL-6 secretion delay tumor growth

Having shown a critical role of TRAPs in the inhibition of anti-tumor immunity, we explored whether inhibition of TRAPs formation by targeting *Becn1*, a gene essential for autophagosome formation, could abolish the generation of tumor-promoting T_{TRAP} (Additional file 2:

Figure S7a, b). *Becn1* knock-down in B16F10 cells diminished intracellular LC3-II accumulation and markedly reduced TRAPs secretion (Additional file 2: Figure S7b, c). The culture media collected from *Becn1* knock-down B16F10 cells had reduced ability to induce IL-6, IL-10, and IL-21 in CD4⁺ T cells (Fig. 7a). In the mice bearing knock-down B16F10 tumors, the frequency of IL-21⁺ and IL-10⁺ CD4⁺ T cells in the tumor draining lymph node and tumor tissue and the serum IL-6 level were significantly reduced as compared to those in the mice bearing control tumors (Fig. 7b–d). Moreover, the frequency of IL-10⁺ B cells and IFN-γ⁺CD4⁺ T cells in mice bearing *Becn1* knock-down tumors was markedly decreased and increased, respectively (Fig. 7e, f). Additionally, *Becn1* knock-down B16F10 cells exhibited significantly slower growth in vivo (Additional file 2: Figure S7d). These results indicate that inhibition of tumor

Fig. 7 (See legend on next page.)

(See figure on previous page.)

Fig. 7 Knockdown of Beclin1 or knockout of IL-6 inhibits tumor growth and alters CD4⁺T and B cells polarization. **a** ELISA of IL-6, IL-10, and IL-21 secretion by CD4⁺ T cells treated with CM, or media from negative control (NC) or *Becn1*-knockdown (KD) B16F10 cells in the presence of anti-CD3/CD28 for 72 h. **b-f** Control (NC) or *Becn1*-KD B16F10 cells were inoculated s.c. into C57BL/6 mice ($n = 6$ per group). Twenty-one days later, serum IL-6 level was measured by ELISA (**b**). The frequency of IL-10⁺ CD4⁺ T cells (**c**), IL-21⁺ CD4⁺ T cells (**d**), IFN- γ ⁺ CD4⁺ T cells (**e**), and IL-10⁺ B cells (**f**) in tumor-draining lymph nodes (dLN) or tumor tissues (Tumor) were analyzed by flow cytometry. **g-j** The tumor-bearing mice ($n = 5$ per group) were depleted of either CD4⁺ or CD8⁺ T cells or CD20⁺ B cells by intravenous administration of 250 μ g/mouse of anti-mouse CD4 or anti-mouse CD8 antibodies twice weekly throughout the course of tumor growth, or 250 μ g of anti-mouse CD20 antibodies, respectively. Control mice were treated with isotype-matched control antibodies. **g** Tumor growth was monitored by calipers. **h, i** The frequency of CD4⁺ IFN- γ ⁺ and CD8⁺ IFN- γ ⁺ T cells in tumor infiltrating lymphocytes were determined by intracellular staining. **j** The frequency of B cells in tumor infiltrating lymphocytes were determined by flow cytometry. **k-n** WT or *Il6*^{-/-} C57BL/6 mice were inoculated s.c. with B16F10 cells. On day 21, the frequencies of IL-10⁺ CD4⁺ T cells (**k**), IL-21⁺ CD4⁺ T cells (**l**), and IL-10⁺ B cells (**m**) in dLN and tumor tissues were evaluated by flow cytometry. **n** Tumor size was measured by caliper. **o, p** B16F10 tumor cells were mixed with WT T_{TRAP} or *Il6*^{-/-} T_{TRAP} and injected s.c. into C57BL/6 mice ($n = 6$ per group). The growth of the tumor was monitored. **o, q** B16F10 tumor cells were i.v. injected into C57BL/6 mice ($n = 6$ per group). Subsequently, WT T_{TRAP} or *Il6*^{-/-} T_{TRAP} were transferred i.v. every other day for 3 times. Three weeks later, tumor nodules in the lungs were examined. **r** A proposed model for the mechanisms and immunosuppressive functions of TRAPs-induced CD4⁺ T cells. Data (mean $\pm$ SEM) represent 3 independent experiments. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ns, not significant, by 1-way ANOVA with the Tukey-Kramer multiple test, 2-tailed unpaired t-test or Mann-Whitney U test

autophagosome formation and release could enhance anti-tumor immunity and inhibit tumor growth in vivo.

Furthermore, the growth of both the negative control and *Becn1* knock-down B16F10 tumors was inhibited in mice depleted of B cells or CD4⁺ T cells (Fig. 7g, Additional file 2: Figure S8). Depletion of CD8⁺ T cells resulted in accelerated growth of *Becn1* knock-down but not negative control tumors (Fig. 7g, Additional file 2: Figure S8). Besides, the frequency of IFN- γ -producing CD4⁺ T cells and CD8⁺ T cells in *Becn1* knock-down tumor tissue was markedly increased (Fig. 7h, i). Notably, B-cell or CD4⁺ T-cell depletion resulted in a significant increase of the percentage of intra-tumoral IFN- γ ⁺ CD4⁺ or CD8⁺ T cells (Fig. 7h, i). The frequency of tumor-infiltrating B cells was markedly reduced upon CD4⁺ T cell depletion (Fig. 7j). These results suggest that the effector function of CD8⁺ T cells in the tumors was dampened by CD4⁺ T cells or B cells. In conclusion, TRAPs-educated CD4⁺ T cells play an important role in promoting tumor growth by inhibiting effector T cell function.

To determine the role of CD4⁺ T cell-derived IL-6 in the differentiation of IL-10- and IL-21-producing CD4⁺ T cells and IL-10-producing Bregs in vivo, WT or *Il6*^{-/-} mice were s.c. inoculated with B16F10 cells. Consistent with previous results, the frequencies of IL-10⁺ and IL-21⁺ CD4⁺ T cells (Fig. 7k, l) and IL-10⁺ B cells (Fig. 7m) in tumor-draining lymph nodes and tumor tissues from *Il6*^{-/-} tumor-bearing mice were significantly decreased. Accordingly, B16F10 tumors grew more slowly in *Il6*^{-/-} mice than in WT mice (Fig. 7n). We then inoculated mice with B16F10 cells together with either WT T_{TRAP} or *Il6*^{-/-} T_{TRAP}. Mice co-inoculated with B16F10 cells and WT T_{TRAP} showed accelerated growth and lung metastasis as compared to those inoculated with B16F10 cells alone (Fig. 7o-q). In contrast, co-inoculation of B16F10 cells with *Il6*^{-/-} T_{TRAP} resulted in no

enhancement of tumor growth and lung metastasis, and the mice even exhibited slightly, albeit not statistically significant, retarded tumor growth (Fig. 7o-q). These results corroborate the conclusion that T_{TRAP} rely on IL-6 to dampen T cell-mediated antitumor immunity and foster tumor progression, and suggest that targeting TRAPs or IL-6 may be an effective therapeutic strategy for improving cancer immunotherapy.

Discussion

In addition to soluble factors, tumor cell-derived extracellular vesicles are being recognized as critical modulators of host anti-tumor immunity during tumor progression [7, 8, 18, 19, 24]. Among them are autophagosomes generated by secretory autophagy. In contrast to canonical autophagy that functions in a primarily degradative capacity to sustain cellular metabolism and homeostasis and is often induced conditions of cellular stress, such as nutrient starvation, organelle damage, and pathogen infection, secretory autophagy is a non-degradative mechanism for cellular trafficking and unconventional protein secretion [10, 11, 13, 14, 25]. Secretory autophagosomes fail to fuse with lysosomes, but are released into the extracellular environment through fusing with the plasma membrane or other pathways [15, 26]. Abundant autophagosomes have been detected in gastrointestinal tumors and invasive melanomas and have been associated with tumor cell proliferation, metastasis, and poor prognosis [27, 28]. Our previous studies showed that extracellular autophagosomes harvested from the supernatant of tumor cells or malignant effusions and ascites of cancer patients, which we have termed as TRAPs, could promote the generation of IL-10⁺ Bregs, reactive oxygen species (ROS)-producing neutrophils, and PD-L1^{hi} macrophages exerting immunoinhibitory activities [18–20].

CD4⁺ T cells that infiltrate advanced solid tumors consist of different effector cells, such as Th1, Th2, Th17, Tfh or regulatory T cells (Tregs), with distinct impact on anti-tumor immunity, immune escape, angiogenesis and metastasis [2, 4, 29], but the influence of the tumors on CD4⁺ effector T cell differentiation remains incompletely understood. Here, we have revealed a TRAPs-mediated regulatory mechanism of CD4⁺ T cells differentiation whereby HSP90α on the surface of TRAPs educate CD4⁺ T cells via a TLR2–autocrine IL-6 cascade to express IL-10 and IL-21 and engender immune suppression to promote tumor growth and metastasis (Fig. 7r). Our findings have revealed TRAPs as one of the tumor-derived extracellular vesicles that could inhibit anti-tumor immune response by enhancing the generation of immunosuppressive cells.

TLRs play crucial roles in the innate host defense as well as the control of adaptive immunity [30, 31]. Our findings indicated TLR2 as a key receptor for TRAPs-mediated IL-6 expression by CD4⁺ T cells. Exogenous pathogen-associated molecular patterns (PAMPs) and endogenous DAMPs can be recognized by TLRs to trigger the production of various inflammatory mediators [30]. The current findings showed that TRAPs-mediated regulation of CD4⁺ T cell differentiation involved membrane-associated Hsp90α. Evidences suggested that extracellular Hsp90α could be released to the extracellular space via unconventional secretion, such as exosomes and necrosis [32]. We observed Hsp90α on the surface of TRAPs, indicating that secretory autophagosomes may also be involved in the release of Hsp90α. Moreover, extracellular Hsp90α was reported to function as a DAMP and provoke biological effects through cell surface receptors, including TLRs and CD91 [23, 33]. Early work showed that heat shock proteins gp96, Hsp90, Hsp70, and calreticulin could function as potential adjuvants to stimulate DC antigen cross-presentation and maturation through the CD91 receptor [33], but Hsp90α was more recently found to also stimulate tumor proliferation and metastasis through binding to cancer cell surface CD91 and be positively correlated with tumor malignancy in cancer patients [34–36]. The present study uncovers a new role of Hsp90α on the surface of TRAPs as a cancer-associated pathological factor that interferes with host anti-tumor immunity.

Chronic inflammation and increased levels of inflammatory mediators at the tumor site can reroute the immunomodulatory response towards a cancer-promoting direction [4, 37, 38]. IL-6 has a profound effect on CD4⁺ T cells survival and proliferation [39]. Otherwise, studies also showed that IL-6 has inhibitory effects via the induction of IL-10-producing T and B cells [40, 41]. Moreover, IL-6 also dampens Th1 differentiation and inhibits CD8⁺ T cell activation and cytokine production [42, 43]. Consistent with the above results, we provided evidences

that TRAPs stimulated IL-10 and IL-21 production in CD4⁺ T cells via an autocrine IL-6 loop. Moreover, IL-6 from T_{TRAP} remarkably suppressed T cell anti-tumor effector function. IL-21 has been identified to be derived mainly from Tfh cells, which was thought to regulate the proliferation, class switching, and plasmacytoid differentiation of B cells and promote the generation and proliferation of human antigen-specific cytotoxic T-cell responses [4, 44, 45]. Mounting evidences have shown that IL-21 also has anti-inflammatory activities by inhibiting DC maturation and stimulating IL-10 production in T and B cells [46–48]. Nonetheless, the role of CD4⁺ T cells in Bregs differentiation in the tumor microenvironment has not been addressed. In our investigation, the IL-21⁺ T_{TRAP} displayed Tfh-associated molecules CXCR5 and Bcl-6. Interestingly, IL-6, IL-10, and IL-21 secretion by T_{TRAP} synergistically enhanced TRAPs-elicited Breg differentiation and immunosuppressive function. These findings together imply that T_{TRAP}-derived IL-21 is a pleiotropic effector that can either facilitate or thwart tumor growth depending on the cytokine milieu in the tumor microenvironment, warranting careful consideration of the selective targeting of IL-6 or IL-21 for the treatment of cancer in the future.

Many recent studies have suggested that inhibiting tumor autophagy may have anti-tumor effects by modulating the tumor microenvironment [49–51]. Consistent with this notion, we found that inhibiting autophagy by targeting the key autophagy gene *Becn1*, which led to a substantial decrease in extracellular TRAPs, could inhibit tumor growth in mice. Of note, inhibiting autophagy resulted in a significant decrease in the frequency of IL-10⁺ B cells, IL-21⁺ and IL-10⁺CD4⁺ T cells, as well as a significant increase in IFN-γ⁺CD4⁺ T cells, in the tumor-draining lymph nodes and tumor tissue. Thus, intervening tumor release of TRAPs could be an effective strategy for cancer therapy.

Conclusions

In this study, we have revealed that TRAPs can educate CD4⁺ T cells to promote tumor growth and metastasis through an HSP90α–TLR2–IL-6–IL-10/IL-21 axis and the induction of IL-10⁺ Bregs. Our study reveals a novel cellular and molecular mechanism of how tumor-derived extracellular vesicles regulate CD4⁺ effector T cell function and highlights TRAPs and their membrane-bound DAMPs as important therapeutic targets to reverse the immunosuppressive tumor microenvironment.

Additional files

Additional file 1: Table S1. Clinical and demographic characteristics of the patients presenting with malignant pleural effusions or ascites. **Table S2.** Antibodies used in flow cytometry. **Table S3.** Primer

sequences for qPCR used in our study. **Table S4.** Antibodies used for immunoblotting. (DOCX 25 kb)

Additional file 2: Figure S1. Effect of TRAPs on differentiation of CD4⁺ T cells. **Figure S2.** Comparison of LC3B⁺ EVs and LC3B⁻ EVs in instructing CD4⁺ T cells. **Figure S3** TRAPs induce Akt/p38/NF- κ B/STAT3 activation in CD4⁺ T cells via the TLR2-MyD88 pathway. **Figure S4** IL-10 and IL-21 production by TRAPs-induced CD4⁺ T cells occurs via autocrine IL-6. **Figure S5** Hsp90 α on TRAPs is essential for the induction of IL-6 secretion from CD4⁺ T cells. **Figure S6.** The secretion of cytokines from TRAPs-induced CD4⁺ T cells (T_{TRAP}) mediate IL-10-producing B cell differentiation. **Figure S7** Evaluation of Beclin-1 expression in B16F10 cells. **Figure S8** Depletion of selected cellular subsets during tumor growth. (DOCX 2.11 mb)

Abbreviations

APCs: Antigen-presenting cells; Bregs: Regulatory B cells; CFSE: Carboxyfluorescein succinimidyl ester; DAMPs: Damage-associated molecular pattern molecules; dLNs: Draining lymph nodes; EVs: Extracellular vesicles; HMGB1: High mobility group box 1; HSP: Heat shock protein; KD: Knock down; KO: Knock out; mAb: Monoclonal antibody; NS: Normal saline; PAMPs: Pathogen-associated molecular patterns; PBMC: Peripheral blood mononuclear cell; ROS: Reactive oxygen species; TB: Tumor-bearing; TF: Tumor-free; TLRs: Toll-like receptors; TRAPs: Tumor cell-released autophagosomes; WT: Wild-type

Acknowledgments

The authors thank Dr. Guozheng Wang (University of Liverpool, Liverpool, UK) for helpful discussion, Dr. Yong Lin (Zhongda Hospital, Medical School of Southeast University) for providing human specimens.

Authors' contributions

LXW, YQC, YLC, KC and HMH designed and discussed this research. YQC, PCL, NP, RG, ZFW, TYZ, FH and FYW performed the experiments. NP, KC and JPZ provided experimental support. YLC provided malignant pleural effusions and ascites from tumor patients. LXW and YQC prepared the figures and wrote the manuscript. KC and HMH contributed to manuscript editing. All authors analyzed and discussed the data. All authors read and approved the final manuscript.

Funding

This study was supported by the National Natural Science Foundation of China (No. 31670918, 31370895 and 31170857 to LX. Wang, No. 81872122 to Y.L. Cai). The Fundamental Research Funds for the Central Universities and Postgraduate Research & Practice Innovation Program of Jiangsu Province (No. KYCX17_0166 to Y.Q. Chen).

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

All animal experiments were approved by the Animal Care and Use Committee of Southeast University. All human experiments were approved by the Ethics Committee for Human Studies of Southeast University and performed under protocol 2016ZDKYSB112. Informed consent was obtained from all patients.

Consent for publication

All authors provide their consent for publication of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Microbiology and Immunology, Medical School of Southeast University, 87 Dingjiaqiao Road, Nanjing 210009, China. ²Department of Obstetrics and Gynecology, Zhongda Hospital, Medical School of Southeast University, 87 Dingjiaqiao Road, Nanjing 210009, China. ³Department of Gastroenterology, Zhongda Hospital, Medical School of Southeast University, Nanjing 210009, China. ⁴Robert W. Franz Cancer Research Center, Earle A. Chiles Research Institute, Providence Portland Medical Center, Portland, OR

97213, USA. ⁵Institutes of Biology and Medical Sciences, Soochow University, Suzhou 215123, China. ⁶Department of Obstetrics and Gynecology and Barbara Ann Karmanos Cancer Institute, Wayne State University, Mucosal Immunology Studies Team, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Detroit, MI 48201, USA. ⁷Jiangsu Province Hospital of Traditional Chinese Medicine, The Affiliated Hospital of Nanjing University of Chinese Medicine, Nanjing 210029, China.

Received: 17 February 2019 Accepted: 19 June 2019

Published online: 12 July 2019

References

- Borst J, Ahrends T, Båbala N, Melief CJM, Kastenmüller W. CD4⁺ T cell help in cancer immunology and immunotherapy. *Nat Rev Immunol*. 2018;18(10):635–47.
- Kim HJ, Cantor H. CD4 T-cell subsets and tumor immunity: the helpful and the not-so-helpful. *Cancer Immunol Res*. 2014;2(2):91–8.
- Fridman WH, Pages F, Sautès-Fridman C, Galon J. The immune contexture in human tumours: impact on clinical outcome. *Nat Rev Cancer*. 2012;12(4):298–306.
- Chen MM, Xiao X, Lao XM, Wei Y, Liu RX, Zeng QH, et al. Polarization of tissue-resident TFH-like cells in human hepatoma bridges innate monocyte inflammation and M2b macrophage polarization. *Cancer Discovery*. 2016; 6(10):1182–95.
- Kumai T, Oikawa K, Aoki N, Kimura S, Harabuchi Y, Celis E, et al. Tumor-derived TGF- β and prostaglandin E2 attenuate anti-tumor immune responses in head and neck squamous cell carcinoma treated with EGFR inhibitor. *J Transl Med*. 2014;12:265.
- Whiteside TL. Exosomes and tumor-mediated immune suppression. *J Clin Invest*. 2016;126(4):1216–23.
- Becker A, Thakur BK, Weiss JM, Kim HS, Peinado H, Lyden D. Extracellular vesicles in Cancer: cell-to-cell mediators of metastasis. *Cancer Cell*. 2016; 30(6):836–48.
- Chen G, Huang AC, Zhang W, Zhang G, Wu M, Xu W, et al. Exosomal PD-L1 contributes to immunosuppression and is associated with anti-PD-1 response. *Nature*. 2018;560(7718):382–6.
- Deretic V, Jiang SY, Dupont N. Autophagy intersections with conventional and unconventional secretion in tissue development, remodeling and inflammation. *Trends Cell Biol*. 2012;22(8):397–406.
- Zhang M, Kenny SJ, Ge L, Xu K, Schekman R. Translocation of interleukin-1 β into a vesicle intermediate in autophagy-mediated secretion. *Elife*. 2015;4.
- Thorburn J, Frankel AE, Thorburn A. Regulation of HMGB1 release by autophagy. *Autophagy*. 2009;5(2):247–9.
- Fader CM, Aguilera MO, Colombo MI. ATP is released from autophagic vesicles to the extracellular space in a VAMP7-dependent manner. *Autophagy*. 2012;8(12):1741–56.
- Nuchel J, Ghatak S, Zuk AV, Illerhaus A, Morgelin M, Schonborn K, et al. TGF β 1 is secreted through an unconventional pathway dependent on the autophagic machinery and cytoskeletal regulators. *Autophagy*. 2018; 14(3):465–86.
- Bel S, Pendse M, Wang YH, Li Y, Ruhn KA, Hassell B, et al. Paneth cells secrete lysozyme via secretory autophagy during bacterial infection of the intestine. *Science*. 2017;357(6355):1047–51.
- Mutsaers Y, Altan-Bonnet N. Enterovirus transmission by secretory autophagy. *Viruses-Basel*. 2018;10(3).
- Li YH, Wang LX, Pang PY, Twitty C, Fox BA, Aung S, et al. Cross-presentation of tumor associated antigens through tumor-derived autophagosomes. *Autophagy*. 2009;5(4):576–7.
- Li YH, Wang LX, Pang PY, Cui ZH, Aung S, Haley D, et al. Tumor-derived autophagosome vaccine: mechanism of cross-presentation and therapeutic efficacy. *Clin Cancer Res*. 2011;17(22):7047–57.
- Zhou M, Wen ZF, Cheng F, Ma J, Li WX, Ren HY, et al. Tumor-released autophagosomes induce IL-10-producing B cells with suppressive activity on T lymphocytes via TLR2-MyD88-NF- κ B signal pathway. *Oncoimmunology*. 2016;5(7):e1180485.
- Gao R, Ma J, Wen Z, Yang P, Zhao J, Xue M, et al. Tumor cell-released autophagosomes (TRAP) enhance apoptosis and immunosuppressive functions of neutrophils. *Oncoimmunology*. 2018;7(6):e1438108.
- Wen Z-F, Liu H, Gao R, Zhou M, Ma J, Zhang Y, et al. Tumor cell-released autophagosomes (TRAPs) promote immunosuppression through induction

- of M2-like macrophages with increased expression of PD-L1. *J Immunother Cancer*. 2018;6(1):151.
21. Chen YQ, Zheng L, Aldarouish M, Zhou ZH, Pan N, Liu JQ, et al. Wnt pathway activator TWS119 enhances the proliferation and cytolytic activity of human gamma delta T cells against colon cancer. *Exp Cell Res*. 2018;362(1):63–71.
 22. Hunter CA, Jones SA. IL-6 as a keystone cytokine in health and disease. *Nat Immunol*. 2015;16(5):448–57.
 23. Calderwood SK, Gong JL, Murshid A. Extracellular HSPs: the complicated roles of extracellular HSPs in immunity. *Front Immunol*. 2016;7:159.
 24. Xu R, Rai A, Chen M, Suwakulsiri W, Greening DW, Simpson RJ. Extracellular vesicles in cancer — implications for future improvements in cancer care. *Nat Rev Clin Oncol*. 2018;15(10):617–38.
 25. Son SM, Cha MY, Choi H, Kang S, Choi H, Lee MS, et al. Insulin-degrading enzyme secretion from astrocytes is mediated by an autophagy-based unconventional secretory pathway in Alzheimer disease. *Autophagy*. 2016;12(5):784–800.
 26. Ponpuak M, Mandell MA, Kimura T, Chauhan S, Cleyrat C, Deretic V. Secretory autophagy. *Curr Opin Cell Biol*. 2015;35:106–16.
 27. Lazova R, Camp RL, Klump V, Siddiqui SF, Amaravadi RK, Pawelek JM. Punctate LC3B expression is a common feature of solid tumors and associated with proliferation, metastasis, and poor outcome. *Clin Cancer Res*. 2012;18(2):370–9.
 28. Katheder NS, Khezri R, O'Farrell F, Schultz SW, Jain A, Rahman MM, et al. Microenvironmental autophagy promotes tumour growth. *Nature*. 2017;541(7637):417–20.
 29. DuPage M, Bluestone JA. Harnessing the plasticity of CD4(+) T cells to treat immune-mediated disease. *Nat Rev Immunol*. 2016;16(3):149–63.
 30. Kugelberg E. Pattern Recognition Receptors Curbing gut inflammation. *Nat Rev Immunol*. 2014;14(9):583.
 31. Mercier BC, Cottalorda A, Coupet CA, Marvel J, Bonnefoy-Berard N. TLR2 engagement on CD8 T cells enables generation of functional memory cells in response to a suboptimal TCR signal. *J Immunol*. 2009;182(4):1860–7.
 32. McCready J, Sims JD, Chan D, Jay DG. Secretion of extracellular hsp90 alpha via exosomes increases cancer cell motility: a role for plasminogen activation. *BMC Cancer*. 2010;10:294.
 33. Basu S, Binder RJ, Ramalingam T, Srivastava PK. CD91 is a common receptor for heat shock proteins gp96, hsp90, hsp70, and calreticulin. *Immunity*. 2001;14(3):303–13.
 34. Shi YK, Liu XQ, Lou JT, Han XH, Zhang LJ, Wang QT, et al. Plasma levels of heat shock protein 90 alpha associated with lung Cancer development and treatment responses. *Clin Cancer Res*. 2014;20(23):6016–22.
 35. Chen JS, Hsu YM, Chen CC, Chen LL, Lee CC, Huang TS. Secreted heat shock protein 90 alpha induces colorectal Cancer cell invasion through CD91/LRP-1 and NF-kappa B-mediated integrin alpha(V) expression. *J Biol Chem*. 2010;285(33):25458–66.
 36. Chen C-C, Chen L-L, Li C-P, Hsu Y-T, Jiang S-S, Fan C-S, et al. Myeloid-derived macrophages and secreted HSP90 α induce pancreatic ductal adenocarcinoma development. *Oncoimmunology*. 2018;7(5):e1424612.
 37. Elinav E, Nowarski R, Thaiss CA, Hu B, Jin CC, Flavell RA. Inflammation-induced cancer: crosstalk between tumours, immune cells and microorganisms. *Nat Rev Cancer*. 2013;13(11):759–71.
 38. Coward J, Kulbe H, Chakravarty P, Leader D, Vassileva V, Leinster DA, et al. Interleukin-6 as a therapeutic target in human ovarian Cancer. *Clin Cancer Res*. 2011;17(18):6083–96.
 39. Teague TK, Marrack P, Kappler JW, Vella AT. IL-6 rescues resting mouse T cells from apoptosis. *J Immunol*. 1997;158(12):5791–6.
 40. Jin JO, Han XZ, Yu Q. Interleukin-6 induces the generation of IL-10-producing Tr1 cells and suppresses autoimmune tissue inflammation. *J Autoimmun*. 2013;40:28–44.
 41. Rosser EC, Oleinika K, Tonon S, Doyle R, Bosma A, Carter NA, et al. Regulatory B cells are induced by gut microbiota-driven interleukin-1 beta and interleukin-6 production. *Nat Med*. 2014;20(11):1334–9.
 42. Wu W, Dietze KK, Gibbert K, Lang KS, Trilling M, Yan HM, et al. TLR ligand induced IL-6 counter-regulates the anti-viral CD8(+) T cell response during an acute retrovirus infection. *Sci Rep UK*. 2015;5:10501.
 43. Tsukamoto H, Senju S, Matsumura K, Swain SL, Nishimura Y. IL-6-mediated environmental conditioning of defective Th1 differentiation dampens antitumour immune responses in old age. *Nat Commun*. 2015;6:6702.
 44. Tangye SG, Ma CS, Brink R, Deenick EK. The good, the bad and the ugly - T-FH cells in human health and disease. *Nat Rev Immunol*. 2013;13(6):412–26.
 45. Crotty S. T follicular helper cell differentiation, function, and roles in disease. *Immunity*. 2014;41(4):529–42.
 46. Brandt K, Bulfone-Paus S, Foster DC, Ruckert R. Interleukin-21 inhibits dendritic cell activation and maturation. *Blood*. 2003;102(12):4090–8.
 47. Spolski R, Kim HP, Zhu W, Levy DE, Leonard WJ. IL-21 mediates suppressive effects via its induction of IL-10. *J Immunol*. 2009;182(5):2859–67.
 48. Yoshizaki A, Miyagaki T, DiLillo DJ, Matsushita T, Horikawa M, Kountikov EI, et al. Regulatory B cells control T-cell autoimmunity through IL-21-dependent cognate interactions. *Nature*. 2012;491(7423):264–8.
 49. Sousa CM, Biancur DE, Wang XX, Halbrook CJ, Sherman MH, Zhang L, et al. Pancreatic stellate cells support tumour metabolism through autophagic alanine secretion. *Nature*. 2016;536(7617):479–83.
 50. Yang AN, Herter-Sprie G, Zhang HK, Lin EY, Biancur D, Wang XX, et al. Autophagy sustains pancreatic Cancer growth through both cell-autonomous and nonautonomous mechanisms. *Cancer Discov*. 2018;8(3):276–87.
 51. Mgrditchian T, Arakelian T, Paggetti J, Noman MZ, Viry E, Moussay E, et al. Targeting autophagy inhibits melanoma growth by enhancing NK cells infiltration in a CCL5-dependent manner. *P Natl Acad Sci USA*. 2017;114(44):E9271–E9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions