Current evidence and future perspectives of immune-checkpoint inhibitors in unresectable malignant pleural mesothelioma

Katsuyuki Hotta,¹ Nobukazu Fujimoto²

To cite: Hotta K, Fujimoto N. Current evidence and future perspectives of immune-checkpoint inhibitors in unresectable malignant pleural mesothelioma. *Journal for ImmunoTherapy of Cancer* 2020;**8**:e000461. doi:10.1136/jitc-2019-000461

Accepted 04 February 2020

ABSTRACT

Platinum-based chemotherapy is commonly used as the standard first-line treatment for unresectable malignant pleural mesothelioma (MPM). However, in recent times, immune-checkpoint inhibitors (ICIs) have led to a paradigm shift. Herein, we review relevant literature and ongoing trials of ICIs used as both first-line and salvage therapies. Specifically, in the Japanese single-arm, phase II trial, the MERIT trial, nivolumab, an antiprogrammed cell death 1 (PD-1) antibody showed favorable efficacy when used as a salvage therapy. Currently, multiple ICI monotherapy or combination therapy trials have been conducted, which could provide further evidence. Among available ICIs, the anti-PD-1 antibody is promising for unresectable MPM, despite the limited efficacy of anti-CTLA4 monotherapy. Ongoing studies will further confirm the potential efficacy of ICIs for MPM, as observed across other malignancies. It is also crucial to identify any clinically useful predictive biomarkers that could reveal ICIs with maximal effects in MPM.

INTRODUCTION

With increasing utilization of asbestos, the incidence of mesothelioma is considered to increase worldwide. Asbestos consumption in the USA has rapidly declined over the last 40 years, which has resulted in a considerable decline in mesothelioma incidence.¹ In Japan, the number of deaths had increased from 500 in 1995 to 1550 in 2016. Mesothelioma manifests mainly in the pleura, peritoneum and pericardium, although most commonly in the pleura.²

The major role of chronic inflammation and local tumor suppression in tumorigenesis observed in some experimental models led to the investigation of immunotherapy for malignant pleural mesothelioma (MPM).³ There have been intensive investigations on the efficacy and safety of immune-checkpoint inhibitors (ICIs) in the treatment of unresectable advanced diseases.^{4 5} Herein, we highlight relevant study results, as well as designs

and concepts of ongoing studies in both first-line and salvage settings.

Known biology

Among approximately 400 different mineral fibers present in nature, six fibers (amphiboles fibers (crocidolite, actinolite, tremolite, anthophyllite and amosite) and serpentine fiber (chrysotile)) are called as ‘asbestos’.⁶ They are carcinogenic and have been associated with mesothelioma.^{6 7} Furthermore, exposure of the chest to therapeutic ionizing radiation, usually performed to treat lymphomas, has been causally linked to mesothelioma, especially in young patients.^{8–10}

The accumulation of genetic aberrations can induce malignancies. Recently, The Cancer Genome Atlas program investigated genetic alterations in mesotheliomas using next-generation sequencing (NGS).¹¹ The results revealed frequent mutations in BAP1, CDKN2A, NF2, TP53, LATS2 and SETD2.^{11 12} Recently, a considerably higher number of genetic alterations in mesotheliomas has been detected than that detected by NGS, including point mutations, minute deletions and copy number changes.^{13 14} Furthermore, the vast array of genetic alterations in mesothelioma may lead to producing neoantigens, which correlate with the clonal expansion of tumor-infiltrating T lymphocytes.^{13 15} These findings suggest that, in contrast to the hypotheses based on NGS studies, mesothelioma may be immunogenic.¹⁵

Rationale for the development of immunotherapy

A hallmark of cancer is immune evasion, in which the immune system does not mount an effective antitumor response.¹⁶ Programmed cell death 1 (PD-1) is a negative costimulatory receptor expressed primarily on the surface of activated T cells^{17 18} and is involved in maintaining peripheral tolerance. The binding of PD-1 to one of its ligands, PD-L1 or PD-L2,

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Center for Innovative Clinical Medicine, Okayama University Hospital, Okayama, Japan

²Department of Medical Oncology and Medicine, Okayama Rosai Hospital, Okayama, Japan

Correspondence to

Dr Katsuyuki Hotta;
khotta@okayama-u.ac.jp

can inhibit a cytotoxic T-cell response.^{19 20} Tumors can co-opt this pathway to escape T-cell-induced antitumor activity.^{21–23}

The biology of MPM shows significant heterogeneity in both tumor and the microenvironment. Several studies, on T-cell-inhibitory receptors and chemokines, have indicated the prognostic role of lymphocytes and the occurrence of immunosuppression in MPM.^{24 25} In a melanoma model, PD-1 blockade increased the proportion of antigen-specific CTLs that recognized melanoma targets by degranulation, suggesting increased recognition efficiency for cognate peptide.²⁶ The increased frequency and absolute number of antigen-specific CTLs by PD-1 blockade resulted from augmented proliferation, and not decreased apoptosis. These findings have led to the extensive development of agents blocking immun checkpoints and their clinical investigation in various malignancies including MPM.

Biomarker in the ICI treatment of MPM

Some sensitive and specific immunohistochemistry markers including calretinin and WT1 are used for diagnosing mesothelioma.⁴ However, markers for treatment efficiency have not been established. Generally, PD-L1 expression level is used as the representative maker for predicting the efficacy of ICIs. In the ICI monotherapy with the salvage setting in non-squamous cell non-small-cell lung cancer, the PD-L1 expression level affected the survival efficacy,²⁷ while its influence was weakened when combined with platinum-based chemotherapy in the first-line setting.²⁸

In MPM, 20%–70% of the specimens tested are usually PD-L1 positive.²⁹ Such a wide range can be attributed to several factors. It could be because tumors are heterogeneous in nature.⁴ It could be partially attributed to the antibodies used; SP-263 is the most commonly used antibody,^{30–32} and the others include clones E1L3N and 28–8.³³ Furthermore, the histological subtype influences its frequency; PD-L1 expression is higher in non-epithelial mesotheliomas.³⁴ The cut-off levels of PD-L1 positivity vary among trials.³⁵ Considering that the positive rates were reported from different small studies with a small number of accrued patients, the data may be limited and actual rates of expression have hardly been studied. In addition to this, whether the ICI efficacy is truly dependent on the PD-L1 expression level is still controversial.

ICIs in the first-line settings

The standard treatment for unresectable, advanced malignant mesothelioma is chemotherapy, although with a very poor prognosis.³⁶ Similar to its use in non-small-cell lung cancer,^{37–44} cisplatin (CDDP) and pemetrexed (PEM) combination therapy (CDDP/PEM) approved by the US Food and Drug Administration (FDA) in 2004, is strongly recommended as the first-line treatment for mesothelioma.⁴⁵ Moreover, molecularly targeted agents have been developed to augment cytotoxic chemotherapy. For instance, a randomized phase III MAPS study showed

that adding bevacizumab to platinum doublets improved survival (HR of overall survival (OS) and progression-free survival (PFS): 0.77 (95% CI: 0.62 to 0.95); $p=0.0167$ and 0.61 (0.50 to 0.75); $p<0.0001$, respectively).⁴⁶ However, this regimen is yet to be approved by the FDA. A double-blind, randomized, placebo-controlled phase III study, the LUME-Meso trial of CDDP and PEM with or without nintedanib, a multikinase inhibitor for unresectable epithelioid MPM, showed that the primary endpoint, PFS, was not met.⁴⁷ Even with such an aggressive chemotherapy, OS for unresectable mesothelioma remains ≤ 12 months.⁴⁸

Given the limitations in the efficacy of existing cytotoxic chemotherapy in MPM and recent advances in tumor immunology across various malignancies, ICIs have been investigated for the treatment of unresectable mesothelioma. A single-arm, Durvalumab with First-line Chemotherapy in Mesothelioma study examined treatment efficacy after adding durvalumab, a PD-L1 inhibitor, to CPPD/PEM, in 54 patients with untreated, unresectable MPM⁴⁹ (table 1). PFS (the primary endpoint) at 6 months was 57%, and the objective response rate (ORR) was 48%, with a median duration of response of 6.5 months. Immune-related adverse events of grade 3 and higher, occurred in eight patients (15%), including lipase elevation ($n=1$), pancreatitis ($n=1$) and renal impairment ($n=1$).

The Canadian Cancer Trials Group has launched a phase II/III study for unresectable MPM, to verify treatment efficacy following the addition of pembrolizumab, a PD-1 antibody, to the standard CPPD/PEM (NCT02784171) (table 2). The use of durvalumab as the first-line immunotherapy is also under evaluation, sponsored by PrECOG (NCT02899195). Japanese investigators are also conducting an exploratory phase II trial, using nivolumab combined with the standard CPPD/PEM, in patients with untreated, unresectable MPM.⁵⁰ Furthermore, a large-scale, randomized phase III study, the CheckMate 743 study is currently investigating the survival advantage of the nivolumab/ipilimumab combination immunotherapy, versus platinum/PEM, in 606 patients with untreated, unresectable MPM (NCT02899299).

Single-agent ICI therapy in the salvage setting

Although the salvage setting is discussed before advancements in the first-line setting, currently available agents in the salvage setting rarely work in MPM, with a median survival time (MST) of ≤ 6 months.⁵¹ Vorinostat, a histone deacetylase inhibitor, was proven not to have any survival advantage in a placebo-controlled randomized phase III trial, the VANTAGE-014 trial,⁵² without earlier trial result confirmation.

Thus far, four ICIs have been tested as an immunotherapy against relapsed tumors (table 1). A single-center, single-arm phase II study, the NivoMes trial, with single-agent nivolumab, an anti-PD-1 antibody showed that 16 (47%) of the 34 registered patients with recurrent MPM achieved disease control at 12 weeks (8 with

Table 1 Relevant trial results

Trial name	Year	Phase	RCT	Drug	Primary endpoint	No	PS 0-1	No of sarcomatoid histology	ORR	mPFS (mo)	MST (mo)	Pneumonitis*	Ref.
Frontline setting													
DREAM			No	Durvalumab	PFS, OR	54	100%	–	48%	6.9	–	NR	49
Salvage setting													
<Single agent>													
MERIT	2018	2	No	Nivolumab	OR	34	100%	3 (9%)	29%	6.1	17.3	6%	54
NivoMes	2018	2	No	Nivolumab	DCR	34	100%	2 (6%)	24%	2.6	11.8	12%	53
KN-028†,‡	2017	1b	No	Pembrolizumab	Safety	25	100%	2 (8%)	20%	5.4	18	NR	55
Chicago group	2018	2	No	Pembrolizumab	OR	65	100%	5 (8%)	19%	4.5	11.5	3%	56
JAVELIN	2019	1b	No	Avelumab	OR	53	100%	2 (4%)	9%	4.1	10.7	6%	57
Italian group	2013	2	No	Tremelimumab	OR	29	79%	3 (10%)	7%	6.2	10.7	NR	58
Italian group	2015	2	No	Tremelimumab	irOR	29	79%	1 (3%)	3%	–	–	NR	59
DETERMINE	2017	2b	Yes	Tremelimumab	OS	382	99%	22 (6%)	5%	2.8	7.7§	NR	60
<Combination>													
NIBIT-MESO-1¶	2018	2	No	Tremelimumab/durvalumab	irOR	40	100%	2 (5%)	28%	5.7	16.6	NR	30
MAPS2	2019	2	Yes	Nivolumab/ipilimumab	DCR	62	98%	9 (15%)**	28%	5.6	15.9	2%	31
INITIATE	2019	2	No	Nivolumab/ipilimumab	DCR	63	97%	11 (17%)**	19%	4.0	11.9	2%	32

*Any grade.

†Those with the following conditions were eligible: (1) failed standard therapy and (2) unable to receive standard therapy.

‡Those with PD-L1-positive tumors were registered.

§OS-HR of 0.92 with a 95% CI 0.76 to 1.12.

¶Subjects who refused the first line platinum-based chemotherapy, or those with disease progression after a maximum of one line of platinum-based therapy, were eligible.

**Including biphasic histology.

DCR, disease control rate; DREAM, Durvalumab with First-Line Chemotherapy in Mesothelioma; irOR, immune-related objective response; MST, median survival time; NR, not reported; ORR, objective response rate; OS, overall survival; PD-L1, programmed cell death ligand 1; PFS, progression-free survival; RCT, randomized controlled trial.

Table 2 Ongoing relevant trials

Trial	Country	Phase	RCT	Regimen	Primary endpoint	No of planned pts	PS	Study start date	Registration no
Front-line setting									
Canadian group	Canada	2/3	Yes	Cis-pem±pembrolizumab	OS	126	0–1	07/10/16	NCT02784171
CM743	Global	3	Yes	Nivolumab/ipilimumab versus p-pem	OS	606	0–1	25/10/16	NCT02899299
PrE0505	USA	2	No	Cis-pem/durvalumab	OS	55	0–1	13/06/17	NCT02899195
JME-001	Japan	2	No	Cis-pem/nivolumab	OR	18	0–1	20/01/18	UMIN000030892
Salvage setting									
Confirm	UK	3	Yes	Nivolumab versus placebo	OS	336	0–1	28/03/17	NCT03063450

Cis-pem, cisplatin and pemetrexed; OS, overall survival; p-pem, platinum (cisplatin or carboplatin) and pemetrexed; PS, performance status; pts, patients; RCT, randomized controlled trial.

partial response (PR) and 8 with stable disease (SD)).⁵³ In this population, PD-L1 expression did not predict treatment responses. A Japanese single-arm phase II study, the MERIT study, also examined the efficacy and safety of nivolumab monotherapy in 34 patients with MPM with a history of prior chemotherapy.⁵⁴ The primary endpoint, ORR, was 29% (10/34), which was dependent on tumor PD-L1 expression, with an ORR of 40% and 8% when PD-L1 expression was $\geq 1\%$ and $< 1\%$, respectively. The median PFS and MST were 6.1 and 17.3 months, respectively. Twenty-six patients (76%) experienced treatment-related adverse events (TRAEs). In essence, these results led to the approval of nivolumab in Japan for unresectable recurrent pleural mesothelioma.

A single-agent pembrolizumab, anti-PD-1 antibody trial (KEYNOTE-028) demonstrated that 5/25 (20%) of previously treated patients with MPM achieved PR, while 13 (52%) had SD, with no treatment-related deaths or discontinuations.⁵⁵ The Chicago group also conducted a pembrolizumab monotherapy phase II trial in 65 patients with pretreated mesothelioma.⁵⁶ Nineteen per cent of the patients achieved PR, without unexpected AEs. The ORR was associated with PD-L1 expression; 7%, 26%, and 31% in patients harboring tumors with PD-L1-expression level of $< 1\%$, 1%–49% and $\geq 50\%$, respectively. The study also showed a median PFS and OS of 4.5 and 11.5 months, respectively.

With avelumab, a human anti-PD-L1 IgG₁ antibody, a phase Ib monotherapy trial (JAVELIN) was conducted in 53 patients with pretreated malignant mesothelioma.⁵⁷ Despite the 9% response in the whole cohort, ORR seemed different, stratified by the PD-L1 expression level in patients with PD-L1-positive (19% (3 of 16)) vs PD-L1-negative tumors (7% (2 of 27)), considering a $\geq 5\%$ PD-L1 cut-off. The median PFS was 4.1 months, whereas the MST extended to > 10 months. Five patients (9%) had grades 3–4 TRAEs, without treatment-related deaths.

Tremelimumab, an anti-CTLA4 antibody, was also evaluated in a salvage setting. In Europe, two single-arm, phase

II monotherapy trials showed preliminary efficacy, with an ORR of 3%–7%.^{58,59} Following these trials, a randomized phase IIb study, the DETERMINE study, revealed that tremelimumab failed to significantly prolong OS compared with that of placebo, in 571 patients with previously treated malignant mesothelioma. The MST showed no difference between treatment groups, with 7.7 and 7.3 months in the tremelimumab and placebo arms, respectively (HR 0.92, 95% CI 0.76 to 1.12).⁶⁰

ICI combination therapy in salvage settings

Given that enhanced immunogenicity can be achieved by combining PD1 or PDL1 and CTLA4 inhibitors,³ several studies evaluating the combination of anti-CTLA-4 and anti-PD-[L]1 antibodies have been reported. A phase II study, the NIBIT-MESO-1 trial, investigated an ICI combination of tremelimumab and durvalumab for unresectable mesothelioma.³⁰ Subjects who had refused first-line platinum-based chemotherapy, or subjects with disease progression after a maximum of one line of platinum-based therapy, were enrolled. Eleven (28%) of 40 patients had an immune-related objective response. The median PFS and MST were 5.7 and 16.6 months, respectively. Baseline tumor PD-L1 expression did not correlate with the immune-related objective response, and seven patients (18%) had grades 3–4 TRAEs.

A combination therapy of nivolumab and ipilimumab, over nivolumab monotherapy, was examined in a randomized phase II trial (IFCT MAPS2).³¹ A total of 125 patients with relapsed MPM were allocated to the combination therapy or monotherapy arm. Disease control rate (DCR), set as the primary endpoint, was 50% and 44%, whereas the ORR was 28% and 19%, respectively. As expected, the combination therapy had an increased risk of AE, with grades 3–4 of 26% and 14%, respectively. Three (5%) of 62 combination group patients had toxicities that led to death (hepatitis, encephalitis and acute kidney failure). When restricted to high PD-L1 tumors ($> 25\%$), either of

the regimens seemed effective, with ORRs of 63%–71% in the post hoc analyses.

Similar to this MAPS2 trial, a single-arm study, the INITIATE study,³² evaluated the efficacy of nivolumab and ipilimumab in mesothelioma refractory to at least one line of platinum-based chemotherapy. Of the 34 patients included in efficacy assessment, 10 (29%) attained PR and 13 (38%) attained SD, resulting in a DCR (primary endpoint) of 68%. Despite the smaller-scale, non-randomized design, this study could reproduce the tolerance and efficacy results obtained from the MAPS2 trial. It also showed a relationship between tumor PD-L1 expression and the efficacy of this combination therapy.

Based on the aforementioned completed trials, several MPM trials are either ongoing or being initiated. The most pivotal is the one initiated by Cancer Research UK: a randomized, double blind placebo controlled CONFIRM trial of nivolumab versus placebo in patients with relapsed mesothelioma (NCT03063450). A total of 336 patients will be recruited from 25 institutes in the UK over a 4-year period. All patients will be treated for 12 months, except in situations of progress or withdrawal. It will be intriguing if this reproduces the Japanese MERIT study results.⁵⁴

Overall, anti-PD-1 antibodies exhibited promising results when used alone as a salvage therapy after the first-line chemotherapy.^{53–56}

Unresolved, unmet needs for MPM ICI therapy

Compared with clinical trials targeting other malignancies, the majority of prior MPM trials employed ‘small-scale’ and ‘single-arm’ designs, and their primary endpoints were set at only ORR or DCR. No clear survival advantage of ICI has been demonstrated through randomized trials. This is mainly because of the extremely small patient population, and mostly exploratory-type trials.⁴ However, favorable responses and survival data could be observed across the studies, which are better than historical data. Considering the current limitations of treatment options in the salvage setting, ICI is now a potential rational and medically useful option for patients with unresectable, relapsed MPM, in the absence of any contraindications. Undoubtedly, well-designed randomized trials provide accurate and consistent data (ie, CONFIRM trial (NCT03063450); table 2). The accumulation of forthcoming relevant data through ongoing clinical trials is important for establishing better ICI use in daily practices.

Among toxicities induced by ICIs, pulmonary toxicity has to be properly managed, as it can be one of the most common causes of ICI-related death. The most common lung toxicity observed in patients receiving ICI treatment is pneumonitis.⁶¹ In our review, as shown in table 1, it occurred in 2%–12% of the patients (median; 6%) in all the trials evaluating ICIs. This seemed almost consistent with that observed in other cancers. The patterns of onset and severity may also vary, and MPM often has characteristics of limited reserve in pulmonary function at the

baseline. These findings suggest the importance of vigilance and rapid response. Thus, physicians still should recognize that the diagnosis of pneumonitis is particularly challenging and failure to detect and treat pneumonitis in a timely manner could lead to poor clinical outcomes.

Another unmet need is the identification of predictive biomarkers of ICI effects. Compared with other malignancies, progress in mesothelioma biomarker research is limited. Some of the single-arm ICI studies reveal the correlation between responses and higher PD-L1 expression. However, as insufficient survival data were generated, more established outcome data are needed to confirm the value of PD-L1 immunohistochemistry as a predictive biomarker for the OS effect. Recently, the tumor mutational burden (TMB) analysis using the whole exome sequence has garnered attention in nivolumab therapy.⁶² Moreover, in lung cancer, no association between TMB and PD-L1 expression was revealed.⁶² Rather, a combination of them would be of value as a predictive biomarker. Nevertheless, only a few precise biomarkers for ICI efficacy assessments seem to exist in MPM clinical trials, besides PD-L1 expression. Further development of new biomarkers is also required for unresectable mesothelioma.

A majority of patients diagnosed with untreated, unresectable mesothelioma exhibit all expected symptoms at the initial presentation, and thus, do not meet the eligibility criteria to participate in clinical trials. Therefore, study results have to be interpreted cautiously, taking into consideration how each of them can be applied per in-care patient, during daily clinical practices.

In the future, more novel immunotherapy results will be made available, which could possibly lead to further drastic changes in unresectable MPM treatment. Our goal is to carefully evaluate any relevant information and deliver better patient treatment.

CONCLUSIONS

MPM prognosis has been poor with the standard platinum chemotherapy. Recently, in the salvage setting, anti-PD-1 antibodies yielded favorable ORR. Nivolumab is approved for use in Japan. Ongoing studies will further confirm the potential efficacy of ICIs for MPM, as observed across other malignancies. It is also crucial to identify any clinically useful predictive biomarkers that could reveal the ICIs with maximal effects in MPM.

Contributors KH and NF carried out the search and assessment for relevant studies. KH drafted the manuscript. Both authors read and approved the final manuscript.

Funding This study was supported by grants-in-aid from the Ministry of Health, Labor and Welfare, Japan.

Competing interests KH has honoraria from AstraZeneca, Ono Pharmaceuticals, BMS, MSD, Eli Lilly Japan, Nihon Kayaku, Taiho Pharmaceuticals and Chugai Pharmaceuticals, as well as unrelated research funding from AstraZeneca, BMS, and Eli Lilly Japan. NF has received consultancy fees from Boehringer Ingelheim, Ono, Bristol-Myers Squibb, Kyorin, and Kissei, and honoraria or research funding from Ono, Boehringer Ingelheim, Bristol-Myers Squibb, Eli Lilly Japan, and MSD

related to this manuscript, as well as unrelated research funding from Hisamitsu, Chugai, Taiho, Novartis and GSK.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Henley SJ, Larson TC, Wu M, *et al.* Mesothelioma incidence in 50 states and the district of Columbia, United States, 2003-2008. *Int J Occup Environ Health* 2013;19:1-10.
- Ministry of the environment government of Japan. Available: <http://www.env.go.jp/air/asbestos/registration/> [Accessed 15 Aug 2019].
- Yap TA, Aerts JG, Popat S, *et al.* Novel insights into mesothelioma biology and implications for therapy. *Nat Rev Cancer* 2017;17:475-88.
- Carbone M, Adusumilli PS, Alexander HR, *et al.* Mesothelioma: scientific clues for prevention, diagnosis, and therapy. *CA Cancer J Clin* 2019;69:402-29.
- Forde PM, Scherpereel A, Tsao AS. Use of immune checkpoint inhibitors in mesothelioma. *Curr Treat Options Oncol* 2019;20:18.
- Baumann F, Ambrosi J-P, Carbone M. Asbestos is not just asbestos: an unrecognised health hazard. *Lancet Oncol* 2013;14:576-8.
- Carbone M, Kanodia S, Chao A, *et al.* Consensus report of the 2015 Weinman International Conference on mesothelioma. *J Thorac Oncol* 2016;11:1246-62.
- Goodman JE, Nascarella MA, Valberg PA. Ionizing radiation: a risk factor for mesothelioma. *Cancer Causes Control* 2009;20:1237-54.
- Vivero M, Bueno R, Chirieac LR. Clinicopathologic and genetic characteristics of young patients with pleural diffuse malignant mesothelioma. *Mod Pathol* 2018;31:122-31.
- Attanoos RL, Churg A, Galateau-Salle F, *et al.* Malignant mesothelioma and its non-asbestos causes. *Arch Pathol Lab Med* 2018;142:753-60.
- Hmeljak J, Sanchez-Vega F, Hoadley KA, *et al.* Integrative molecular characterization of malignant pleural mesothelioma. *Cancer Discov* 2018;8:1548-65.
- Bueno R, Stawiski EW, Goldstein LD, *et al.* Comprehensive genomic analysis of malignant pleural mesothelioma identifies recurrent mutations, gene fusions and splicing alterations. *Nat Genet* 2016;48:407-16.
- Mansfield AS, Peikert T, Smadbeck JB, *et al.* Neoantigenic potential of complex chromosomal rearrangements in mesothelioma. *J Thorac Oncol* 2019;14:276-87.
- Yoshikawa Y, Emi M, Hashimoto-Tamaoki T, *et al.* High-Density array-CGH with targeted NGS unmask multiple noncontiguous minute deletions on chromosome 3p21 in mesothelioma. *Proc Natl Acad Sci U S A* 2016;113:13432-7.
- Carbone M, Yang H, Goldino G. Does chromothripsis make mesothelioma an immunogenic cancer? *J Thorac Oncol* 2019;14:157-9.
- Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011;144:646-74.
- Sharpe AH, Freeman GJ. The B7-CD28 superfamily. *Nat Rev Immunol* 2002;2:116-26.
- Keir ME, Butte MJ, Freeman GJ, *et al.* PD-1 and its ligands in tolerance and immunity. *Annu Rev Immunol* 2008;26:677-704.
- Freeman GJ, Long AJ, Iwai Y, *et al.* Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med* 2000;192:1027-34.
- Dong H, Strome SE, Salomao DR, *et al.* Tumor-Associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion. *Nat Med* 2002;8:793-800.
- Schreiber RD, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in cancer suppression and promotion. *Science* 2011;331:1565-70.
- Vesely MD, Kershaw MH, Schreiber RD, *et al.* Natural innate and adaptive immunity to cancer. *Annu Rev Immunol* 2011;29:235-71.
- Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 2012;12:252-64.
- Ujii H, Kadota K, Nitadori J-I, *et al.* The tumoral and stromal immune microenvironment in malignant pleural mesothelioma: a comprehensive analysis reveals prognostic immune markers. *Oncoimmunology* 2015;4:e1009285.
- Chéné A-L, d'Almeida S, Blondy T, *et al.* Pleural effusions from patients with mesothelioma induce recruitment of monocytes and their differentiation into M2 macrophages. *J Thorac Oncol* 2016;11:1765-73.
- Wong RM, Scotland RR, Lau RL, *et al.* Programmed death-1 blockade enhances expansion and functional capacity of human melanoma antigen-specific CTLs. *Int Immunol* 2007;19:1223-34.
- Borghaei H, Paz-Ares L, Horn L, *et al.* Nivolumab versus docetaxel in advanced Nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015;373:1627-39.
- Gandhi L, Rodríguez-Abreu D, Gadgeel S, *et al.* Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N Engl J Med* 2018;378:2078-92.
- Combaz-Lair C, Galateau-Sallé F, McLeer-Florin A, *et al.* Immune biomarkers PD-1/PD-L1 and TLR3 in malignant pleural mesotheliomas. *Hum Pathol* 2016;52:9-18.
- Calabrò L, Morra A, Giannarelli D, *et al.* Tremelimumab combined with durvalumab in patients with mesothelioma (NIBIT-MESO-1): an open-label, non-randomised, phase 2 study. *Lancet Respir Med* 2018;6:451-60.
- Scherpereel A, Mazieres J, Greillier L, *et al.* Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial. *Lancet Oncol* 2019;20:239-53.
- Disselhorst MJ, Quispel-Janssen J, Lalezari F, *et al.* Ipilimumab and nivolumab in the treatment of recurrent malignant pleural mesothelioma (initiate): results of a prospective, single-arm, phase 2 trial. *Lancet Respir Med* 2019;7:260-70.
- Cedrès S, Ponce-Aix S, Zugazagoitia J, *et al.* Analysis of expression of programmed cell death 1 ligand 1 (PD-L1) in malignant pleural mesothelioma (MPM). *PLoS One* 2015;10:e0121071.
- Mansfield AS, Roden AC, Peikert T, *et al.* B7-H1 expression in malignant pleural mesothelioma is associated with sarcomatoid histology and poor prognosis. *J Thorac Oncol* 2014;9:1036-40.
- Nowak AK, McDonnell A, Cook A. Immune checkpoint inhibition for the treatment of mesothelioma. *Expert Opin Biol Ther* 2019;19:697-706.
- Mutti L, Peikert T, Robinson BWS, *et al.* Scientific advances and new frontiers in mesothelioma therapeutics. *J Thorac Oncol* 2018;13:1269-83.
- Hotta K, Matsuo K, Ueoka H, *et al.* Meta-Analysis of randomized clinical trials comparing cisplatin to carboplatin in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2004;22:3852-9.
- Hotta K, Matsuo K, Ueoka H, *et al.* Addition of platinum compounds to a new agent in patients with advanced non-small-cell lung cancer: a literature based meta-analysis of randomised trials. *Ann Oncol* 2004;15:1782-9.
- Hotta K, Matsuo K. Long-Standing debate on cisplatin- versus carboplatin-based chemotherapy in the treatment of advanced non-small cell lung cancer. *J Thorac Oncol* 2007;2:96.
- Hotta K, Fujiwara Y, Matsuo K, *et al.* Recent improvement in the survival of patients with advanced nonsmall cell lung cancer enrolled in phase III trials of first-line, systemic chemotherapy. *Cancer* 2007;109:939-48.
- Hotta K, Takigawa N, Hisamoto-Sato A, *et al.* Reappraisal of short-term low-volume hydration in cisplatin-based chemotherapy: results of a prospective feasibility study in advanced lung cancer in the Okayama lung cancer Study Group trial 1002. *Jpn J Clin Oncol* 2013;43:1115-23.
- Ninomiya K, Hotta K, Hisamoto-Sato A, *et al.* Short-Term low-volume hydration in cisplatin-based chemotherapy for patients with lung cancer: the second prospective feasibility study in the Okayama lung cancer Study Group trial 1201. *Int J Clin Oncol* 2016;21:81-7.
- Hotta K, Ninomiya K, Takigawa N, *et al.* Reappraisal of short-term low-volume hydration in cisplatin-based chemotherapy; hoping for it as a public domain. *Jpn J Clin Oncol* 2015;45:603-4.
- Nishii K, Hotta K, Ninomiya K, *et al.* Programmed cell death-ligand 1 expression and efficacy of cisplatin-based chemotherapy in lung cancer: a sub-analysis of data from the two Okayama lung cancer Study Group prospective feasibility studies. *Respir Investig* 2019;57:460-5.
- Vogelzang NJ, Rusthoven JJ, Symanowski J, *et al.* Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *J Clin Oncol* 2003;21:2636-44.

- 46 Zalcman G, Mazieres J, Margery J, *et al.* Bevacizumab for newly diagnosed pleural mesothelioma in the mesothelioma Avastin cisplatin pemetrexed study (maps): a randomised, controlled, open-label, phase 3 trial. *Lancet* 2016;387:1405–14.
- 47 Scagliotti GV, Gaafar R, Nowak AK, *et al.* Nintedanib in combination with pemetrexed and cisplatin for chemotherapy-naïve patients with advanced malignant pleural mesothelioma (LUME-Meso): a double-blind, randomised, placebo-controlled phase 3 trial. *Lancet Respir Med* 2019;7:569–80.
- 48 Tsao AS, Lindwasser OW, Adjei AA, *et al.* Current and future management of malignant mesothelioma: a consensus report from the National cancer Institute thoracic malignancy Steering Committee, International association for the study of lung cancer, and mesothelioma applied research Foundation. *J Thorac Oncol* 2018;13:1655–67.
- 49 Nowak A, Kok P, Lesterhuis W, *et al.* OA08.02 DREAM - A Phase 2 Trial of Durvalumab with First Line Chemotherapy in Mesothelioma: Final Result. *J Thorac Oncol* 2018;13:S338–9.
- 50 Fujimoto N, Aoe K, Kozuki T, *et al.* A phase II trial of first-line combination chemotherapy with cisplatin, pemetrexed, and nivolumab for unresectable malignant pleural mesothelioma: a study protocol. *Clin Lung Cancer* 2018;19:e705–7.
- 51 Kindler HL, Ismaila N, Armato SG, *et al.* Treatment of malignant pleural mesothelioma: American Society of clinical oncology clinical practice guideline. *J Clin Oncol* 2018;36:1343–73.
- 52 Krug LM, Kindler HL, Calvert H, *et al.* Vorinostat in patients with advanced malignant pleural mesothelioma who have progressed on previous chemotherapy (VANTAGE-014): a phase 3, double-blind, randomised, placebo-controlled trial. *Lancet Oncol* 2015;16:447–56.
- 53 Quispel-Janssen J, van der Noort V, de Vries JF, *et al.* Programmed death 1 blockade with nivolumab in patients with recurrent malignant pleural mesothelioma. *J Thorac Oncol* 2018;13:1569–76.
- 54 Okada M, Kijima T, Aoe K, *et al.* Clinical Efficacy and Safety of Nivolumab: Results of a Multicenter, Open-label, Single-arm, Japanese Phase II study in Malignant Pleural Mesothelioma (MERIT). *Clin Cancer Res* 2019;25:5485–92.
- 55 Alley EW, Lopez J, Santoro A, *et al.* Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1B trial. *Lancet Oncol* 2017;18:623–30.
- 56 Desai A, Karrison T, Rose B, *et al.* Phase II trial of pembrolizumab (NCT02399371) in previously treated malignant mesothelioma: final analysis, in IASLC (ED). 19th IASLC World Conference on Lung Cancer Toronto, Canada, 2018.
- 57 Hassan R, Thomas A, Nemunaitis JJ, *et al.* Efficacy and safety of avelumab treatment in patients with advanced unresectable mesothelioma: phase 1B results from the javelin solid tumor trial. *JAMA Oncol* 2019;5:351–7.
- 58 Calabrò L, Morra A, Fonsatti E, *et al.* Tremelimumab for patients with chemotherapy-resistant advanced malignant mesothelioma: an open-label, single-arm, phase 2 trial. *Lancet Oncol* 2013;14:1104–11.
- 59 Calabrò L, Morra A, Fonsatti E, *et al.* Efficacy and safety of an intensified schedule of tremelimumab for chemotherapy-resistant malignant mesothelioma: an open-label, single-arm, phase 2 study. *Lancet Respir Med* 2015;3:301–9.
- 60 Maio M, Scherpereel A, Calabrò L, *et al.* Tremelimumab as second-line or third-line treatment in relapsed malignant mesothelioma (determine): a multicentre, international, randomised, double-blind, placebo-controlled phase 2B trial. *Lancet Oncol* 2017;18:1261–73.
- 61 Puzanov I, Diab A, Abdallah K, *et al.* Managing toxicities associated with immune checkpoint inhibitors: consensus recommendations from the Society for immunotherapy of cancer (SITC) toxicity management Working group. *J Immunother Cancer* 2017;5:95.
- 62 Hellmann MD, Ciuleanu T-E, Pluzanski A, *et al.* Nivolumab plus ipilimumab in lung cancer with a high tumor mutational burden. *N Engl J Med* 2018;378:2093–104.