Evolving impact of long-term survival results on metastatic melanoma treatment

Olivier Michielin,^{1,2} Michael B Atkins ,³ Henry B Koon,⁴ Reinhard Dummer,⁵ Paolo Antonio Ascierto ⁶

To cite: Michielin O, Atkins MB, Koon HB, *et al.* Evolving impact of long-term survival results on metastatic melanoma treatment. *Journal for ImmunoTherapy of Cancer* 2020;**8**:e000948. doi:10.1136/jitc-2020-000948

Accepted 19 August 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Oncology Department, Precision Oncology Center, Lausanne, Switzerland

²Oncology Department, Lausanne University Hospital, Lausanne, Switzerland

³Medical Oncology, Georgetown Lombardi Comprehensive Cancer Center and Oncology Academic Department, Georgetown University Medical Center, Washington, DC, USA

⁴Clinical Research, Bristol Myers Squibb, Princeton, New Jersey, USA

⁵Dermatology, Universitäts Spital, Zurich, Switzerland

⁶Melanoma Cancer Immunotherapy and Innovative Therapy Unit, Istituto Nazionale Tumori IRCCS Fondazione Pascale, Naples, Italy

Correspondence to

Dr Paolo Antonio Ascierto; paolo.ascierto@gmail.com

ABSTRACT

Melanoma treatment has been revolutionized over the past decade. Long-term results with immuno-oncology (I-O) agents and targeted therapies are providing evidence of durable survival for a substantial number of patients. These results have prompted consideration of how best to define long-term benefit and cure. Now more than ever, oncologists should be aware of the long-term outcomes demonstrated with these newer agents and their relevance to treatment decision-making. As the first tumor type for which I-O agents were approved, melanoma has served as a model for other diseases. Accordingly, discussions regarding the value and impact of long-term survival data in patients with melanoma may be relevant in the future to other tumor types. Current findings indicate that, depending on the treatment, over 50% of patients with melanoma may gain durable survival benefit. The best survival outcomes are generally observed in patients with favorable prognostic factors, particularly normal baseline lactate dehydrogenase and/or a low volume of disease. Survival curves from melanoma clinical studies show a plateau at 3 to 4 years, suggesting that patients who are alive at the 3-year landmark (especially in cases in which treatment had been stopped) will likely experience prolonged cancer remission. Quality-of-life and mixture-cure modeling data, as well as metrics such as treatment-free survival, are helping to define the value of this long-term survival. In this review, we describe the current treatment landscape for melanoma and discuss the long-term survival data with immunotherapies and targeted therapies, discussing how to best evaluate the value of long-term survival. We propose that some patients might be considered functionally cured if they have responded to treatment and remained treatment-free for at least 2 years without disease progression. Finally, we consider that, while there have been major advances in the treatment of melanoma in the past decade, there remains a need to improve outcomes for the patients with melanoma who do not experience durable survival.

INTRODUCTION

Despite intensive efforts, the goal of curing cancer has proven to be elusive for most advanced solid tumors, including melanoma. To achieve this goal, an in-depth

understanding of melanoma biology, tumor development, and the body's innate defense against tumor progression is required. Advances in our understanding of melanoma have facilitated the development and subsequent approval of various novel cancer therapies with the ability to substantially improve outcomes.

Melanoma is the paradigm of how the treatment landscape can be revolutionized over a decade. Since 2011, a series of novel immuno-oncology (I-O) agents, targeted therapies, and combination regimens have been approved.¹ Robust data from phase III trials^{2–7} and subsequent real-world evidence^{8–10} have unequivocally demonstrated overall survival (OS) and progression-free survival (PFS) benefits with these agents. As the follow-up of pivotal trials continues, a substantial body of long-term data is emerging that supports the durability of survival with some agents.^{11–17} These data are altering practice in ways almost inconceivable a decade ago, raising new clinical questions and challenges.

In this review, we consider the current treatment landscape for metastatic melanoma, including the following questions: Are patients with sustained benefit actually cured? Can treatment and/or follow-up of these patients be safely stopped, allowing a return to their lives before diagnosis? How is the value of long-term survival best assessed, and can these metrics guide treatment choice? Does the demonstration of durable outcomes for patients treated in the metastatic setting impact the role of local or regional therapies? How can outcomes be improved for patients with melanoma who do not experience durable survival? Because melanoma was the first tumor type for which I-O agents were approved, it has served as a guiding example for other tumors; therefore, these discussions may

prove to be informative for the treatment of different tumor types in the future.

DISCUSSION

The evolution of melanoma treatment

Supported by innovative research tools and techniques, our understanding of tumor biology has improved, particularly in regard to the mechanisms tumors use to escape detection by the immune system. Fundamental to greater understanding has been the development of the immunoediting theory, which describes the interaction between the tumor and the immune system as a three-step process (referred to as the three Es): (1) elimination of tumors at an early stage (also known as the immune surveillance hypothesis); (2) equilibrium (when the immune system controls the tumor); and (3) escape (when tumor cells are fully immunoedited and grow without immune control). The three Es are now accepted as a template for understanding the interactions of cancer cells with the host immune system.¹⁸ This concept has important therapeutic consequences because it implies that fully developed tumors are inherently immune-resistant. This has shifted our therapeutic strategies away from stimulating the immune system by vaccination, interleukin (IL)-2, or other strategies and toward counteracting immune escape mechanisms. The first agents were designed to block the inhibitory immune checkpoint receptors cytotoxic T-lymphocyte associated antigen-4 (CTLA-4) and programmed death 1 (PD-1). CTLA-4 is expressed on T cells where it acts to regulate the magnitude of the early, priming stages of T-cell activation. Once the T-cell receptor binds to an antigen, cluster of differentiation (CD)28, a co-stimulatory receptor, amplifies the signal to activate the T cells. CD28 and CTLA-4 share ligands (CD80 (B7.1) and CD86 (B7.2)), for which CTLA-4 has a much greater affinity than CD28. Therefore, expression of CTLA-4 on the T-cell surface outcompetes CD28 for ligand binding, delivering inhibitory signals to the T cell and reducing activation.¹⁹ Unlike CTLA-4, the primary role of PD-1 is to limit the activity of T cells during the effector phase of the immune response in the peripheral tissue, limiting autoimmunity. The engagement of PD-1 by its ligands, programmed death-ligand 1 (PD-L1) and PD-L2, results in an inhibitory signal that reduces T-cell proliferation, cytotoxicity, and cytokine production.¹⁹ The PD-1 and CTLA-4 pathways are distinct, acting at different times and locations to regulate T-cell activation.²⁰ In tumors, inhibiting these pathways has different effects: blocking CTLA-4 induces de novo antitumor T-cell responses, while PD-1 inhibition restores antitumor T-cell function.^{19–23} These distinct but potentially complementary effects led to the clinical evaluation of immune checkpoint inhibitor-based combinations.

While immune checkpoint inhibitors are now a mainstay of treatment for patients with metastatic melanoma, other therapeutic approaches designed to modulate a patient's immune system have also been developed. Adoptive

Figure 1 US Food and Drug Administration approval of melanoma therapies.^{31 32} Agents shown are approved for metastatic melanoma unless stated and italicized. ^aFor patients whose tumors express *BRAF* V600E or V600K. ^bFor patients whose tumors express *BRAF* V600E.

cell transfer involves the infusion of autologous tumor-infiltrating lymphocytes that have been expanded in vitro with IL-2.²⁴ Development of this approach began in the 1980s and responses of long duration were first achieved with the use of lymphoid depletion prior to adoptive cell transfer.^{25 26} Subsequently, various regimens and production processes have subsequently been evaluated in larger clinical trials. Findings from a recent meta-analysis showed that, especially when combined with high-dose IL-2, adoptive cell transfer achieves durable clinical benefit in a substantial minority of patients; however, the complex production process means that this approach is currently used in a small, but expanding number of treatment centers worldwide.²⁴ Vaccination is a strategy to expand tumor-directed lymphocytes in vivo. Melanoma vaccines have a long history, with numerous approaches having been investigated: the majority of studies have evaluated melanoma vaccines used alone and have not demonstrated clinical efficacy.^{27–30} In 2015, talimogene laherparepvec (TVEC), an oncolytic virotherapy, was approved by the US Food and Drug Administration (FDA) for the local treatment of unresectable lesions in patients with melanoma recurrence after initial surgery (figure 1).^{31 32} It has been speculated that the presence of this genetically modified virus in the melanoma cells

results in an individualized vaccine. Approval was based on the results of a randomized phase III trial showing significantly improved durable response rates with TVEC versus subcutaneous granulocyte-macrophage colony-stimulating factor in patients with unresectable stage IIIB to stage IV melanoma.³³ The final analysis from this trial showed a 33% OS rate at 5 years.³⁴ A number of other vaccines are under evaluation for the treatment of melanoma and may play a role in combination therapy with immune checkpoint inhibitors and/or targeted therapies in the future.³⁵

At the same time while research into immune checkpoint receptors was being pursued, various tumorigenic mutations were discovered. In particular, *BRAF*-activating mutations were identified in more than 50% of melanomas and found to play a critical role in driving tumor growth and development. This led to the development of vemurafenib, dabrafenib, and encorafenib which target oncogenic activating mutations in the MAP kinase (MAPK) pathway, and trametinib, cobimetinib, and binimetinib, which target MEK.^{31, 36} Clinical experience has demonstrated that the majority of patients treated with a *BRAF* inhibitor alone developed resistant disease, primarily involving reactivation of the MAPK pathway.³ Some patients also developed secondary malignancies, such as cutaneous squamous-cell carcinoma, attributed to paradoxical *BRAF*-inhibitor-induced activation of the MAPK pathway in *RAS*-mutant tumors. Further research has shown that combined *BRAF* and MEK inhibition delayed the emergence of resistance and reduced the incidence of cutaneous hyperproliferative lesions compared with single-agent *BRAF* inhibition, which led to the clinical evaluation of combination regimens that demonstrated improved therapeutic efficacy compared with monotherapy.³

In 2011, the FDA approved the first immune checkpoint inhibitor, ipilimumab, and the first targeted therapy, vemurafenib, for melanoma.³¹ Subsequently, several other agents have been approved, based on the demonstration of significant OS benefit in randomized trials.^{1, 31} These agents were initially approved for use as monotherapy (figure 1);^{31, 32} however, strong scientific and clinical rationale for combination therapy resulted in the subsequent approval of checkpoint-inhibitor-based and targeted therapy-based combination regimens. More recently, several adjuvant treatments for patients with fully resected stage III or stage IV disease have been approved (figure 1).^{31, 37–40}

Metastatic melanoma treatment today

The newer therapies for metastatic melanoma have undoubtedly improved outcomes for many patients, but they have also created new treatment selection challenges for both patients and oncologists. This is particularly true for patients with *BRAF*-mutant melanoma, for which both I-O agents and targeted therapies are approved.

For patients with *BRAF*-mutant melanoma, robust data have demonstrated a significant benefit for combination targeted therapy over *BRAF*-inhibitor monotherapy; so, when a targeted therapy is chosen, combination *BRAF*/MEK inhibitor treatment is preferred for most

patients.^{3, 4, 6, 7, 41} When considering treatment with an I-O agent, note that data from randomized trials has clearly shown that a PD-1 inhibitor is superior to ipilimumab (a CTLA-4 inhibitor) in the first-line setting in terms of both efficacy and safety, regardless of tumor *BRAF*-mutation status.^{2, 5, 41, 42} In the absence of head-to-head studies, the choice between the approved PD-1 inhibitors (nivolumab or pembrolizumab) is often based on factors such as treatment schedule, institutional formulary options, and the preference of the treating oncologist. However, as long-term data become available, they may also inform decision-making (table 1).^{5, 11–17, 25, 43–46}

When deciding whether to prescribe nivolumab plus ipilimumab combination therapy or PD-1-inhibitor monotherapy, consider that descriptive data from the CheckMate 067 trial showed superior median OS with the combination versus nivolumab alone (>60 months (median not reached) versus 36.9 months; HR 0.83); this difference was maintained with up to 5 years of follow-up.¹² The HR for benefit with the combination over nivolumab was particularly robust in patients with *BRAF*-mutant disease (0.70; 5-year OS rate 60% vs 46%), elevated (ie, two times the upper limit of normal) lactate dehydrogenase (LDH; 0.73; 5-year OS rate 28% vs 14%), or M1c stage per American Joint Committee on Cancer, seventh edition (0.82; 5-year OS rate 43% vs 35%), respectively. The frequency and severity of treatment-related adverse events (AEs), however, was higher with the combination than with nivolumab alone (any grade 96% vs 87%; grade 3/4 59% vs 23%).¹² Consequently, the treatment decision is complex and multifactorial, involving careful consideration and discussion about the risks and potential benefits of the treatment options and, in our experience, with patient preference often being an important factor.

The choice between I-O and targeted therapy for treatment-naïve patients with *BRAF*-mutant disease is difficult because of the lack of data from head-to-head trials. Most clinicians rely on practical experience, real-world and trial data, and expert-opinion publications. However, health economic/quality-of-life (QoL) data and outcomes from other analyses, such as data modeling, may help to inform decision-making. For example, in a recent matching-adjusted indirect comparison, clinical outcomes with nivolumab plus ipilimumab were compared with dabrafenib plus trametinib or vemurafenib plus cobimetinib combination treatments in patients with *BRAF*-mutant advanced melanoma.⁴⁷ After adjusting for differences in baseline characteristics, PFS and response outcomes were similar among the treatments; but, beyond 12 months of treatment, nivolumab plus ipilimumab improved OS compared with targeted therapy. This suggests that the survival benefit with combination immunotherapy emerges later and may be more durable than with targeted therapies. The incidences of AEs (any-grade, grade 3/4) were lowest with dabrafenib plus trametinib (89.5%, 31.6%) and similar between vemurafenib plus cobimetinib (97.6%, 59.5%) and nivolumab plus ipilimumab (94.6%, 54.1%), although the nature of the

Table 1 Long-term survival in clinical trials with I-O agents or targeted therapies in patients with advanced/metastatic melanoma

Study	Phase	Patient population*	Duration of survival follow-up, months	Arm† (n), dosing schedule	Median PFS, months (95% CI)	Longest landmark PFS rate	Median OS, months (95% CI)	Longest landmark OS rate
Immune checkpoint inhibitors								
CheckMate 067 ¹²	III	Any <i>BRAF</i>	60 (minimum)	NIVO+IPI (314), NIVO 1 mg/kg+IPI 3 mg/kg Q3W×4, then NIVO 3 mg/kg Q2W	11.5 (8.7 to 19.3)	5 years: 36%	NR (38.2 to NR)	5 years: 52%
				NIVO (316), NIVO 3 mg/kg Q2W	6.9 (5.1 to 10.2)	5 years: 29%	36.9 (28.2 to 58.7)	5 years: 44%
				IPI (315), IPI 3 mg/kg Q3W×4	2.9 (2.8 to 3.2)	5 years: 8%	19.9 (16.8 to 24.6)	5 years: 26%
CheckMate 066 ¹⁶	III	<i>BRAF</i> wild-type	60 (minimum)	NIVO (210), NIVO 3 mg/kg Q2W	5.1 (3.5 to 12.2)	5 years: 28%	37.3 (25.4 to 51.6)	5 years: 39%
KEYNOTE-006 ⁵	III	Any <i>BRAF</i> Data shown are for the treatment-naïve patient subgroup	57.7 (median)	PEMBRO (368), PEMBRO 10 mg/kg Q2W/Q3W	11.6 (8.2 to 16.4)	4 years: 26% (Q2W), 28% (Q3W)	38.7 (27.3 to 50.7)	5 years: 41% (Q2W), 45% (Q3W)
				IPI (181), IPI 3 mg/kg Q3W×4	3.7 (2.8 to 4.3)	NA	17.1 (13.8 to 26.2)	NA
CA184-169 ⁴⁴	III	Any <i>BRAF</i>	43 (minimum)	IPI 10 mg/kg (365), IPI 3 mg/kg (362), Q3W×4	2.8 (2.8 to 3.0), 2.8 (2.8 to 2.8)	NA	15.7 (11.6 to 17.8), 11.5 (9.9 to 13.3)	3 years: 31%, 23%
KEYNOTE-001 ¹¹	I	Any <i>BRAF</i> Data shown are for the treatment-naïve patient subgroup	55 (median)	PEMBRO (151), PEMBRO 2 mg/kg Q3W/10 mg Q2W/10 mg Q3W	16.9 (9.3 to 35.5)	5 years: 29%	38.6 (27.2 to NR)	5 years: 41%
CA209-004 ⁴⁵	I	Any <i>BRAF</i> No prior immune checkpoint inhibitor	43.1 (median)	NIVO+IPI (94), Dose-escalation study†	NA	NA	NR (40.9 to NR)	4 years: 57%
CA209-003 ¹⁷	I	Previously treated, advanced melanoma (any <i>BRAF</i> , <i>RCC</i> , or <i>NSCLC</i> (only melanoma data shown))	58.3 (minimum)	NIVO (107), Dose escalation 0.1 to 10 mg/ kg Q2W	NA	NA	20.3 (12.5 to 37.9)	5 years: 34%
IPI-pooled analysis ⁴⁶	II/III	Various	Approx. 11 (median)	IPI (1861), Most patients received 3 or 10 mg/kg	NA	NA	11.4 (10.7 to 12.1)	3 years: 22%
Other immunotherapy								
IL-2 pooled analysis ¹⁹	Various	Various	62 (median)	High-dose IL-2 (270), 600,000 or 720,000 IU/kg	NA	NA	11.4 (NA)	NA

Continued

Table 1 Continued

Study	Phase	Patient population*	Duration of survival follow-up, months	Arm† (n), dosing schedule	Median PFS, months (95% CI)	Longest landmark PFS rate	Median OS, months (95% CI)	Longest landmark OS rate
ACT+IL-2 ²⁵	Various	Previously treated, advanced melanoma	62 (median)	Autologous tumor-infiltrating lymphocytes+IL-2 after a lymphodepletion	NA	NA	NA	5 years: 29%
Combination-targeted therapy								
COMBI-d and COMBI-v ¹⁵	III	BRAF-mutant	22 (median)	DAB +TRAM (563), DAB 150 mg twice daily+TRAM 2 mg once daily	11.1 (9.5 to 12.8)	5 years: 19%	25.9 (22.6 to 31.5)	5 years: 34%
COLUMBUS ¹³	III	BRAF-mutant	48.8 (median)	ENCO+BINI (192), ENCO 450 mg once daily+BINI 45 mg twice daily	14.9 (11.0 to 20.2)	3 years: 29%	33.6 (24.4 to 39.2)	3 years: 47%
				VEM (191), VEM 960 mg twice daily	7.3 (5.6 to 7.9)	3 years: 14%	16.9 (14.0 to 24.5)	3 years: 31%
				ENCO (194), ENCO 300 mg once daily	9.6 (7.4 to 14.8)	3 years: 25%	23.5 (19.6 to 33.6)	3 years: 41%
				COBI+VEM (247), COBI 60 mg once daily+VEM 960 mg twice daily	12.6 (9.5 to 14.8)	5 years: 14%	22.5 (20.3 to 28.8)	5 years: 31%
coBRIM ¹⁴	III	BRAF-mutant	NA	VEM (248), VEM 960 mg twice daily	7.2 (5.6 to 7.5)	5 years: 10%	17.4 (14.5 to 19.8)	5 years: 26%

*Trials were conducted solely in treatment-naïve patients unless indicated.

†Data for control arms are only included if ipilimumab or a targeted therapy was the comparator.

‡NIVO+IPI Q3W×4 (mg/kg) in one of the following cohorts: (1) NIVO 0.3+IPI 3; (2) NIVO 1+IPI 3; (2a) NIVO 3+IPI 3; (8) NIVO 1+IPI 3. Cohorts 1 to 3 received maintenance with NIVO Q3W×4, then NIVO+IPI Q12W×8.

ACT, adoptive cell transfer; BINI, binimetinib; COBI, cobimetinib; DAB, dabrafenib; ENCO, encorafenib; IL-2, interleukin-2; I-O, immuno-oncology; IPI, ipilimumab; NA, not available; NIVO, nivolumab; NR, not reached; NSCLC, non-small cell lung cancer; OS, overall survival; PEMBRO, pembrolizumab; PFS, progression-free survival; Q2W, every 2 weeks; Q3W, every 3 weeks; RCC, renal cell carcinoma; TRAM, trametinib; VEM, vemurafenib.

Figure 2 Long-term OS in clinical trials with immuno-oncology agents and targeted therapies in patients with advanced melanoma. The data presented represent first-line treatment options, with the exception of those for IL-2 and pooled ipilimumab. The data were compiled from the references indicated. IL-2, interleukin 2; OS, overall survival.

toxicities is very different between immune checkpoint inhibitors and targeted therapies.⁴⁸

Several randomized trials have been ongoing to determine the most effective sequence and administration of I-O and targeted therapies for all comers and biomarker-defined subsets of treatment-naïve patients with *BRAF*-mutant melanoma.^{49–51} The randomized phase II SECOMBIT trial (NCT02631447)⁴⁹ was designed to determine the best sequential approach with ipilimumab plus nivolumab and encorafenib plus binimetinib. In the phase III DREAMseq trial (NCT02224781),⁵⁰ patients will be receiving ipilimumab and nivolumab followed by dabrafenib and trametinib or the reverse sequence at the time of documented disease progression. The phase II EBIN trial (NCT03235245)⁵¹ was designed to compare encorafenib plus binimetinib followed by nivolumab plus ipilimumab versus nivolumab plus ipilimumab. The phase II ImmunoCobiVem trial (NCT02902029)⁵² was designed to evaluate cobimetinib plus vemurafenib followed by a programmed death ligand 1 (PD-L1) inhibitor.

Long-term survival outcomes

As patients are followed for 5 years and beyond in the pivotal trials with I-O agents and targeted therapies, long-term data are becoming available and provide new metrics that can contribute to treatment decisions. Available long-term data are summarized in [table 1](#) and [figure 2](#).^{5 11–17 25 43–46} Because these studies differed in their patient populations and other parameters, it is not appropriate to compare the trials directly. Additionally, different trial start dates may have impacted the patient populations enrolled and the subsequent therapies received. However, when assessing IL-2 or ipilimumab treatment, the improvement in clinical outcomes

in patients treated with PD-1-inhibitor monotherapy, targeted therapy, or nivolumab plus ipilimumab becomes clear.

Across studies, the best survival outcomes have generally been observed in patients with favorable prognostic factors, particularly a normal or low LDH level and/or a low volume of disease. In a pooled survival analysis of dabrafenib and trametinib (in patients with *BRAF*-mutant tumors), the 5-year survival rate was 43% in patients with LDH at or below the upper limit of normal versus 34% in the general population.¹⁵ The highest OS rate, 55%, was found in patients with low/normal LDH and fewer than three sites of disease. In CheckMate 067, there was a similar trend for better survival in patients with favorable prognostic factors: the highest 5-year survival rates with nivolumab plus ipilimumab were observed in patients with normal LDH (60%) or normal LDH with fewer than three sites of disease (64%), compared with 52% in all patients (the 5-year survival rate was 60% for patients with *BRAF*-mutant tumors).¹²

When long-term data with immune checkpoint inhibitors became available in a pooled analysis of ipilimumab trials in which some patients were followed for more than 10 years,⁴⁶ attention was drawn to the shape of the OS curves, specifically the plateau or flattening of the tail of the curve. As shown in [figure 2](#), the immune checkpoint inhibitor and IL-2 curves were similar, plateauing at 3 or 4 years. The OS curves for targeted therapies also appeared to plateau. However, the contributions of patients receiving a subsequent therapy were unclear. Based on current data, the plateaus for the I-O and the targeted agents are expected to be sustained with longer follow-up. This leads to the question of whether the

patients represented in the tails of these curves can be considered to be functionally cured.

OS curves are known to be impacted by subsequent therapies. Many patients treated with BRAF plus MEK inhibitor combinations who are still alive may have received, or are still receiving, subsequent therapy with immune checkpoint inhibitors; patients in immune checkpoint inhibitor trials may have received subsequent BRAF plus MEK inhibitor combinations. In the pooled analysis of the COMBI-d and COMBI-v trials, of the 161 patients who were alive at 5 years, 72 (45%) had received further treatment, the most common being immunotherapy (52 patients, 78%).¹⁵ In the CheckMate 067 trial, of the 281 patients in the nivolumab-containing arms who were alive at 5 years, 58 (21%) had received subsequent therapy.¹² Data were similar for patients with BRAF-mutant melanoma: of the 97 patients who were alive at 5 years, 25 (26%) had received subsequent therapy.⁵³ To fully evaluate the contribution of an agent to OS, it may be helpful to consider the tail of the PFS curve. PFS offers an alternative to OS that is available in a timely manner and is not confounded by subsequent therapies or causes of mortality unrelated to cancer.^{54 55} In CheckMate 067 and the COMBI-d and COMBI-v pooled analyses, the PFS curves appeared to flatten after approximately 3 years. In CheckMate 067, PFS rates for the overall population at 3, 4, and 5 years were 39%, 37%, and 36% with nivolumab plus ipilimumab, and 32%, 31%, and 29% with nivolumab alone.^{12 56 57} PFS rates for patients with BRAF-mutant melanoma at 3, 4, and 5 years were 40%, 39%, and 38% with nivolumab plus ipilimumab, and 22%, 23%, and 22% with nivolumab alone. In COMBI-d and COMBI-v, PFS rates were 24% at 3 years, 21% at 4 years, and 19% at 5 years.¹⁵ In the KEYNOTE-006 trial, PFS rates for patients who received either dose of pembrolizumab as a first-line therapy declined from 37% at 2 years, to 33% at 3 years, and 27% at 4 years; the lack of data beyond 4 years makes it unclear whether the curve continued to flatten.⁵

Defining cure in advanced cancer

Defining cure has not been necessary for most advanced cancers. In 1985, Schipper and colleagues proposed the term 'functional cure,' acknowledging that, given the biology of cancer, eradicating all cancer cells or proving that they had been eradicated may not be possible, and seeking to gain long-term control may be more realistic.⁵⁸ Based on current long-term data, patients could be regarded as functionally cured after 3 or 4 years when the OS curve plateaus; however, at that time, depending on the agent, some patients may still be receiving treatment. For example, in CheckMate 067, stopping nivolumab after a specific time period had not been mandated and, at 5 years, 13% of the patients in the nivolumab-containing arms who were alive were still receiving study treatment. However, the median duration of treatment was short in both arms (3.6 months (range, 0.0 to 57.0) with nivolumab plus ipilimumab combination therapy and 7.6 months (range, 0.0 to 62.9) with nivolumab monotherapy).⁵⁹ In

contrast, in the COMBI-d and COMBI-v pooled analysis, at 5 years, 43% of the patients who were alive were still on study treatment.¹⁵ Furthermore, one-half of the patients with complete response (CR) on these medications were shown to relapse at median of 6.6 months after treatment was stopped.⁶⁰ Findings from a retrospective analysis of the long-term outcomes of 396 patients treated with a PD-1 inhibitor at a single institution showed that most patients discontinued treatment at the time of CR. Most CRs were durable, but the probability of treatment failure 3 years after best overall response was 27%.⁶¹

The appropriate time to stop treatment has been the subject of much debate, particularly because different agents were developed with different stopping rules. For example, in the pembrolizumab KEYNOTE-006 trial, patients received up to 2 years of treatment⁵; whereas, in the CheckMate 067 trial, patients continued until progression, unacceptable toxicity, or withdrawal of consent.⁵⁷ If a maximum duration of therapy has not been mandated, the decision to stop may be supported by a biopsy showing either the absence of viable tumor or negative ¹⁸F-2-fluoro-2-deoxy-D-glucose-positron emission tomography (FDG-PET) imaging. An FDG-PET scan, which provides functional information alongside standard anatomical imaging, may help to determine whether residual lesions in patients with a partial response (PR) or stable disease (SD) contain any viable tumor, thereby identifying patients achieving a complete metabolic response.⁶² Results of a study designed to determine whether FDG-PET imaging is a better predictor of long-term outcomes than a standard CT scan showed that, after 1 year of immune checkpoint inhibitor treatment, 28% of patients had achieved CR on CT according to Response Evaluation Criteria in Solid Tumors V.1.1.⁶² In comparison, 68% of patients with a PR on CT had a complete metabolic response according to FDG-PET imaging. Almost all of the patients with a complete metabolic response at 1 year had an ongoing response to therapy thereafter. Acknowledging that longer follow-up is needed, these findings suggest that FDG-PET imaging could help to predict long-term survival and guide the decision to stop therapy. Other studies are planned to investigate the ideal duration of therapy for a specific subset of patients. The subject of how long to treat patients requires data, discussion, and development of best-practice guidelines, along with the identification of surrogate markers that indicate when stopping treatment is safe.

A further consideration is whether a documented radiologic CR is necessary for a patient to be considered functionally cured. Given the concept of a functional cure and the mechanism of action of I-O agents in particular, the long-term absence of progression rather than the documentation of a CR may be a more appropriate metric. Long-term data suggest that achieving a CR is not necessary for a sustained survival benefit, although patients who achieve a CR have more durable benefit over time.^{15 53 63} In the COMBI-d and COMBI-v pooled analysis, the 5-year OS rates were 71% in patients with a CR, 32% in patients

with a PR, and 16% in patients with SD.¹⁵ In comparison, the 5-year OS rates in CheckMate 067 were 90%, 65%, and 24% with nivolumab plus ipilimumab combination therapy and 93%, 63%, and 40% with nivolumab monotherapy.⁵³ Data obtained with pembrolizumab in a pooled analysis of KEYNOTE-001 and KEYNOTE-006 showed a similar trend, with 4-year OS rates of 98%, 75%, and 66% in patients with a CR, PR, or SD at week 24, respectively.⁶⁴

After taking into consideration both the data above and our clinical experience, we propose that patients diagnosed with advanced melanoma may be considered functionally cured if they have responded to treatment and have been off treatment for at least 2 years without disease progression. In our opinion, a patient still receiving treatment cannot be considered functionally cured. Furthermore, while a patient is still on treatment, it cannot be determined whether ongoing treatment is necessary to control their disease.

Evaluating the value of long-term survival

As these data for melanoma treatments continue to mature, landmark survival rates at 5 and 10 years will become increasingly relevant. It is also becoming important to understand what long-term survival means to patients, physicians, payers, and other healthcare professionals. In this regard, new analytical approaches may help to describe long-term survival and inform decision-making. Treatment-free survival (TFS) and treatment-free interval (TFI) are metrics that may be considered alongside traditional efficacy measures. TFS characterizes the proportion of time without both treatment or toxicity on a particular regimen; it can be further separated into time with, and time without treatment-related toxicity of a specified grade.⁶⁵ TFI captures the time interval from the last dose of study drug to subsequent systemic therapy.¹² For example, data suggest that TFI positively impacts QoL and may improve the cost-effectiveness of immune checkpoint inhibitors.^{66 67} TFI data may help to determine the optimal duration of treatment. In CheckMate 067, the median TFI for patients receiving nivolumab plus ipilimumab combination therapy was 18.1 months compared with 1.8 months with nivolumab monotherapy.¹² These data indicate that a short duration of highly active combination therapy may be sufficient for many patients to achieve durable tumor control without requiring ongoing treatment. Additionally, survival at 5 years was similar in patients who discontinued combination therapy due to AEs during the induction phase and in the overall population.¹² Emerging data from studies in the neoadjuvant setting also suggest that a short course of active therapy may be sufficient to achieve durable disease control.^{68 69}

The quality of prolonged survival is a potentially important relative determinant of value. The 5-year results from CheckMate 067 showed that QoL was maintained during the TFI after discontinuation of treatment with nivolumab plus ipilimumab combination therapy or nivolumab monotherapy.¹² Another analysis (Q-TWiST) provided evidence that longer quality-adjusted time

without symptoms or toxicity is attainable for patients treated with nivolumab plus ipilimumab than for those treated with ipilimumab alone.⁷⁰ QoL data for patients with metastatic melanoma are available from key studies with pembrolizumab⁷¹ and dabrafenib plus trametinib,^{72 73} but the analysis period to date is limited to 2 years after treatment initiation.

For cancer therapies yielding long-term survival and hence functional cures in a proportion of patients, traditional statistical analyses may not accurately estimate OS due to heterogeneity in patient outcomes. Mixture-cure models can be used to capture survival heterogeneity among patients by assuming that a certain proportion of patients are ‘cured’; the mortality risk among these patients is considered to be the same as that of the general population. Mixture-cure models have been used in various cancer settings, including melanoma in the adjuvant setting, to complement standard survival analyses. By assuming that a proportion of patients in a study will never experience an event of interest, these models allow a more accurate quantification of the proportion of patients who gain a long-term survival benefit (ie, are cured).^{74 75} Determining the ‘cure fraction’ with agents in the metastatic setting may provide another valuable comparative metric.^{76 77}

Opportunities to improve survival in more patients with metastatic melanoma

While some patients achieve long-term survival, improving outcomes for those who do not remains an important goal (figure 3).⁷⁸ One approach is to identify biomarkers or characteristics that identify the patients most likely to achieve long-term survival. Beyond established prognostic factors such as baseline LDH and disease burden/sites of disease, analyses of data from key phase III studies have failed to identify reliable predictors of poor survival.^{12 15 79} Analysis of long-term data from a phase I study of nivolumab in patients with advanced solid tumors, including melanoma, suggested that higher absolute lymphocyte count or a higher number of circulating regulatory T cells at baseline was related to survival at 5 years, warranting further evaluation.¹⁷ Identifying predictive biomarkers remains a focus of research; potential biomarkers under evaluation include tumor mutational burden and neoantigen expression, T-cell receptor sequencing, immune-related gene expression, and tumor T-cell infiltration.⁷⁹ Other opportunities to improve long-term survival include identifying effective adjuvant treatments for patients with early-stage disease, improving the efficacy of adjuvant treatment in high-risk patients (perhaps through treatment in the neoadjuvant setting or with novel combinations), and the development of treatment regimens designed to overcome the innate and acquired mechanisms of tumor resistance.

When considering new treatment combinations as a way to improve survival, one approach under evaluation is the use of I-O plus targeted therapy in patients with *BRAF*-mutant melanoma, based on reports indicating

Figure 3 Improving long-term survival in patients with metastatic melanoma. LDH, lactate dehydrogenase. (Figure from Ascierto and Dummer. Immunological effects of BRAF+MEK inhibition.⁷⁸ Adapted with permission from Taylor & Francis Ltd, <http://www.tandfonline.com>).

that BRAF and MEK inhibitors may impact tumor antigen expression, T-cell infiltration, and immune-cell populations, making the tumor microenvironment more amenable to I-O therapy.⁸⁰ Several studies are in progress, and phase III results are becoming available. The phase III IMspire 170 trial (NCT03273153) was designed to evaluate atezolizumab (PD-L1 inhibitor) plus cobimetinib (MEK inhibitor) versus pembrolizumab in untreated patients with *BRAF* wild-type tumors: unfortunately, it failed to meet its primary endpoint and PFS was not improved.⁸¹ Recently, positive results were presented from a phase III triplet trial, IMspire150 (NCT02908672), in which cobimetinib plus vemurafenib was evaluated with or without atezolizumab in untreated patients with *BRAF*-mutant melanoma. The triplet combination significantly improved PFS compared with cobimetinib plus vemurafenib (15.1 vs 10.6 months), although objective response rates (ORRs) were similar; median OS data, although not mature, suggested a benefit with the triplet regimen. The safety profile of the triplet combination was tolerable and manageable.⁸² In the phase II KEYNOTE-022 triplet study of dabrafenib plus trametinib with or without pembrolizumab, the difference in median PFS between the arms (16.9 vs 10.7 months) did

not reach statistical significance. At 24 months, the PFS rate was numerically higher in the triplet compared with the doublet arm (41% vs 16%), the overall response rate was lower (ORR; 63% vs 72%), the duration of response was longer (55% vs 16% at 24 months), and there was a higher incidence of grade ≥ 3 treatment-related AEs (58% vs 25%), respectively. Median OS was not reached with the triplet compared with 26.3 months with doublet and the 2-year OS rate was higher with the triplet (63% vs 52%). These data suggest a potential benefit with the triplet combination that will need to be confirmed with longer follow-up.^{83 84} In COMBI-i (NCT02967692), a phase III study of dabrafenib plus trametinib with or without spartalizumab (PD-1 inhibitor), results from the non-randomized part of the trial included an ORR of 78% and grade ≥ 3 treatment-related AEs in 72% of patients treated with the triplet therapy.⁸⁵ While initial ORRs from some trials are high, no information has been reported regarding the durability of these responses and whether they will translate into an OS benefit relative to BRAF plus MEK inhibition or I-O agents alone, or with these approaches used in sequence. The generally high incidences of grade ≥ 3 AEs with triplet regimens is also a concern. In the future, a more appropriate control arm

for such trials may be nivolumab plus ipilimumab, which represents an alternative and possibly more powerful standard of care than the comparators used so far.

IMPemBra (NCT02625337) was a phase II study designed to evaluate several dabrafenib plus trametinib dosing schedules in combination with pembrolizumab.⁸⁶ Early results from this small trial suggest that intermittent, short-term administration of dabrafenib plus trametinib in combination with pembrolizumab may provide the efficacy benefits expected with triplet therapy while minimizing both toxicity and the development of resistance.⁸⁷ Results have recently been presented from the single-arm, phase II TRIDeNT study (NCT02910700), which includes treatment with a combination of nivolumab, dabrafenib, and trametinib in patients with *BRAF*-mutant melanoma.⁸⁸ Unlike other comparable trials, patients with disease that was refractory to previous PD-1 inhibitor therapy or those with brain metastases are eligible for inclusion. The incidence of grade 3 or 4 treatment-related AEs (65% of patients) was similar to those of other triplet trials. Efficacy data showed an ORR of 100% in PD-1 inhibitor-naïve patients (including two CRs) and an ORR of 83% in patients with refractory disease (including one CR); ORRs were similar in patients with and without brain metastases (88%, including two CRs, in patients with brain metastases; 93%, including one CR, in those without). These data suggest that triplet therapy may have particular value, and merits further evaluation in patients with PD-1 inhibitor-refractory disease and/or those with brain metastases.

With the efficacy of triplet regimens still to be determined, a rational approach may be to use what is thought to be the most effective treatment first. Ongoing trials such as SECOMBIT, DREAMseq, and EBIN should ultimately help to determine which sequence of I-O and targeted agents is most active for particular subsets of patients and whether using a PD-1 inhibitor with intermittent or short-term targeted therapy is effective.

Other agents designed to target different components of the immune system or oncogenic processes/pathways are under evaluation, either alone or in combination with a PD-1 inhibitor. In the phase III LEAP-3 study, pembrolizumab is being evaluated in combination with lenvatinib, a multireceptor tyrosine kinase inhibitor, in previously untreated patients with advanced melanoma (NCT03820986).⁸⁹ Another approach is to combine an I-O agent with a vaccine. A phase III trial (KEYNOTE-034/MASTERKEY265/NCT02263508)⁹⁰ designed to evaluate TVEC plus pembrolizumab was started after results from a phase Ib study showed that the vaccine followed by combination therapy gave a response rate of 62% in patients with advanced melanoma, with no dose-limiting toxicities reported.⁹¹

Agents are under evaluation that target other immune checkpoint receptors, such as lymphocyte-activation gene-3 (LAG-3; eg, relatlimab)⁹² and glucocorticoid-induced tumor necrosis factor receptor (GITR).⁹³ A phase II/III study of relatlimab plus nivolumab versus

nivolumab in patients with untreated metastatic unresectable melanoma was initiated (NCT03470922)⁹⁴ after phase I/IIa results showed responses with the combination in a heavily pretreated population of patients with melanoma and progression on prior anti-PD-1/anti-PD-L1 therapy. Data from this initial trial also showed that the combination had a safety profile similar to that of nivolumab monotherapy.⁹² Bempegaldesleukin is a pegylated IL-2 pathway agonist. In the phase I/II PIVOT-02 study, an ORR of 53% (34% CRs) was reported with first-line bempegaldesleukin plus nivolumab treatment of patients with metastatic melanoma; responses were durable and were observed regardless of PD-L1 expression and the combination was well tolerated.⁹⁵ Bempegaldesleukin plus nivolumab is now under evaluation in a randomized phase III study in patients with previously untreated, unresectable, or metastatic melanoma (NCT03635983).⁹⁶ Another approach involves using toll-like receptor-9 agonists to stimulate innate and adaptive antitumor immune responses. These agents, including tiltsotolimod and CMP-001, are being studied in melanoma.^{97,98} A randomized, phase II trial with intradermal tiltsotolimod in pT3-4 cN0M0 melanoma (INTRIM) is in progress (NCT04126876).⁹⁹ Data from phase I studies showed acceptable tolerability and evidence of antitumor activity alone (intratumoral tiltsotolimod) and in combination with pembrolizumab (CMP-001).^{97,98} Other agents are designed to target the tumor, although often with the goal of mediating immunomodulatory effects. These include entinostat, a histone deacetylase inhibitor that leads to the downregulation of immunosuppressive cell types in the tumor microenvironment.¹⁰⁰ BMS-986179 is an anti-CD73 antibody that prevents the conversion of AMP to adenosine, with the goal of stopping adenosine-associated immunosuppression and tumor progression.¹⁰¹ Another approach involves inhibiting heat shock protein 90 (HSP90), for example, with XL888; preclinical data suggest that inhibiting HSP90 may help to overcome resistance to BRAF plus MEK inhibitor combinations.¹⁰²

CONCLUSION

The potential for long-term survival in patients with metastatic melanoma is an important decision-making factor and, accordingly, associated outcomes need to be assessed appropriately and accurately. Metrics that encompass economic or QoL parameters may be particularly helpful in describing the long-term value of agents used to treat these patients. The current data indicate that several treatment approaches offer the potential for long-term survival. Combined CTLA-4 and PD-1 inhibitor immunotherapy results in the highest OS plateau (up to 52% of patients), and in our opinion, may achieve long-term survival (functional cure) in the greatest number of patients. We propose that patients may be considered functionally cured if they have responded to treatment and have been off treatment for at least 2 years without disease progression. There remains a need to improve

outcomes for those patients who do not experience long-term OS.

Acknowledgements Professional medical writing and editorial assistance were provided by Rebecca Turner and Michele Salernitano at Ashfield Healthcare Communications, funded by Bristol Myers Squibb.

Contributors OM, MBA, HBK, RD, and PAA all contributed to the conception, content, writing and revisions, and have read and approved the submitted manuscript.

Funding This work was supported by Bristol Myers Squibb.

Competing interests OM: Consulting or advisory role: Amgen, Bristol Myers Squibb, Merck Sharp & Dohme, Novartis, and Roche. Travel, accommodations, and expenses: Bristol Myers Squibb and Merck Sharp & Dohme. Expert testimony: Bristol Myers Squibb. Research funding: Bristol Myers Squibb, Merck Sharp & Dohme, and NeraCare GmbH. MBA: Advisory board: Arrowhead, Bristol Myers Squibb, Exelixis, Fathom, Leads, Merck, Novartis, PACT, Pfizer, Pneuma, Pyxis Oncology, and Werewolf. Consultant: Array, Aveo, Boehringer Ingelheim, Bristol Myers Squibb, Cota, Eisai, Galactone (personal fees), Genentech/Roche, ImmunoCore, Iovance, Merck, Newlink, Novartis, Pfizer, and Surface. Research support to institution: Bristol Myers Squibb, Merck, and Pfizer. Clinical trial involvement: Aveo (Tivo 3), Bristol Myers Squibb (CM-214, CM-004, CM-067, CM-204, CM-218, CM-238, CM-905, X4P-X4 RCCA, X4-RCCB), Genentech/Roche (ImMotion 150, ImMotion 151), Merck (KN-426, KN-427, KN-029, KN-564), and Pfizer (029 Trial, Javelin 101). Speaker's bureau: None. Stock options: Pyxis Oncology and Werewolf. HBK: Employment: Bristol Myers Squibb. Stock and other ownership interests: Bristol Myers Squibb. RD: Consulting or advisory role: Amgen, Bristol Myers Squibb, Merck Sharp & Dohme, Novartis, Pierre Fabre, Roche, Sanofi, Sun Pharma, and Takeda. Honoraria: Amgen, Bristol Myers Squibb, Merck Sharp & Dohme, Novartis, Pierre Fabre, Roche, Sanofi, Sun Pharma, and Takeda. Research funding: Amgen, Bristol Myers Squibb, Merck Sharp & Dohme, Novartis, and Roche. PAA: Consulting or advisory role: ASC, Alkermes, Amgen, Array BioPharma, AstraZeneca, Bristol Myers Squibb, Genmab, Idera, Immunocore, Incyte, Italfarmaco, MedImmune, Merck Serono, Merck Sharp & Dohme, Nektar, NewLink Genetics, Novartis, Pierre Fabre, Roche/Genentech, Sandoz, Sanofi, Sun Pharma, Syndax, and Ultimovacs. Travel, accommodations, and expenses: Merck Sharp & Dohme. Stock and other ownership interests: PrimeVax. Research funding: Array BioPharma, Bristol Myers Squibb, and Roche/Genentech.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer-reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Michael B Atkins <http://orcid.org/0000-0003-3901-9924>

Paolo Antonio Ascierto <http://orcid.org/0000-0002-8322-475X>

REFERENCES

- Schvartsman G, Taranto P, Glitz IC, et al. Management of metastatic cutaneous melanoma: updates in clinical practice. *Ther Adv Med Oncol* 2019;11:1758835919851663.
- Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Combined nivolumab and ipilimumab or monotherapy in untreated melanoma. *N Engl J Med* 2015;373:23–34.
- Long GV, Stroyakovskiy D, Gogas H, et al. Combined BRAF and MEK inhibition versus BRAF inhibition alone in melanoma. *N Engl J Med* 2014;371:1877–88.
- Robert C, Karaszewska B, Schachter J, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med* 2015;372:30–9.
- Robert C, Ribas A, Schachter J, et al. Pembrolizumab versus ipilimumab in advanced melanoma (KEYNOTE-006): post-hoc 5-year results from an open-label, multicentre, randomised, controlled, phase 3 study. *Lancet Oncol* 2019;20:1239–51.
- Ascierto PA, McArthur GA, Dréno B, et al. Cobimetinib combined with vemurafenib in advanced BRAF(V600)-mutant melanoma (coBRIM): updated efficacy results from a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2016;17:1248–60.
- Dummer R, Ascierto PA, Gogas HJ, et al. Overall survival in patients with BRAF-mutant melanoma receiving encorafenib plus binimetinib versus vemurafenib or encorafenib (COLUMBUS): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol* 2018;19:1315–27.
- Bocquet-Tremoureux S, Scharbarg E, Nguyen J-M, et al. Efficacy and safety of nivolumab in metastatic melanoma: real-world practice. *Eur J Dermatol* 2019;29:315–21.
- Cowey CL, Liu FX, Boyd M, et al. Real-world treatment patterns and clinical outcomes among patients with advanced melanoma: a retrospective, community oncology-based cohort study (a STROBE-compliant article). *Medicine* 2019;98:e16328.
- Liu FX, Ou W, Dieder SJ, et al. Real-world experience with pembrolizumab in patients with advanced melanoma: a large retrospective observational study. *Medicine* 2019;98:e16542.
- Hamid O, Robert C, Daud A, et al. 5-year survival outcomes in patients with advanced melanoma treated with pembrolizumab in KEYNOTE-001. *Presented at American Society of Clinical Oncology (ASCO) Annual Meeting*, 2018.
- Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Five-year survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 2019;381:1535–46.
- Ascierto PA, Dummer R, Gogas HJ, et al. Update on tolerability and overall survival in COLUMBUS: landmark analysis of a randomised phase 3 trial of encorafenib plus binimetinib vs vemurafenib or encorafenib in patients with BRAF V600-mutant melanoma. *Eur J Cancer* 2020;126:33–44.
- McArthur GA, Dréno B, Larkin J, et al. 5-year survival update of cobimetinib plus vemurafenib in BRAFV600 mutation-positive advanced melanoma: final analysis of the coBRIM study. *Presented at Society for Melanoma Research (SMR)*, Salt Lake City, UT, 2019.
- Robert C, Grob JJ, Stroyakovskiy D, et al. Five-year outcomes with dabrafenib plus trametinib in metastatic melanoma. *N Engl J Med* 2019;381:626–36.
- Robert C, Long GV, Brady B, et al. Five-year survival outcomes in patients with BRAF wild-type advanced melanoma who received nivolumab (NIVO) monotherapy in the phase 3 CheckMate 066 study. *Presented at Society for Melanoma Research (SMR)*, Salt Lake City, UT, 2019.
- Topalian SL, Hodi FS, Brahmer JR, et al. Five-year survival and correlates among patients with advanced melanoma, renal cell carcinoma, or Non-Small cell lung cancer treated with nivolumab. *JAMA Oncol* 2019;5:1411–20.
- Dunn GP, Old LJ, Schreiber RD. The three ES of cancer immunoeediting. *Annu Rev Immunol* 2004;22:329–60.
- Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 2012;12:252–64.
- Topalian SL, Weiner GJ, Pardoll DM. Cancer immunotherapy comes of age. *J Clin Oncol* 2011;29:4828–36.
- Wei SC, Duffy CR, Allison JP. Fundamental mechanisms of immune checkpoint blockade therapy. *Cancer Discov* 2018;8:1069–86.
- Wang C, Thudium KB, Han M, et al. In vitro characterization of the anti-PD-1 antibody nivolumab, BMS-936558, and in vivo toxicology in non-human primates. *Cancer Immunol Res* 2014;2:846–56.
- Brahmer JR, Drake CG, Wolner I, et al. Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates. *J Clin Oncol* 2010;28:3167–75.
- Dafni U, Michielin O, Lluesma SM, et al. Efficacy of adoptive therapy with tumor-infiltrating lymphocytes and recombinant interleukin-2 in advanced cutaneous melanoma: a systematic review and meta-analysis. *Ann Oncol* 2019;30:1902–13.
- Rosenberg SA, Yang JC, Sherry RM, et al. Durable complete responses in heavily pretreated patients with metastatic melanoma using T-cell transfer immunotherapy. *Clin Cancer Res* 2011;17:4550–7.
- Topalian SL, Muul LM, Solomon D, et al. Expansion of human tumor infiltrating lymphocytes for use in immunotherapy trials. *J Immunol Methods* 1987;102:127–41.
- Slingluff CL, Lee S, Zhao F, et al. A randomized phase II trial of multi-peptide vaccination with melanoma peptides for cytotoxic T cells and helper T cells for patients with metastatic melanoma (E1602). *Clin Cancer Res* 2013;19:4228–38.
- Testori A, Richards J, Whitman E, et al. Phase III comparison of vitespen, an autologous tumor-derived heat shock protein gp96 peptide complex vaccine, with physician's choice of treatment for stage IV melanoma: the C-100-21 Study Group. *J Clin Oncol* 2008;26:955–62.

- 29 Faries MB, Mozzillo N, Kashani-Sabet M, *et al.* Long-term survival after complete surgical resection and adjuvant immunotherapy for distant melanoma metastases. *Ann Surg Oncol* 2017;24:3991–4000.
- 30 Eggermont AMM, Suciu S, Rutkowski P, *et al.* Adjuvant ganglioside GM2-KLH/QS-21 vaccination versus observation after resection of primary tumor > 1.5 mm in patients with stage II melanoma: results of the EORTC 18961 randomized phase III trial. *J Clin Oncol* 2013;31:3831–7.
- 31 Drugs@FDA: FDA-Approved Drugs. Available: <https://www.accessdata.fda.gov/scripts/cder/daf/> [Accessed 17 Dec 2019].
- 32 U.S. Food & Drug Administration (FDA). IMLYGIC (talimogene laherparepvec). Available: <https://www.fda.gov/vaccines-blood-biologics/cellular-gene-therapy-products/imlygic-talimogene-laherparepvec> [Accessed 26 Feb 2020].
- 33 Andtbacka RHI, Kaufman HL, Collichio F, *et al.* Talimogene laherparepvec improves durable response rate in patients with advanced melanoma. *J Clin Oncol* 2015;33:2780–8.
- 34 Andtbacka RHI, Collichio F, Harrington KJ, *et al.* Final analyses of OPTIM: a randomized phase III trial of talimogene laherparepvec versus granulocyte-macrophage colony-stimulating factor in unresectable stage III–IV melanoma. *J Immunother Cancer* 2019;7:145.
- 35 Kwak M, Leick KM, Melssen MM, *et al.* Vaccine strategy in melanoma. *Surg Oncol Clin N Am* 2019;28:337–51.
- 36 Russo A, Ficili B, Candido S, *et al.* Emerging targeted therapies for melanoma treatment (review). *Int J Oncol* 2014;45:516–24.
- 37 Long GV, Hauschild A, Santinami M, *et al.* Adjuvant dabrafenib plus trametinib in stage III BRAF-mutated melanoma. *N Engl J Med* 2017;377:1813–23.
- 38 Weber J, Mandala M, Del Vecchio M, *et al.* Adjuvant nivolumab versus ipilimumab in resected stage III or IV melanoma. *N Engl J Med Overseas Ed* 2017;377:1824–35.
- 39 Eggermont AMM, Chiarion-Sileni V, Grob J-J, *et al.* Prolonged survival in stage III melanoma with ipilimumab adjuvant therapy. *N Engl J Med* 2016;375:1845–55.
- 40 Eggermont AMM, Blank CU, Mandala M, *et al.* Adjuvant pembrolizumab versus placebo in resected stage III melanoma. *N Engl J Med* 2018;378:1789–801.
- 41 National Comprehensive Cancer Network®. NCCN clinical practice guidelines in oncology (NCCN Guidelines®): cutaneous melanoma version 1.2020 — December 19, 2019. Available: https://www.nccn.org/professionals/physician_gls/pdf/cutaneous_melanoma.pdf [Accessed 27 Feb 2020].
- 42 Postow MA, Chesney J, Pavlick AC, *et al.* Nivolumab and ipilimumab versus ipilimumab in untreated melanoma. *N Engl J Med* 2015;372:2006–17.
- 43 Atkins MB, Lotze MT, Dutcher JP, *et al.* High-Dose recombinant interleukin 2 therapy for patients with metastatic melanoma: analysis of 270 patients treated between 1985 and 1993. *J Clin Oncol* 1999;17:2105–16.
- 44 Ascierto PA, Del Vecchio M, Robert C, *et al.* Ipilimumab 10 mg/kg versus ipilimumab 3 mg/kg in patients with unresectable or metastatic melanoma: a randomised, double-blind, multicentre, phase 3 trial. *Lancet Oncol* 2017;18:611–22.
- 45 Atkins MB, Kirkwood JM, Wolchok JD, *et al.* Long-Term follow-up of CA209-004, a phase I dose-escalation study of combined nivolumab (NIVO) and ipilimumab (IPI) in patients with advanced melanoma. Presented at American Society of Clinical Oncology (ASCO) Annual Meeting, Chicago, IL, 2019.
- 46 Schadendorf D, Hodi FS, Robert C, *et al.* Pooled analysis of long-term survival data from phase II and phase III trials of ipilimumab in unresectable or metastatic melanoma. *J Clin Oncol* 2015;33:1889–94.
- 47 Atkins MB, Tarhini A, Rael M, *et al.* Comparative efficacy of combination immunotherapy and targeted therapy in the treatment of BRAF-mutant advanced melanoma: a matching-adjusted indirect comparison. *Immunotherapy* 2019;11:617–29.
- 48 Yushak M, Chapman P, Robert C, *et al.* Systemic therapy options for patients with unresectable melanoma. *Am Soc Clin Oncol Educ Book* 2017;37:661–72.
- 49 ClinicalTrials.gov. A Three Arms Prospective, Randomized Phase II Study to Evaluate the Best Sequential Approach With Combo Immunotherapy (Ipilimumab/Nivolumab) and Combo Targeted Therapy (LGX818/MEK162) in Patients With Metastatic Melanoma and BRAF Mutation. Available: <https://clinicaltrials.gov/ct2/show/NCT02631447> [Accessed 21 Oct 2019].
- 50 ClinicalTrials.gov. DREAMseq (Doublet, Randomized Evaluation in Advanced Melanoma Sequencing) a Phase III Trial. Available: <https://clinicaltrials.gov/ct2/show/NCT02224781> [Accessed 21 Oct 2019].
- 51 ClinicalTrials.gov. Combination of Targeted Therapy (Encorafenib and Binimetinib) Followed by Combination of Immunotherapy (Ipilimumab and Nivolumab) vs Immediate Combination of Immunotherapy in Patients With Unresectable or Metastatic Melanoma With BRAF V600 Mutation an EORTC Randomized Phase II Study (EBIN). Available: <https://clinicaltrials.gov/ct2/show/NCT03235245> [Accessed 21 Oct 2019].
- 52 ClinicalTrials.gov. Evaluating the Efficacy and Safety of a Sequencing Schedule of Cobimetinib Plus Vemurafenib Followed by Immunotherapy With an Anti- PD-L1 Antibody in Patients With Unresectable or Metastatic BRAF V600 Mutant Melanoma (ImmunoCobiVem). Available: <https://clinicaltrials.gov/ct2/show/NCT02902029> [Accessed 21 Oct 2019].
- 53 Long GV, Hodi FS, Chiarion-Sileni V, *et al.* Characteristics of long-term survivors and subgroup analyses with combination nivolumab plus ipilimumab for advanced melanoma (CheckMate 067). Presented at Society for Melanoma Research (SMR), Salt Lake City, UT, 2019.
- 54 Ascierto PA, Long GV. Progression-Free survival landmark analysis: a critical endpoint in melanoma clinical trials. *Lancet Oncol* 2016;17:1037–9.
- 55 Imai H, Kaira K, Minato K. Clinical significance of post-progression survival in lung cancer. *Thorac Cancer* 2017;8:379–86.
- 56 Hodi FS, Chiarion-Sileni V, Gonzalez R, *et al.* Nivolumab plus ipilimumab or nivolumab alone versus ipilimumab alone in advanced melanoma (CheckMate 067): 4-year outcomes of a multicentre, randomised, phase 3 trial. *Lancet Oncol* 2018;19:1480–92.
- 57 Wolchok JD, Chiarion-Sileni V, Gonzalez R, *et al.* Overall survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 2017;377:1345–56.
- 58 Schipper H, Goh CR, Wang TL. Shifting the cancer paradigm: must we kill to cure? *J Clin Oncol* 1995;13:801–7.
- 59 Larkin JMG, Chiarion-Sileni V, Gonzalez R, *et al.* 5-Year survival outcomes of the CheckMate 067 phase 3 trial of nivolumab plus ipilimumab (NIVO+IPI) combination therapy in advanced melanoma. Presented at the European Society for Medical Oncology (ESMO) Congress, Barcelona, Spain, 2019.
- 60 Carlino MS, Vanella V, Girgis C, *et al.* Cessation of targeted therapy after a complete response in BRAF-mutant advanced melanoma: a case series. *Br J Cancer* 2016;115:1280–4.
- 61 Betof Warner A, Palmer JS, Shoushtari AN, *et al.* Long-Term outcomes and responses to retreatment in patients with melanoma treated with PD-1 blockade. *J Clin Oncol* 2020;38:1655–63.
- 62 Tan AC, Emmett L, Lo S, *et al.* FDG-PET response and outcome from anti-PD-1 therapy in metastatic melanoma. *Ann Oncol* 2018;29:2115–20.
- 63 Osgood C, Mulkey F, Mishra-Kalyani PS, *et al.* FDA analysis of depth of response (DPR) and survival across 10 randomized controlled trials in patients with previously untreated unresectable or metastatic melanoma (UMM) by therapy type. Presented at American Society of Clinical Oncology (ASCO) Annual Meeting, Chicago, IL, 2019.
- 64 Hamid O, Robert C, Daud A, *et al.* Long-term outcomes in patients who achieved clinical benefit (stable disease, partial response, complete response) with pembrolizumab: pooled analysis of KEYNOTE-001 and KEYNOTE-006. Presented at 16th International Congress of the Society for Melanoma Research (SMR), 2019.
- 65 Regan MM, Werner L, Rao S, *et al.* Treatment-free survival: a novel outcome measure of the effects of immune checkpoint inhibition-A pooled analysis of patients with advanced melanoma. *J Clin Oncol* 2019;37:3350–8.
- 66 Acaster S, Gaugris S, Velikova G, *et al.* Impact of the treatment-free interval on health-related quality of life in patients with multiple myeloma: a UK cross-sectional survey. *Support Care Cancer* 2013;21:599–607.
- 67 Tarhini A, McDermott D, Ambavane A, *et al.* Clinical and economic outcomes associated with treatment sequences in patients with BRAF-mutant advanced melanoma. *Immunotherapy* 2019;11:283–95.
- 68 Amaria RN, Reddy SM, Tawbi HA, *et al.* Neoadjuvant immune checkpoint blockade in high-risk resectable melanoma. *Nat Med* 2018;24:1649–54.
- 69 Rozeman EA, Menzies AM, van Akkooi ACJ, *et al.* Identification of the optimal combination dosing schedule of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma (OpACIN-neo): a multicentre, phase 2, randomised, controlled trial. *Lancet Oncol* 2019;20:948–60.
- 70 McDermott DF, Shah R, Gupte-Singh K, *et al.* Quality-adjusted survival of nivolumab plus ipilimumab or nivolumab alone versus ipilimumab alone among treatment-naïve patients with advanced

- melanoma: a quality-adjusted time without symptoms or toxicity (Q-TWIST) analysis. *Qual Life Res* 2019;28:109–19.
- 71 Petrella TM, Robert C, Richtig E, et al. Patient-reported outcomes in KEYNOTE-006, a randomised study of pembrolizumab versus ipilimumab in patients with advanced melanoma. *Eur J Cancer* 2017;86:115–24.
 - 72 Grob JJ, Amonkar MM, Karaszewska B, et al. Comparison of dabrafenib and trametinib combination therapy with vemurafenib monotherapy on health-related quality of life in patients with unresectable or metastatic cutaneous BRAF Val600-mutation-positive melanoma (COMBI-v): results of a phase 3, open-label, randomised trial. *Lancet Oncol* 2015;16:1389–98.
 - 73 Schadendorf D, Amonkar MM, Stroyakovskiy D, et al. Health-related quality of life impact in a randomised phase III study of the combination of dabrafenib and trametinib versus dabrafenib monotherapy in patients with BRAF V600 metastatic melanoma. *Eur J Cancer* 2015;51:833–40.
 - 74 Hauschild A, Dummer R, Schadendorf D, et al. Longer follow-up confirms relapse-free survival benefit with adjuvant dabrafenib plus trametinib in patients with resected BRAF V600-mutant stage III melanoma. *J Clin Oncol* 2018;36:3441–9.
 - 75 Ouwens MJNM, Mukhopadhyay P, Zhang Y, et al. Estimating lifetime benefits associated with immuno-oncology therapies: challenges and approaches for overall survival extrapolations. *Pharmacoeconomics* 2019;37:1129–38.
 - 76 Weber JS, Kurt M, Edmondson-Jones M, et al. Mixture-cure modeling for resected stage III/IV melanoma in the phase III CheckMate 238 trial. *Presented at the European Society for Medical Oncology (ESMO) Congress*, 2019.
 - 77 Othus M, Bansal A, Koepf L, et al. Accounting for cured patients in cost-effectiveness analysis. *Value Health* 2017;20:705–9.
 - 78 Ascierto PA, Dummer R. Immunological effects of BRAF+MEK inhibition. *Oncoimmunology* 2018;7:e1468955.
 - 79 Tarhini A, Kudchadkar RR. Predictive and on-treatment monitoring biomarkers in advanced melanoma: moving toward personalized medicine. *Cancer Treat Rev* 2018;71:8–18.
 - 80 Pelster MS, Amaria RN. Combined targeted therapy and immunotherapy in melanoma: a review of the impact on the tumor microenvironment and outcomes of early clinical trials. *Ther Adv Med Oncol* 2019;11:1758835919830826.
 - 81 Arance AM, Gogas H, Dreno B, et al. Combination treatment with cobimetinib (C) and atezolizumab (A) vs pembrolizumab (P) in previously untreated patients (pts) with BRAFV600 wild type (wt) advanced melanoma: primary analysis from the phase III IMspire170 trial. *Ann Oncol* 2019;30:v906–34.
 - 82 McArthur GA, Stroyakovskiy D, Gogas H, et al. Evaluation of atezolizumab, cobimetinib, and vemurafenib in previously untreated patients with BRAF V600 mutation-positive advanced melanoma: primary results from the phase 3 IMspire150 trial. *Presented at American Association for Cancer Research (AACR) Virtual Annual Meeting*, 2020.
 - 83 Ascierto PA, Ferrucci PF, Fisher R, et al. Dabrafenib, trametinib and pembrolizumab or placebo in BRAF-mutant melanoma. *Nat Med* 2019;25:941–6.
 - 84 Ferrucci PF, Ascierto PA, Maio M, et al. Updated survival in patients with BRAF-mutant melanoma administered pembrolizumab, dabrafenib, and trametinib. *Presented at Society for Melanoma Research (SMR)*, Salt Lake City, UT, 2019.
 - 85 Long GV, Lebbé C, Atkinson V, et al. The anti-PD-1 antibody spartalizumab (S) in combination with dabrafenib (D) and trametinib (T) in previously untreated patients (pts) with advanced BRAF V600-mutant melanoma: Updated efficacy and safety from parts 1 and 2 of COMBI-i. *J Clin Oncol* 2019;37:9531.
 - 86 ClinicalTrials.gov. Study comparing pembrolizumab with dual MAPK pathway inhibition plus pembrolizumab in melanoma patients (IMPemBra). Available: <https://clinicaltrials.gov/ct2/show/NCT02625337> [Accessed 24 Feb 2020].
 - 87 Rozeman EA, Versluis JM, Sikorska K, et al. IMPEMBRA: phase 2 study comparing pembrolizumab with intermittent/short-term dual MAPK pathway inhibition plus PEMbrolizumab in patients harboring the BRAFV600 mutation. *Oral presentation at Melanoma Bridge*, Naples, Italy, 2019.
 - 88 Burton EM. Safety and efficacy of triplet combination of nivolumab with dabrafenib and trametinib [TRIDENT] in patients with BRAF-mutated metastatic melanoma: a single center phase II study. *Presented at Melanoma Bridge*, Naples, Italy, 2019.
 - 89 ClinicalTrials.gov. Safety and efficacy study of pembrolizumab (MK-3475) combined with lenvatinib (MK-7902/E7080) as first-line intervention in adults with advance melanoma (MK-7902-003/E7080-G000-312/LEAP-003). Available: <https://clinicaltrials.gov/ct2/show/NCT03820986> [Accessed 30 May 2020].
 - 90 ClinicalTrials.gov. Pembrolizumab with or without Talimogene Laherparepvec or Talimogene Laherparepvec placebo in unresected melanoma (KEYNOTE-034). Available: <https://clinicaltrials.gov/ct2/show/NCT02263508> [Accessed 30 May 2020].
 - 91 Ribas A, Dummer R, Puzanov I, et al. Oncolytic virotherapy promotes intratumoral T cell infiltration and improves anti-PD-1 immunotherapy. *Cell* 2017;170:e10:1109–19.
 - 92 Ascierto PA, Bono P, Bhatia S, et al. Efficacy of BMS-986016, a monoclonal antibody that targets lymphocyte activation gene-3 (LAG-3), in combination with nivolumab in patients with melanoma who progressed during prior anti-PD-1/PD-L1 therapy in all-comer and biomarker-enriched populations. *Presented at European Society for Medical Oncology (ESMO) Congress*, Madrid, Spain, 2017.
 - 93 Papadopoulos KP, Autio KA, Golan T, et al. Phase 1 study of MK-4166, an anti-human glucocorticoid-induced tumor necrosis factor receptor (GITR) antibody, as monotherapy or with pembrolizumab (pembro) in patients (pts) with advanced solid tumors. *J Clin Oncol* 2019;37:9509.
 - 94 A randomized, double-blind phase 2/3 study of relatlimab combined with nivolumab versus nivolumab in participants with previously untreated metastatic or unresectable melanoma. Available: <https://clinicaltrials.gov/ct2/show/NCT03470922> [Accessed 24 Feb 2020].
 - 95 Diab A, Puzanov I, Maio M, et al. Clinical activity of BEMPEG plus NIVO in previously untreated patients with metastatic melanoma: updated results from the phase 1/2 PIVOT-02 study. *Presented at 34th Annual Meeting Society of the Immunotherapy of Cancer (SITC)*, National Harbor, MD, 2019.
 - 96 ClinicalTrials.gov. A study of NKTR-214 combined with nivolumab vs nivolumab alone in participants with previously untreated inoperable or metastatic melanoma. Available: <https://clinicaltrials.gov/ct2/show/NCT03635983> [Accessed 29 May 2020].
 - 97 Babiker HM, Borazanci E, Subbiah V, et al. Activation of innate and adaptive immunity using intratumoral tilosotolimod (IMO-2125) as monotherapy in patients with refractory solid tumors: a phase 1B study (ILLUMINATE-101). *Presented at American Association for Cancer Research (AACR) Conference*, Atlanta, GA, 2019.
 - 98 Milhem M, Gonzales R, Medina T, et al. Intratumoral toll-like receptor 9 (TLR9) agonist, CMP-001, in combination with pembrolizumab can reverse resistance to PD-1 inhibition in a phase Ib trial in subjects with advanced melanoma. *Presented at American Association for Cancer Research (AACR) Annual Meeting*, Chicago, IL, 2018.
 - 99 ClinicalTrials.gov. A randomized controlled phase II trial with intradermal IMO-2125 in pathological tumor stage (P) T3-4 cN0M0 melanoma (INTRIM). Available: <https://clinicaltrials.gov/ct2/show/NCT04126876> [Accessed 24 Feb 2020].
 - 100 Sullivan RJ, Moschos SJ, Johnson ML, et al. Efficacy and safety of entinostat (ENT) and pembrolizumab (PEMBRO) in patients with melanoma previously treated with anti-PD1 therapy. *Presented at American Association for Cancer Research (AACR) Annual Meeting*, Atlanta, GA, 2019.
 - 101 Siu LL, Burris B, Le DT, et al. Preliminary phase 1 profile of BMS-986179, an anti-CD73 antibody, in combination with nivolumab in patients with advanced solid tumors. *Presented at American Association for Cancer Research (AACR) Annual Meeting*, 2018.
 - 102 Eroglu Z, Chen YA, Gibney GT, et al. Combined BRAF and HSP90 inhibition in patients with unresectable BRAF V600E-mutant melanoma. *Clin Cancer Res* 2018;24:5516–24.