Prognostic significance and immune correlates of CD73 expression in renal cell carcinoma

Abhishek Tripathi ^{1,2}, Edwin Lin,³ Wanling Xie,⁴ Abdallah Flaifel,⁵ John A Steinharter,² Emily N Stern Gatof,⁶ Gabrielle Bouchard,² Justin H Fleischer,² Nieves Martinez-Chanza ², Connor Gray,⁶ Charlene Mantia,⁶ Linda Thompson,⁷ Xiao X Wei,² Marios Giannakis,⁸ Bradley A McGregor,² Toni K Choueiri ², Neeraj Agarwal ³, David F McDermott,⁶ Sabina Signoretti,⁵ Lauren C Harshman ²

To cite: Tripathi A, Lin E, Xie W, *et al.* Prognostic significance and immune correlates of CD73 expression in renal cell carcinoma. *Journal for ImmunoTherapy of Cancer* 2020;**8**:e001467. doi:10.1136/jitc-2020-001467

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jitc-2020-001467>).

AT and EL are joint first authors.

Accepted 02 October 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Lauren C Harshman; lharshman@gmail.com

ABSTRACT

Background CD73–adenosine signaling in the tumor microenvironment is immunosuppressive and may be associated with aggressive renal cell carcinoma (RCC). We investigated the prognostic significance of CD73 protein expression in RCC leveraging nephrectomy samples. We also performed a complementary analysis using The Cancer Genome Atlas (TCGA) dataset to evaluate the correlation of CD73 (ecto-5'-nucleotidase (*NT5E*), CD39 (ectonucleoside triphosphate diphosphohydrolase 1 (*ENTPD1*)) and A2 adenosine receptor (A2AR; *ADORA2A*) transcript levels with markers of angiogenesis and antitumor immune response.

Methods Patients with RCC with available archived nephrectomy samples were eligible for inclusion. Tumor CD73 protein expression was assessed by immunohistochemistry and quantified using a combined score (CS: % positive cells×intensity). Samples were categorized as CD73_{negative} (CS=0), CD73_{low} or CD73_{high} (< and ≥median CS, respectively). Multivariable Cox regression analysis compared disease-free survival (DFS) and overall survival (OS) between CD73 expression groups. In the TCGA dataset, samples were categorized as low, intermediate and high *NT5E*, *ENTPD1* and *ADORA2A* gene expression groups. Gene expression signatures for infiltrating immune cells, angiogenesis, myeloid inflammation, and effector T-cell response were compared between *NT5E*, *ENTPD1* and *ADORA2A* expression groups.

Results Among the 138 patients eligible for inclusion, 'any' CD73 expression was observed in 30% of primary tumor samples. High CD73 expression was more frequent in patients with M1 RCC (29% vs 12% M0), grade 4 tumors (27% vs 13% grade 3 vs 15% grades 1 and 2), advanced T-stage (≥T3: 22% vs T2: 19% vs T1: 12%) and tumors with sarcomatoid histology (50% vs 12%). In the M0 cohort (n=107), patients with CD73_{high} tumor expression had significantly worse 5-year DFS (42%) and 10-year OS (22%) compared with those in the CD73_{negative} group (DFS: 75%, adjusted HR: 2.7, 95% CI 1.3 to 5.9, p=0.01; OS: 64%, adjusted HR: 2.6, 95% CI 1.2 to 5.8, p=0.02) independent of tumor stage and grade. In the TCGA analysis, high *NT5E* expression was associated with significantly worse 5-year OS (p=0.008). *NT5E* and *ENTPD1* expression correlated with higher regulatory

T cell (Treg) signature, while *ADORA2A* expression was associated with increased Treg and angiogenesis signatures.

Conclusions High CD73 expression portends significantly worse survival outcomes independent of stage and grade. Our findings provide compelling support for targeting the immunosuppressive and proangiogenic CD73–adenosine pathway in RCC.

INTRODUCTION

Immune checkpoint inhibitors targeting the programmed cell death-1 (PD-1) and cytotoxic T-lymphocyte-associated antigen-4 (CTLA-4) pathways have significantly advanced the treatment of metastatic renal cell carcinoma (mRCC). Recent studies combining PD-1/L1 inhibitors with either the vascular endothelial growth factor (VEGF) receptor tyrosine kinase inhibitor axitinib or the anti-CTLA-4 antibody ipilimumab have demonstrated improved objective response rates (ORRs), progression-free survival (PFS) and overall survival (OS) compared with sunitinib.^{1–4} Given these results, most patients with clear cell renal cell carcinoma (ccRCC) will receive a PD-1/L1 inhibitor-based regimen either in combination with axitinib or, if intermediate or poor-risk disease, ipilimumab, as first-line therapy. Although immune checkpoint blockade represents a significant therapeutic advance, approximately 20% of patients are primary treatment-refractory to these agents, and the majority who experience clinical benefit eventually progress as evidenced by a median PFS of 12–14 months on any of these regimens.^{1–4} The lack of response in a significant proportion of patients and eventual disease progression in most patients suggests the existence of multiple non-redundant mechanisms of de novo or acquired

resistance to immune checkpoint inhibition in the tumor microenvironment.

CD73 (ecto-5'-nucleotidase (*NT5E*)) and CD39 (ectonucleoside triphosphate diphosphohydrolase 1 (*ENTPDI*)) mediate sequential dephosphorylation of extracellular ATP to adenosine with CD73 catalyzing the rate-limiting step.⁵ A hypoxic tumor microenvironment significantly increases CD73 expression on tumor cells and tumor-infiltrating immune cells through hypoxia-inducible factor (HIF)-1 α .^{6–8} Adenosine generated by CD73 and CD39 binds to and activates G protein-coupled A1, A2A, A2B, and A3 receptors.⁹ In preclinical models, increased adenosine signaling attenuates the antitumor immune response through the proliferation of regulatory T cells (Tregs) and myeloid-derived suppressor cells (MDSCs) and differentiation of tumor-associated macrophages into the immunosuppressive M2 phenotype.^{10–13} Adenosine signaling may also stimulate angiogenesis through increased vasodilation, release of proangiogenic factors such as VEGF, and recruitment of endothelial progenitor cells in the tumor microenvironment.^{14–16}

Previously, our group found that CD73 was more frequently expressed in mRCC specimens compared with primary tumors.¹⁷ As CD73 mediates the rate-limiting step in the generation of immunosuppressive adenosine, we hypothesized that higher CD73 expression correlates with more aggressive disease in renal cell carcinoma (RCC). Employing an institutional dataset of RCC samples, we investigated the prognostic significance of CD73 expression in localized RCC. Using gene expression data from The Cancer Genome Atlas (TCGA) Kidney Renal Clear Cell Carcinoma (KIRC) dataset, we also evaluated the correlation of mRNA expression of CD73 (*NT5E*), CD39 (*ENTPDI*) and A2AR (*ADORA2A*) with gene expression signatures reflecting angiogenesis, myeloid inflammation and effector T-cell response (T_{eff}) and infiltrating immune cell subsets.

METHODS

Institutional dataset

The study used previously established RCC tissue microarrays (TMAs) which included primary tumor samples from patients with localized or de novo mRCC (T1–4, N0–1, M0–1), who underwent nephrectomy at Dana-Farber Cancer Institute/Brigham and Women's Hospital and Beth Israel Deaconess Medical Center between January 2002 and May 2006. Immunohistochemistry (IHC) for CD73 was performed on the formalin-fixed paraffin-embedded tumor tissue from the TMAs. Rehydrated tissue sections were boiled in citrate buffer pH 6.0 (Life Technologies, Frederick, Maryland, USA) with a pressure cooker (Biocare Medical, Pacheco, California, USA) for 30 s at 125°C. After cooling down at room temperature (RT), tissue sections were successively incubated with a peroxidase block (Dual Endogenous Enzyme Block, Agilent) and a protein block (Serum Free Block, Agilent) for 10 min each. Sections were next incubated for 1 hour

at RT with the rabbit monoclonal anti-CD73 antibody (clone D7F9A, 1:25; Cell Signaling Technology) diluted in antibody diluent with background reducing components (Agilent). Tissue sections were then incubated for 1 hour at RT with a polyclonal mouse anti-rabbit antibody (Agilent, 1:750) diluted in antibody diluent with background reducing components (Agilent) followed by incubation for 30 min at RT with EnVision anti-mouse horseradish peroxidase (HRP)-conjugated antibody (Agilent). HRP visualization was performed by applying 3,3-diaminobenzidine+substrate (Agilent) for 1 min and 30 s. Between steps, tissue sections were washed for 5 min in washing buffer (0.1 mM Tris, pH 7.4+0.05% Tween 20). Nuclei were counterstained with hematoxylin.

von Hippel-Lindau deficient and proficient University of Michigan-Renal Carcinoma-2 cells were used as positive and negative controls, respectively. Percentage of cell staining positive for CD73 and the intensity of staining (1+, 2+ or 3+) were assessed manually by one observer (SS). CD73 expression for each sample was quantified using a combined score (CS), which was calculated by multiplying the intensity of staining (1+, 2+, or 3+) with the percentage of tumor cells staining positive. For patients with multiple tumor samples with evaluable CD73 expression, the mean CS of all available samples was calculated. CD73 positivity was defined as any CD73 expression on tumor cells irrespective of percentage of cells or the intensity of staining. Patients with CD73 positivity were categorized into CD73_{low} and CD73_{high} subgroups. Patients with CS less than the median CS of all CD73-positive patients were designated as CD73_{low}. Conversely, those with expression equal to or higher than the median CS were characterized as CD73_{high}. Clinical and pathological characteristics such as tumor stage, Fuhrman grade, presence or absence of sarcomatoid features, along with outcomes data, were cataloged through retrospective chart review by investigators blinded to the CD73 expression data.

TCGA dataset

RNA-seq data for 538 ccRCC samples derived from primary tumors were obtained from the TCGA-KIRC dataset.¹⁸ Gene expression of CD73 (*NT5E*), CD39 (*ENTPDI*) and A2AR (*ADORA2A*) was quantified by HT-seq and measured in upper-quartile normalized fragments per kilobase million.¹⁹ Patients were split into low (≤ -1 SD from the overall mean), intermediate (-1 to 1 SD from the mean), and high (≥ 1 SD from the mean) *NT5E*, *ENTPDI* and *ADORA2A* expression groups. Averaged log₂-transformed expression of previously validated genes was used to calculate angiogenesis (*VEGFA*, *KDR*, *ESM1*, *PECAM1*, *ANGPTL4* and *CD34*), myeloid inflammation (*IL-6*, *CXCL1*, *CXCL2*, *CXCL3*, *CXCL8* and *PTGS2*), and T_{eff} (*CD8A*, *EOMES*, *PRFI*, *IFNG* and *CD274*) gene expression signatures.²⁰ Gene signatures for tumor-infiltrating immune cells such as B cells, CD4⁺ T cells, CD8⁺ T cells, neutrophils, NK cells, macrophages, and dendritic cells were calculated based on averaged

Table 1 Baseline characteristics of patients (N=138)

	Localized RCC (N=107)		De novo mRCC (N=31)		Total (N=138)	
	n	%	n	%	n	%
Gender						
Female	43	40	14	45	57	41
Male	64	60	17	55	81	59
Histology						
ccRCC	78	73	25	81	103	75
nccRCC	28	26	5	16	33	24
Chromophobe	14	13	.	.	14	10
Papillary not subtyped	6	6	1	3	7	5
Papillary type 1	4	4	1	3	5	4
Papillary type 2	3	3	1	3	4	3
RCC unclassified	1	1	2	7	3	2
Unknown	1	1	1	3	2	1
Pathological T stage						
T1	58	54	2	6	60	44
T2	21	20	7	23	28	20
T3	21	20	21	68	42	30
T4	3	3	1	3	4	3
Unknown	4	4	.	.	4	3
Pathological N stage						
N0	33	31	11	36	44	32
N1	8	8	6	19	14	10
Nx	62	58	14	45	76	55
Unknown	4	4	.	.	4	3
AJCC stage						
I	57	53	.	.	57	41
II	19	18	.	.	19	14
III (T3 or N1)	26	24	.	.	26	19
IV (T4 or M1)	3	3	31	100	34	25
Unknown	2	2	.	.	2	1
Fuhrman grade						
G1	7	7	.	.	7	5
G2	59	55	9	29	68	49
G3	26	24	12	39	38	28
G4	13	12	9	29	22	16
Unknown	2	2	1	3	3	2
Sarcomatoid features						
No	100	94	24	77	124	90
Yes	4	4	6	19	10	7
Unknown	3	3	1	3	4	3

Continued

Table 1 Continued

	Localized RCC (N=107)		De novo mRCC (N=31)		Total (N=138)	
	n	%	n	%	n	%
Adjuvant therapy						
No	103	96	31	100	134	97
Yes	3	3	.	.	3	2
Unknown	1	1	.	.	1	1

AJCC, American Joint Committee on Cancer; ccRCC, clear cell renal cell carcinoma; mRCC, metastatic renal cell carcinoma; nccRCC, non-clear cell renal cell carcinoma; RCC, renal cell carcinoma.

log-transformed expression levels of previously validated immune metagenes.²¹

Statistical analysis

Baseline patient characteristics were summarized using descriptive statistics. The distribution of CD73 expression in the primary tumor (CD73_{negative} vs CD73_{low} or CD73_{high}) by baseline clinical and pathologic characteristics was compared using the Cochran-Armitage Trend test (for variables with two categories) and Jonckheere-Terpstra test (for variables three or more ordered categories). The primary survival analysis focused on patients with localized RCC for which disease-free survival (DFS) and OS were primary outcome measures. DFS was defined as the time from nephrectomy to disease recurrence or death from any cause. OS was defined as the time from nephrectomy to death from any cause. DFS and OS were censored at the date of the last follow-up. An exploratory analysis evaluated outcomes between CD73 expression groups in patients with de novo mRCC. For this analysis, OS was the primary outcome measure and defined as the time from cytoreductive nephrectomy to death from any cause. Distribution of DFS and OS were estimated using the Kaplan Meier method. Multivariable Cox regression analysis was used to estimate the association of CD73 expression with DFS and OS adjusted for known prognostic factors tumor grade (grade ≥ 3 vs < 3) and the American Joint Committee on Cancer (AJCC) disease stage (III/IV vs I/II). All statistical tests were two-sided and statistical significance was considered at $p < 0.05$.

In the TCGA dataset, the Mann-Whitney U test was used to compare AJCC disease stage and gene expression signature scores between low, intermediate, and high *NT5E*, *ENTPD1* and *ADORA2A* expression groups. Bonferroni correction was used to control false discovery rate to $\alpha < 0.05$. Significant differences were defined by a fold change > 4 and adjusted $p < 0.05$. OS was compared between low, intermediate, and high *NT5E* and *ADORA2A* groups using the Log rank test stratified by AJCC stage. All bioinformatic analyses and figures using TCGA data were produced using Pandas, NumPy, SciPy, and Matplotlib software packages in a Python 3.6 environment.

RESULTS

CD73 expression and outcomes in the Dana-Farber/Harvard Cancer Center institutional dataset

We interrogated TMAs consisting of tumor samples from 199 patients with localized or de novo mRCC. For our analysis, patients were excluded if pathologic diagnosis was not RCC (n=17), they had unevaluable CD73 expression (n=13), or there was inadequate linked clinical and pathological data (n=19) (online supplemental figure 1). In order to minimize potential heterogeneity induced by higher CD73 expression in metastatic samples compared with primary tumors,¹⁷ we excluded patients who only had samples from metastatic sites (n=12). Of the 138 eligible patients with primary tumor samples, 107 patients had localized RCC, and 31 patients had de novo mRCC. ccRCC was the most common histologic subtype (75%, n=103; table 1) with 24% non-clear cell disease (papillary, n=16; chromophobe, n=14). High nuclear grade (≥ 3) and sarcomatoid component was present in 43% (n=60) and 7% (n=10) respectively. The majority (n=88, 64%) had pT1-T2 tumors while 33% (n=46) were $\geq$ pT3 and 10% had nodal involvement (n=14). Locally advanced disease defined as AJCC stage III or IV was present in 44% (n=60) of patients.

In the overall cohort (n=138), any CD73 expression (CS >0) was seen in 30% (n=42, table 2) of samples with a median CS of 11.8 (range 0.1–210, figure 1). In patients with ccRCC, CD73 positivity was seen in 27% of tumors (median CS: 6.3, range 0.1–125) and 13% had high CD73 expression. In patients with non-clear cell histology, CD73 positivity was observed in 39% with 24% being CD73_{high}. Compared with patients with localized (M0) RCC, those with de novo mRCC were more likely to be CD73 positive (55% vs 23%; p=0.002) and have high CD73 expression (29% vs 12%, figure 2, p=0.002). Similarly, high CD73 expression was more frequent in grade four tumors (27%) compared with grade 3 (13%) or grade 1–2 (15%, $p_{\text{trend}}=0.035$, figure 2), pathologic stage $\geq$ T3 (22%) compared with T2 (19%) and T1 (12%; $p_{\text{trend}}=0.004$) and locally advanced disease stage (AJCC stage, IV: 29% vs III: 12% vs II: 21% vs I: 9%; $p_{\text{trend}} < 0.0001$). In tumors with sarcomatoid features (n=10), 90% (n=9) had any CD73 expression with 50% (n=5) being CD73_{high}. In contrast, in tumors without sarcomatoid features, only 24% (n=30) demonstrated any CD73 expression and only 12% (n=15) were CD73_{high} (p<0.001).

Median follow-up was 10 years (range <0.1–15.5 years) for patients with localized RCC (n=107). DFS at 5 years for patients in the CD73_{negative}, CD73_{low} and CD73_{high} groups was 75%, 50% and 42%, respectively (figure 3A). On multivariable analysis adjusting for Fuhrman grade (graded 1–2 vs grades 3–4) and AJCC stage (stage I/II vs III/IV), high CD73 expression was associated with a significantly worse DFS (table 3; adjusted HR: 2.72, 95% CI 1.27 to 5.85, p=0.01). Similarly, the CD73_{high} group experienced significantly worse 10-year OS compared with the CD73_{negative} group (22% vs 64%, adjusted HR: 2.59, 95% CI 1.15 to 5.84, p=0.021; figure 3B). The survival

outcomes were not significantly different between the CD73_{negative} and CD73_{low} groups (DFS HR: 1.22, 95% CI 0.50 to 2.98; OS HR: 0.78, 95% CI 0.27 to 2.30). Additional analysis combining the CD73 negative and low groups was performed and the significant differences in DFS and OS remained. The adjusted HR for DFS was 2.89 (95% CI: 1.37 to 6.07, p=0.005) and for OS was 2.86 (95% CI 1.30 to 6.29, p=0.009) in favor of no or low expression (CD73_{high} vs CD73_{negative+low}) (table 3). In an exploratory analysis evaluating OS in patients with de novo mRCC (n=31), there was no statistically significant difference in 2-year OS between patients in the CD73_{negative} (43%), CD73_{low} (50%) and CD73_{high} groups (52%, online supplemental figure 2; p=0.52).

Clinical and genomic correlates of *NT5E*, *ENTPD1* and *ADORA2A* gene expression in the TCGA dataset

In the complementary analysis using TCGA RNA-seq data, high *NT5E* expression was associated with significantly lower 5-year OS compared with intermediate or low *NT5E* expression (figure 4A) in AJCC stage IV tumors. No significant differences in OS were observed between *NT5E* expression groups in patients with AJCC stage I–III disease (online supplemental figure 3A–C). *NT5E* expression was similar across AJCC stages (online supplemental figure 3D). Similarly, there was no significant correlation of *ADORA2A* expression with OS or AJCC disease stage (online supplemental figure 4). Treg gene expression signature was positively correlated with *NT5E*, *ENTPD1* and *ADORA2A* expression (figure 4). The angiogenesis gene expression signature positively correlated with *ADORA2A* (figure 4D) but not *NT5E* or *ENTPD1* expression (online supplemental tables 1–3). T_{eff} myeloid inflammation, and other immune cell signatures (eg, CD8⁺ T cells, B cells, neutrophils, and macrophages) were not significantly different between the low intermediate and high *NT5E*, *ENTPD1* or *ADORA2A* expression groups.

DISCUSSION

In this biomarker-based analysis, CD73 protein expression was present in 30% of all RCC nephrectomy tumor specimens. Higher expression correlated with more aggressive disease as defined by higher Fuhrman nuclear grade, advanced stage at diagnosis, presence of sarcomatoid histology, and in patients with localized RCC, significantly worse DFS and OS. In a complementary analysis using the TCGA gene expression dataset, high *NT5E* (CD73) gene expression was associated with significantly worse OS in AJCC stage IV tumors. *NT5E* and *ENTPD1* expression correlated with increased expression of immunosuppressive Treg markers, while A2AR (*ADORA2A*) expression was associated with significantly higher angiogenesis and Treg gene expression signatures.

There is a sparse but growing literature on the prevalence and clinical significance of CD73 expression in RCC. Yu *et al* analyzed CD73 expression by IHC in patients with ccRCC, the majority of whom had localized

Table 2 Associations of CD73 expression with baseline characteristics of patients (N=138)

Baseline characteristics (N)	CD73 expression						P value*
	Positive (N=42)						
	High (n=22)		Low (n=20)		Negative (N=96)		
	n	%	n	%	n	%	
Metastatic status at presentation							0.002
Localized RCC (n=107)	13	12	12	11	82	77	
De novo mRCC (n=31)	9	29	8	26	14	45	
Histology							0.11†
ccRCC (n=103)	13	13	15	15	75	73	
nccRCC (n=33)	8	24	5	15	20	61	
Papillary (n=16)	6	38	5	31	5	31	
Chromophobe (n=14)	2	14	.	.	12	86	
RCC unclassified (n=3)	.	.	.	.	3	100	
Unknown (n=2)	1	50			1	50	
Sarcomatoid features							<0.0001
No (n=124)	15	12	15	12	94	76	
Yes (n=10)	5	50	4	40	1	10	
Unknown (n=4)	2	50	1	25	1	25	
Fuhrman nuclear grade							0.035
G1-2 (n=75)	11	15	7	9	57	76	
G3 (n=38)	5	13	8	21	25	66	
G4 (n=22)	6	27	5	23	11	50	
Unknown (n=3)					3	100	
Pathological T stage							0.004
T1 (n=60)	7	12	2	3	51	85	
T2 (n=28)	5	19	7	25	16	57	
T3-4 (n=46)	10	22	9	20	27	59	
Unknown (n=4)			2	50	2	50	
AJCC stage							<0.0001
I (n=57)	5	9	2	4	50	88	
II (n=19)	4	21	4	21	11	58	
III (n=26)	3	12	4	15	19	73	
IV (n=34)	10	29	9	27	15	44	
Unknown (n=2)			1	50	1	50	

*Cochran-Armitage Trend test (for variables with two categories) and Jonckheere-Terpstra test (for variables with ≥ 3 ordered categories); unknown group was excluded from the comparison.

†P value is for comparison of ccRCC versus nccRCC.

AJCC, American Joint Committee on Cancer; ccRCC, clear cell renal cell carcinoma; mRCC, metastatic renal cell carcinoma; nccRCC, non-clear cell renal cell carcinoma; RCC, renal cell carcinoma.

(T1–4, N0, M0) disease.²² Nearly half of tumors (48%) expressed CD73, which was associated with high tumor grade and T stage. In our prior work in RCC, we observed CD73 expression in 19% of primary tumors and that expression was significantly more common in metastatic tumor samples (67%).¹⁷ CD73 expression in the primary tumor was associated with a trend towards higher nuclear grade and numerically worse 5-year OS. Our current analysis included tumor samples from 107 patients with

localized RCC and 31 patients with de novo mRCC from three high-volume kidney cancer centers with well-annotated long-term clinical outcomes (median ~10 years from nephrectomy). Confirming the results of our prior study, tumors from patients with de novo mRCC had a significantly higher propensity to express CD73 (55%) compared with localized disease (23%, $p=0.002$). The relatively higher prevalence of CD73 expression demonstrated by Yu *et al* might be due to different antibodies used

Figure 1 Representative images demonstrating membranous CD73 expression on tumor cells in the CD73_{negative} (A), CD73_{low} (B) and CD73_{high} (C) groups.

for IHC in their analysis (ab115289; Abcam, Cambridge, Massachusetts, USA). The prevalence of CD73 positivity seen in the current analysis is consistent with our prior study using the same anti-CD73 antibody (clone D7F9A, Cell Signaling Technology).

CD73 expression also correlated with adverse pathologic features and was independently associated with worse prognosis after adjusting for other prognostic features such as tumor grade and AJCC stage in patients with localized RCC. Similarly, in the TCGA dataset, high *NT5E* expression was associated with worse outcomes in patients with AJCC stage IV RCC. These findings suggest

that CD73–adenosine signaling may be a potential driver of aggressive disease in patients with both localized and advanced RCC. As several agents targeting this pathway are in early clinical development, these findings provide compelling rationale for targeting this pathway not only in mRCC but also as perioperative therapy in patients with localized disease and high CD73 expression.

We also characterized CD73 expression in non-clear cell subtypes (papillary n=16, chromophobe n=14) and tumors with a sarcomatoid component (n=10), RCC disease states where expression of inhibitory immune checkpoints has been shown to play a significant immunosuppressive

Figure 2 Distribution of CD73 expression (negative, low and high) by baseline clinical and pathological characteristics. AJCC, American Joint Committee on Cancer; RCC, renal cell carcinoma.

Figure 3 Kaplan-Meier plots of (A) DFS and (B) OS according to CD73 expression in patients with localized renal cell carcinoma. DFS, disease-free survival; OS, overall survival.

role.^{23–26} Approximately 5%–30% of non-ccRCCs express PD-L1, and PD-1/L1 blockade has demonstrated encouraging efficacy in non-clear cell histologies with an ORR of approximately 25%.^{23–24} Similarly, the presence of sarcomatoid differentiation has been associated with increased responsiveness to immune checkpoint inhibition.^{25–26} The CD73–adenosine pathway could be a complementary therapeutic target in these histological subtypes where there exists an even greater clinical need for more effective therapies. We observed CD73 positivity in 39% of our non-ccRCC samples with 24% demonstrating high CD73 expression. In our dataset, 90% of tumors with a sarcomatoid component demonstrated CD73 expression, and 50% were CD73_{high}. Although limited by the small sample sizes and in need of validation, to our knowledge, this is the first study characterizing CD73 expression in non-clear cell and sarcomatoid RCC.

In the TCGA dataset, we also found a strong correlation between CD73 (*NT5E*), CD39 (*ENTPD1*) and A2AR (*ADORA2A*) expression, and Treg gene expression signature. CD73 and CD39 activation on tumor and stromal cells generates extracellular adenosine, which exerts an immunosuppressive effect antagonistic to PD-1 inhibitors. Adenosine activates the high-affinity A2AR receptor, which in turn inhibits infiltrating NK cells and cytotoxic T lymphocyte activity and increases Treg proliferation.^{11–27–30} In addition to tumor cells, CD73, CD39, and

Table 3 Multivariable analysis comparing DFS and OS between CD73 expression groups in patients with localized RCC

CD73 expression groups	DFS			OS		
	N	Event (n)	5-year DFS, % (95% CI)	Adjusted HR (95% CI)*	Adjusted P value	Adjusted HR (95% CI)*
CD73 _{negative}	82	32	75 (63 to 83)	Reference		Reference
CD73 _{low}	12	7	50 (21 to 74)	1.22 (0.50 to 2.98)	0.659	0.78 (0.27 to 2.30)
CD73 _{high}	13	10	42 (15 to 67)	2.72 (1.27 to 5.85)	0.010	2.59 (1.15 to 5.84)
CD73 _{high} vs CD73 _{low+negative}				2.89 (1.37 to 6.07)	0.005	2.86 (1.30 to 6.29)

Results in bold indicate statistically significant results

*Adjusted for Fuhrman nuclear grade (G1–2 vs G3–4) and AJCC stage (stage I/II vs III/IV). Multivariable models excluded four patients with unknown grade or stage. AJCC, American Joint Committee on Cancer; DFS, disease-free survival; OS, overall survival; RCC, renal cell carcinoma.

Figure 4 Genomic and clinical correlates of CD73 (*NT5E*), A2aR (*ADORA2A*) and CD39 (*ENTPD1*) gene expression in the TCGA cohort. AJCC, American Joint Committee on Cancer; FPKM, fragments per kilobase million; Int, intermediate; OS, overall survival.

A2AR can also be expressed on infiltrating immune cells such as Tregs and effector T cells with resultant autocrine production of adenosine further promoting immunosuppression through Treg proliferation.^{27 31} The strong correlation of *NT5E*, *ENTPD1*, and *ADORA2A* expression with the Treg signature observed in our analysis supports the findings of prior preclinical studies¹¹ and provides additional mechanistic support for the immunosuppressive role of CD73–adenosine signaling in RCC.

Adenosine signaling also mediates the recruitment and proliferation of granulocytic MDSCs.¹² Fong *et al* reported that a gene expression signature incorporating genes encoding CXCR2 ligands (eg, *CXCL1*, *CXCL2*, *CXCL3*, *CXCL5*, and *CXCL6*) and mediators of neutrophil and MDSC biology (*ILB*, *IL1B*, and *PTGS2*) could

be a potential biomarker of CD73–adenosine signaling.³² The myeloid-inflammation gene signature used in our study includes several genes included in the adenosine signature (*CXCL1*, *CXCL2*, *CXCL3*, *CXCL8*, and *PTGS2*). However, we did not find a correlation between *NT5E* expression and myeloid inflammation gene expression signature. Further, there was no significant difference in MDSC signature between low and high *NT5E* expression groups ($p = 3.50 \times 10^{-1}$). It is possible that CD73 protein expression or cell type-specific expression (tumor vs immune cell) might have differential effects on these signatures, which was not captured in the bulk RNA-seq data derived from the TCGA dataset.

Preclinical evidence also suggests that CD73–adenosine signaling may promote angiogenesis through vasodilation, release of proangiogenic factors such as VEGF, and recruitment of endothelial progenitor cells.^{14–16} While we did not observe a correlation between *NT5E* expression and angiogenesis marker genes, high expression of *ADORA2A* was strongly associated with high angiogenesis gene expression signature. Dysregulation of the proangiogenic HIF pathway is a critical oncogenic driver in RCC, and A2AR has been identified as a downstream proangiogenic target unique to HIF-2 α .^{33 34} A2AR activation can result in increased angiogenesis through decreased secretion of thrombospondin-1 (TSP-1) and by promoting differentiation of macrophages to the M2 phenotype resulting in increased expression of proangiogenic factors such as VEGF, IL-10 and nitric oxide synthase.^{35–37} Our findings, in corroboration with mechanistic insights from preclinical studies,^{11 28 29 34 38 39} support the hypothesis that adenosine signaling contributes to both tumor immune evasion and angiogenesis in RCC.

Several agents targeting the CD73–adenosine pathway are in clinical development either as monotherapy or in combination with approved anti-PD-1/L1 agents across several cancers, including mRCC. In a first in human, phase I dose-escalation study, the oral competitive A2AR inhibitor ciferadenant (CPI-444) was evaluated as monotherapy and in combination with atezolizumab in patients with advanced refractory cancers.³² Among patients with treatment-refractory mRCC, ciferadenant demonstrated encouraging efficacy, and disease control lasting at least 6 months was seen in 39% of patients treated with the combination. Multiple ongoing trials are investigating adenosine pathway blockade via inhibition of CD73, A2AR, and CD39 as monotherapy and combinations (eg, NCT04148937, NCT03549000, NCT03454451, NCT03835949, NCT03884556, NCT03381274, and NCT04336098). Evaluation of predictive biomarkers such as CD73 expression or adenosine pathway gene expression signatures has the potential to optimize patient selection.

Our study has limitations inherent to institutional retrospective analyses. The relatively small number of patients with non-clear cell histology and de novo mRCC limited our investigation of the prognostic significance of CD73 expression in these subgroups. CD73 expression on

stromal cells or infiltrating immune cells was not assessed, which could provide additional insights into the role of CD73–adenosine signaling in the tumor microenvironment. Although our analysis demonstrates that CD73 expression is associated with poor prognosis at both the mRNA and protein level, additional experimentation will be required to determine the correlation between CD73 protein and mRNA expression. Lastly, protein expression of A2AR and CD39 was not assessed in the current dataset but is worthy of future study of optimal biomarker signatures for adenosine signaling in RCC.

CONCLUSIONS

The CD73–adenosine pathway is an emerging therapeutic target in RCC with extensive preclinical data and growing clinical evidence supporting its role in immunosuppression and angiogenesis. We found that a significant proportion of both clear cell and non-ccRCC tumors express CD73 with a propensity for higher expression in de novo mRCC and sarcomatoid tumors. Higher expression correlated with worse DFS and OS in patients with localized disease independent of stage and grade. In addition, gene expression of CD73, CD39 and A2AR was associated with increased immunosuppressive cell markers and while A2AR expression correlated with the angiogenesis signature in the TCGA cohort. Our findings support the growing investigation of this pathway in advanced RCC.

Author affiliations

¹University of Oklahoma Health Sciences Center, Stephenson Cancer Center, Oklahoma City, Oklahoma, USA

²Lank Center for Genitourinary Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts, USA

³University of Utah, Huntsman Cancer Institute, Salt Lake City, Utah, USA

⁴Department of Data Sciences, Dana Farber Cancer Institute, Boston, Massachusetts, USA

⁵Brigham and Women's Hospital, Boston, Massachusetts, USA

⁶Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA

⁷Oklahoma Medical Research Foundation, Oklahoma City, Oklahoma, USA

⁸Dana-Farber Cancer Institute, Boston, Massachusetts, USA

Twitter Abhishek Tripathi @AbhiTrip87, Toni K Choueiri @DrChoueiri and Lauren C Harshman @LaurenCHarshman

Acknowledgements We thank Abhishek A Chakraborty, PhD, at the Cleveland Clinic's Lerner College of Medicine for providing the Von Hippel-Lindau (VHL) proficient and deficient UMRC2 cells used for positive and negative controls for the immunohistochemistry.

Funding NCI SPORE Grant no: P50 CA101942.

Competing interests AT: advisory role for Foundation Medicine, Pfizer; research funding to institution from EMD Serono, Bayer, Clovis Oncology, Aravive Inc, WindMIL therapeutics, Corvus Pharmaceuticals; EL, WX, AF, JAS, ESG, GB, JHF, CG, CM and LT: none; NM-C: support for research travel from Pfizer and Ipsen, and consulting fees from BMS and Bayer; XXW: research funding to institution from Bristol-Myers Squibb (BMS); MG: research funding from BMS and Merck; BAM: payment for consulting with Bayer, Astellas, Astra Zeneca, Seattle Genetics, Exelixis, Nektar, Pfizer, Janssen, Genentech, Eisai and EMD Serono, and received research support to Dana Farber Cancer Institute (DFCI) from BMS, Calithera, Exelixis, Seattle Genetics. TKC: supported in part by the Dana-Farber/Harvard Cancer Center Kidney SPORE and Program, the Kohlberg Chair at Harvard Medical School and the Trust Family, Michael Brigham, and Loker Pinard Funds for Kidney Cancer Research at DFCI. Research (institutional and personal): AstraZeneca, Alexion, Bayer, BMS/ER Squibb and sons LLC, Cerulean, Eisai, Foundation Medicine Inc, Exelixis,

Ipsen, Tracoon, Genentech, Roche, Roche Products Limited, F. Hoffmann-La Roche, GlaxoSmithKline, Lilly, Merck, Novartis, Peloton, Pfizer, Prometheus Labs, Corvus, Calithera, Analysis Group, Sanofi/Aventis, Takeda. Honoraria: AstraZeneca, Alexion, Sanofi/Aventis, Bayer, BMS/ER Squibb and sons LLC, Cerulean, Eisai, Foundation Medicine Inc, Exelixis, Genentech, Roche, Roche Products Limited, F. Hoffmann-La Roche, GlaxoSmithKline, Merck, Novartis, Peloton, Pfizer, EMD Serono, Prometheus Labs, Corvus, Ipsen, Up-to-Date, NCCN, Analysis Group, NCCN, Michael J. Hennessy (MJH) Associates, Inc (Healthcare Communications Company with several brands such as OnClive, PeerView and PER), Research to Practice, L-path, Kidney Cancer Journal, Clinical Care Options, Platform Q, Navinata Healthcare, Harborside Press, American Society of Medical Oncology, NEJM, Lancet Oncology, Heron Therapeutics, Lilly Oncology. Consulting or advisory Role: AstraZeneca, Alexion, Sanofi/Aventis, Bayer, Bristol Myers-Squibb/ER Squibb and sons LLC, Cerulean, Eisai, Foundation Medicine Inc., Exelixis, Genentech, Heron Therapeutics, Lilly, Roche, GlaxoSmithKline, Merck, Novartis, Peloton, Pfizer, EMD Serono, Prometheus Labs, Corvus, Ipsen, Up-to-Date, NCCN, Analysis Group, Pionyr, Tempest, Lilly Ventures. Stock ownership: Pionyr, Tempest. Other present or past leadership roles: Director of GU Oncology Division at Dana-Farber and past President of medical Staff at Dana-Farber), member of NCCN Kidney panel and the GU Steering Committee, past chairman of the Kidney Cancer Association Medical and Scientific Steering Committee), KidneyCan Advisory board, Kidney cancer Research Summit co-chair (2019-). Patents, royalties or other intellectual properties: related to biomarkers of immune checkpoint blockers. Travel, accommodations, expenses, in relation to consulting, advisory roles, or honoraria: Medical writing and editorial assistance support may have been funded by Communications companies funded by pharmaceutical companies (ClinicalThinking, Envision Pharma Group, Fishawack Group of Companies, Health Interactions, Parexel, Oxford PharmaGenesis, and others). The institution (Dana-Farber Cancer Institute) may have received additional independent funding of drug companies or/and royalties potentially involved in research around the subject matter. CV provided upon request for scope of clinical practice and research. Mentored several non-US citizens on research projects with potential funding (in part) from non-US sources/Foreign Components. NA: consultancy to Astellas, Astra Zeneca, Bayer, Bristol Myers Squibb, Clovis, Eisai, Eli Lilly, EMD Serono, Exelixis, Foundation Medicine, Genentech, Janssen, Merck, Nektar, Novartis, Pfizer, Pharmacyclics, and Seattle Genetics. Research funding to my institution: Astra Zeneca, Bavarian Nordic, Bayer, Bristol Myers Squibb, Calithera, Celldex, Clovis, Eisai, Eli Lilly, EMD Serono, Exelixis, Genentech, Glaxo Smith Kline, Immunomedics, Janssen, Medivation, Merck, Nektar, New Link Genetics, Novartis, Pfizer, Prometheus, Rexahn, Roche, Sanofi, Seattle Genetics, Takeda, and Tracoon. DFM: Honorarium for consultancy from BMS, Pfizer, Merck, Alkermes Inc, EMD Serono, Eli Lilly, Iovance and Eisai. Research support from BMS, Merck, Genentech, Pfizer, Exelixis, X4 Pharma, Alkermes. SS: reports receiving commercial research grants from Bristol-Myers Squibb, AstraZeneca, Exelixis and Novartis; is a consultant/advisory board member for Merck, AstraZeneca, Bristol-Myers Squibb, CRISPR Therapeutics AG, AACR, and NCI; and receives royalties from Biogenex. LCH: Reports consulting fees from Genentech, Dendreon, Pfizer, Medivation/Astellas, Exelixis, Bayer, Kew Group, Corvus, Merck, Novartis, BMS, Michael J Hennessy Associates (Healthcare Communications Company and several brands such as OnLive and PER), Jounce, EMD Serrano, ASIM CME, and Ology Medical Education. Research funding from Bayer, Sotio, BMS, Merck, Takeda, Dendreon/Valient, Janssen, Medivation/Astellas, Genentech, Pfizer, Endocyte (Novartis). Support for research travel from Bayer and Genentech. Currently employed by Surface Oncology.

Patient consent for publication Not required.

Ethics approval Institutional ethics committee approval was obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. The datasets generated during and/or analyzed during the current study are available from the corresponding author on request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Abhishek Tripathi <http://orcid.org/0000-0002-5198-0673>

Nieves Martinez-Chanza <http://orcid.org/0000-0002-2340-7415>

Toni K Choueiri <http://orcid.org/0000-0002-9201-3217>

Neeraj Agarwal <http://orcid.org/0000-0003-1076-0428>

Lauren C Harshman <http://orcid.org/0000-0002-7636-1588>

REFERENCES

- Motzer RJ, Tannir NM, McDermott DF, *et al.* Nivolumab plus ipilimumab versus sunitinib in advanced renal-cell carcinoma. *N Engl J Med* 2018;378:1277–90.
- Tannir NM, Frontera OA, Hammers HJ, *et al.* Thirty-month follow-up of the phase III CheckMate 214 trial of first-line nivolumab + ipilimumab (N+I) or sunitinib (S) in patients (PTS) with advanced renal cell carcinoma (aRCC). *JCO* 2019;37:547.
- Rini BI, Plimack ER, Stus V, *et al.* Pembrolizumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med* 2019;380:1116–27.
- Motzer RJ, Penkov K, Haanen J, *et al.* Avelumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med* 2019;380:1103–15.
- Zimmermann H. 5'-Nucleotidase: molecular structure and functional aspects. *Biochem J* 1992;285 (Pt 2):345–65.
- Sitkovsky MV, Kjaergaard J, Lukashev D, *et al.* Hypoxia-adenosinergic immunosuppression: tumor protection by T regulatory cells and cancerous tissue hypoxia. *Clin Cancer Res* 2008;14:5947–52.
- Synnestvedt K, Furuta GT, Comerford KM, *et al.* Ecto-5'-Nucleotidase (CD73) regulation by hypoxia-inducible factor-1 mediates permeability changes in intestinal epithelia. *J Clin Invest* 2002;110:993–1002.
- Semenza GL. Hypoxia-Inducible factors in physiology and medicine. *Cell* 2012;148:399–408.
- Yegutkin GG. Nucleotide- and nucleoside-converting ectoenzymes: important modulators of purinergic signalling cascade. *Biochim Biophys Acta* 2008;1783:673–94.
- Takayama H, Trenn G, Sitkovsky MV. Locus of inhibitory action of cAMP-dependent protein kinase in the antigen receptor-triggered cytotoxic T lymphocyte activation pathway. *J Biol Chem* 1988;263:2330–6.
- Ohta A, Kini R, Ohta A, *et al.* The development and immunosuppressive functions of CD4(+) CD25(+) FoxP3(+) regulatory T cells are under influence of the adenosine-A2A adenosine receptor pathway. *Front Immunol* 2012;3:190.
- Ryzhov S, Novitskiy SV, Goldstein AE, *et al.* Adenosinergic regulation of the expansion and immunosuppressive activity of CD11b+Gr1+ cells. *J Immunol* 2011;187:6120–9.
- Csóka B, Selmeczy Z, Koscsó B, *et al.* Adenosine promotes alternative macrophage activation via A2A and A2B receptors. *Faseb J* 2012;26:376–86.
- Smits P, Williams SB, Lipson DE, *et al.* Endothelial release of nitric oxide contributes to the vasodilator effect of adenosine in humans. *Circulation* 1995;92:2135–41.
- Feoktistov I, Ryzhov S, Zhong H, *et al.* Hypoxia modulates adenosine receptors in human endothelial and smooth muscle cells toward an A2B angiogenic phenotype. *Hypertension* 2004;44:649–54.
- Montesinos MC, Shaw JP, Yee H, *et al.* Adenosine A(2A) receptor activation promotes wound neovascularization by stimulating angiogenesis and vasculogenesis. *Am J Pathol* 2004;164:1887–92.
- Tripathi A, Johnston SE, Zhao YD, *et al.* CD73 expression in primary and metastatic renal cell carcinoma (RCC). *Journal of Clinical Oncology* 2018;36:643.
- Cancer Genome Atlas Research Network. Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature* 2013;499:43–9.
- Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 2015;31:166–9.
- McDermott DF, Huseini MA, Atkins MB, *et al.* Clinical activity and molecular correlates of response to atezolizumab alone or in combination with bevacizumab versus sunitinib in renal cell carcinoma. *Nat Med* 2018;24:749–57.
- Charoentong P, Finotello F, Angelova M, *et al.* Pan-Cancer Immunogenomic analyses reveal Genotype-Immunophenotype relationships and predictors of response to checkpoint blockade. *Cell Rep* 2017;18:248–62.
- Yu YI, Wang W, Song L, *et al.* Ecto-5'-Nucleotidase expression is associated with the progression of renal cell carcinoma. *Oncol Lett* 2015;9:2485–94.
- Choueiri TK, Fay AP, Gray KP, *et al.* PD-L1 expression in nonclear-cell renal cell carcinoma. *Ann Oncol* 2014;25:2178–84.
- McGregor BA, McKay RR, Braun DA, *et al.* Results of a multicenter phase II study of Atezolizumab and bevacizumab for patients with metastatic renal cell carcinoma with variant histology and/or Sarcomatoid features. *J Clin Oncol* 2020;38:63–70.
- Rini BI, Huseini M, Atkins MB, *et al.* Molecular correlates differentiate response to atezolizumab (atezo) + bevacizumab (bev) vs sunitinib (sun): results from a phase III study (IMmotion151) in untreated metastatic renal cell carcinoma (mRCC). *Annals of Oncology* 2018;29:viii724–5.
- McDermott DF, Choueiri TK, Motzer RJ, *et al.* CheckMate 214 post-hoc analyses of nivolumab plus ipilimumab or sunitinib in IMDC intermediate/poor-risk patients with previously untreated advanced renal cell carcinoma with sarcomatoid features. *JCO* 2019;37:4513.
- Chambers AM, Wang J, Lupo KB, *et al.* Adenosinergic signaling alters natural killer cell functional responses. *Front Immunol* 2018;9:2533.
- Sitkovsky MV, Lukashev D, Apasov S, *et al.* Physiological control of immune response and inflammatory tissue damage by hypoxia-inducible factors and adenosine A2A receptors. *Annu Rev Immunol* 2004;22:657–82.
- Clayton A, Al-Taei S, Webber J, *et al.* Cancer exosomes express CD39 and CD73, which suppress T cells through adenosine production. *J Immunol* 2011;187:676–83.
- Wang L, Fan J, Thompson LF, *et al.* CD73 has distinct roles in nonhematopoietic and hematopoietic cells to promote tumor growth in mice. *J Clin Invest* 2011;121:2371–82.
- Deaglio S, Dwyer KM, Gao W, *et al.* Adenosine generation catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune suppression. *J Exp Med* 2007;204:1257–65.
- Fong L, Hotson A, Powderly JD, *et al.* Adenosine 2A receptor blockade as an immunotherapy for treatment-refractory renal cell cancer. *Cancer Discov* 2020;10:40–53.
- Ricketts CJ, Crooks DR, Linehan WM. Targeting HIF2α in clear-cell renal cell carcinoma. *Cancer Cell* 2016;30:515–7.
- Ahmad A, Ahmad S, Glover L, *et al.* Adenosine A2A receptor is a unique angiogenic target of HIF-2α in pulmonary endothelial cells. *Proc Natl Acad Sci U S A* 2009;106:10684–9.
- Desai A, Victor-Vega C, Gadangi S, *et al.* Adenosine A2A receptor stimulation increases angiogenesis by down-regulating production of the antiangiogenic matrix protein thrombospondin 1. *Mol Pharmacol* 2005;67:1406–13.
- Ferrante CJ, Pinhal-Enfield G, Elson G, *et al.* The adenosine-dependent angiogenic switch of macrophages to an M2-like phenotype is independent of interleukin-4 receptor alpha (IL-4Rα) signaling. *Inflammation* 2013;36:921–31.
- Grinberg S, Hasko G, Wu D, *et al.* Suppression of PLCβ2 by endotoxin plays a role in the adenosine A(2A) receptor-mediated switch of macrophages from an inflammatory to an angiogenic phenotype. *Am J Pathol* 2009;175:2439–53.
- Ohta A, Gorelik E, Prasad SJ, *et al.* A2A adenosine receptor protects tumors from antitumor T cells. *Proc Natl Acad Sci U S A* 2006;103:13132–7.
- Csóka B, Himer L, Selmeczy Z, *et al.* Adenosine A2A receptor activation inhibits T helper 1 and T helper 2 cell development and effector function. *Faseb J* 2008;22:3491–9.