



Immunogenic cell death pathway polymorphisms for predicting oxaliplatin efficacy in metastatic colorectal cancer

Hiroyuki Arai ¹, Yi Xiao ², Fotios Loupakis,³ Natsuko Kawanishi,¹ Jingyuan Wang,¹ Francesca Battaglin,¹ Shivani Soni,¹ Wu Zhang,¹ Christoph Mancao,⁴ Bodour Salhia,⁵ Shannon M Mumenthaler,⁶ Daniel J Weisenberger,⁷ Gangning Liang,⁸ Chiara Cremolini,⁹ Alfredo Falcone,⁹ Joshua Millstein,² Heinz-Josef Lenz¹

To cite: Arai H, Xiao Y, Loupakis F, *et al.* Immunogenic cell death pathway polymorphisms for predicting oxaliplatin efficacy in metastatic colorectal cancer. *Journal for ImmunoTherapy of Cancer* 2020;**8**:e001714. doi:10.1136/jitc-2020-001714

Accepted 11 October 2020

ABSTRACT

Background Immunogenic cell death (ICD) is a tumor cell death involving both innate and adaptive immune responses. Given published findings that oxaliplatin, but not irinotecan, drives ICD, we investigated whether single nucleotide polymorphisms (SNPs) in the ICD pathway are associated with the efficacy of oxaliplatin-based chemotherapy in metastatic colorectal cancer (mCRC).

Methods Two randomized clinical trials data were analyzed: discovery cohort, FOLFOX/bevacizumab arm (MAVERICC); validation cohort, FOLFOXIRI/bevacizumab arm (TRIBE); and two control cohorts, FOLFIRI/bevacizumab arms (both trials). Genomic DNA extracted from blood samples was genotyped. Ten SNPs in the ICD pathway were tested for associations with clinical outcomes.

Results In total, 648 patients were included. In the discovery cohort, three SNPs were significantly associated with clinical outcomes in univariate analysis: *CALR* rs1010222 with progression-free survival (G/G vs any A, HR=0.61, 95% CI 0.43–0.88), *ANXA1* rs1050305 with overall survival (OS) (A/A vs any G, HR=1.87, 95% CI 1.04–3.35), and *LRP1* rs1799986 with OS (C/C vs any T, HR=1.69, 95% CI 1.07–2.70). Multivariate analysis confirmed the trend, but statistical significance was not reached. In the validation cohort, *ANXA1* rs1050305, and *LRP1* rs1799986 were validated to have the significant associations with clinical outcome. No significant associations of these SNPs were observed in the two control cohorts. Treatment-by-SNP interaction test confirmed the predictive values.

Conclusions The predictive utility of ICD-related SNPs for the efficacy of oxaliplatin-based chemotherapy was demonstrated, warranting further validation studies to be translated into personalized treatment strategies using conventional cytotoxic agents in mCRC.

BACKGROUND

The current standard of care for patients with microsatellite-stable metastatic colorectal cancer (mCRC) consists of the combination

of a cytotoxic chemotherapy backbone and a monoclonal antibody based on *RAS*/*BRAF* testing.^{1 2} In clinical practice, there are two choices for cytotoxic agent, namely oxaliplatin or irinotecan in combination with 5-fluorouracil, and two for monoclonal antibodies targeting vascular endothelial growth factor (VEGF) or epidermal growth factor receptor (EGFR). Most research has focused on the efficacy of monoclonal antibodies, leading to the successful identification of predictive and prognostic biomarkers for EGFR inhibitors such as *RAS* and *BRAF* mutation status and primary tumor sidedness.^{3–6} However, there are no predictive biomarkers for the selection of oxaliplatin or irinotecan which have distinct mechanisms of action: oxaliplatin is a platinum compound forming intra-strand and inter-strand cross-links; irinotecan is a topoisomerase I (TOP1) inhibitor trapping TOP1 cleavage complex with DNA replication and transcription. The choice of these two drugs remains an important clinical question for optimizing treatment in individual patients.

Immunogenic cell death (ICD) is a novel concept of tumor cell death that engages both innate and adaptive immune responses, thereby conferring an additional immunogenic antitumor effect to cytotoxic agents.^{7–9} However, most chemotherapeutics kill tumor cells in a non-immunogenic manner, and only a few cytotoxic agents can prime antitumor immunity by inducing ICD.^{10 11} The mechanisms of ICD induction are characterized by the ability to induce endoplasmic reticulum stress, which in turn leads to the release of damage-associated molecular patterns (DAMPs) from dying tumor cells and the subsequent



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Heinz-Josef Lenz;
lenz@usc.edu

activation of pattern recognition receptors (PRRs) of host innate immune cells such as dendritic cells.¹² The engagement of innate immune responses cross-primes adaptive responses controlled by cytotoxic T lymphocytes through proinflammatory cytokine release, potentially leading to the continuous elimination of tumor cells.^{12,13} Oxaliplatin has been recently well-recognized as an ICD inducer, similarly as other drugs such as anthracyclines and bortezomib.⁷ In a preclinical study, CRC cells treated with oxaliplatin exhibited increased pre-apoptotic exposure of calreticulin (CALR) and post-apoptotic release of high-mobility group box 1 (HMGB1), which are the signals of DAMPs required for ICD induction.¹⁴ By contrast, irinotecan lacks the mechanism of action that drives ICD, and thus, its cytotoxic effects depend on non-immunogenic mechanisms.^{10,15} These findings suggested that genetic variants in the ICD pathway are related to the efficacy of oxaliplatin but not irinotecan, prompting us to test whether these novel biomarkers will be useful for patient selection. Therefore, we investigated the predictive utility of single nucleotide polymorphisms (SNPs) in ICD-related genes encoding essential DAMPs and PRRs using genetic and clinical data from two first-line randomized clinical trials comparing oxaliplatin- (or oxaliplatin-containing) and irinotecan-based treatments in patients with mCRC.

METHODS

Patient population and study design

This study included patients with mCRC enrolled in two first-line randomized trials: MAVERICC (NCT01374425)¹⁶ and TRIBE (NCT00719797).¹⁷ In the MAVERICC trial, patients were randomized to treatment with either

FOLFOX plus bevacizumab or FOLFIRI plus bevacizumab. In the TRIBE trial, patients were randomized to the FOLFOXIRI plus bevacizumab or FOLFIRI plus bevacizumab arm. Patients without sufficient peripheral whole blood samples for analyzes were excluded from our study. To assess whether the ICD pathway is specifically related to the efficacy of oxaliplatin, we set the cohorts as follows: (1) discovery cohort, FOLFOX plus bevacizumab arm in MAVERICC; (2) validation cohort, FOLFOXIRI plus bevacizumab arm in TRIBE; (3) control cohort 1, FOLFIRI plus bevacizumab arm in MAVERICC; and (4) control cohort 2, FOLFIRI plus bevacizumab arm in TRIBE (figure 1). All patients provided informed consent for molecular research prior to study enrollment.

Genotyping and selecting polymorphisms

Genomic DNA was extracted from peripheral whole blood collected before treatment initiation using a QIAmp Kit (Qiagen, Valencia, California, USA) in accordance with the manufacturer's protocol (www.qiagen.com). The OncoArray of 530 K SNPs was used for genotyping (Illumina, San Diego, California, USA).¹⁸ After genotyping, we planned this exploratory analysis focusing on five genes encoding DAMPs or PRRs related to chemotherapy-induced ICD,¹² including three DAMP-encoding genes (*CALR*, *HMGB1*, and *ANXA1*) and two PRR-encoding genes (*LRP1* and *P2RX7*) (online supplementary table S1). Although *FPRI*, *TLR3*, and *TLR4* are also key PRR-encoding genes, these genes were not included in this study because they were not predictive of oxaliplatin efficacy in patients with CRC in another large biomarker study.¹⁹ The candidate SNPs for this study were arbitrarily selected according to the following criteria: (1)




Figure 1 Consort diagram. BEV, bevacizumab; SNP, single nucleotide polymorphism.

Table 1 Associations between ICD-related SNPs and clinical outcomes

Genotype	N	TR			PFS			OS		
		ORR (%)	P value*	P	Univariate analysis			Univariate analysis		
					HR (95% CI)	HR (95% CI)	P value†	HR (95% CI)	HR (95% CI)	P value†
CALR rs1010222										
Discovery cohort										
G/G	72	66.2	0.48	0.008	1	1	0.10	1	1	0.07
Any A	87	60.7			0.61 (0.43 to 0.88)	0.67 (0.42 to 1.07)		0.67 (0.44 to 1.04)	0.78 (0.46 to 1.35)	
Validation cohort										
G/G	44	71.4	0.33	0.63	1	1	0.61	1	1	0.76
Any A	65	62.3			0.90 (0.57 to 1.41)	0.87 (0.52 to 1.46)		0.93 (0.60 to 1.44)	0.79 (0.49 to 1.29)	
Control cohort 1										
G/G	81	67.5	1.00	0.98	1	1	0.88	1	1	0.41
Any A	82	67.5			0.99 (0.69 to 1.44)	0.96 (0.58 to 1.60)		0.81 (0.49 to 1.34)	0.97 (0.50 to 1.89)	
Control cohort 2										
G/G	90	63.6	0.16	0.37	1	1	0.56	1	1	0.75
Any A	119	53.9			1.16 (0.84 to 1.58)	1.11 (0.79 to 1.56)		1.05 (0.77 to 1.44)	0.96 (0.69 to 1.33)	
ANXA1 rs1050305										
Discovery cohort										
A/A	138	63.4	0.99	0.26	1	1	0.47	1	1	0.03
Any G	22	63.6			1.33 (0.81 to 2.21)	1.29 (0.66 to 2.51)		1.87 (1.04 to 3.35)	1.96 (0.93 to 4.12)	
Validation cohort										
A/A	91	70.6	0.03	0.009	1	1	0.17	1	1	<0.001
Any G	18	44.4			2.17 (1.20 to 3.92)	1.62 (0.83 to 3.16)		2.69 (1.56 to 4.61)	2.00 (1.07 to 3.72)	
Control cohort 1										
A/A	128	66.9	0.67	0.72	1	1	0.85	1	1	0.82
Any G	33	71.0			1.08 (0.70 to 1.67)	1.07 (0.54 to 2.11)		1.07 (0.60 to 1.92)	0.81 (0.33 to 1.98)	
Control cohort 2										
A/A	184	57.6	0.70	0.92	1	1	0.72	1	1	0.23
Any G	31	61.3			1.02 (0.66 to 1.58)	1.09 (0.68 to 1.74)		0.76 (0.48 to 1.19)	0.67 (0.42 to 1.09)	
LRP1 rs1799986										
Discovery cohort										
C/C	111	64.2	0.99	0.22	1	1	0.53	1	1	0.03
Any T	44	64.3			1.30 (0.85 to 1.97)	1.18 (0.71 to 1.94)		1.69 (1.07 to 2.70)	1.55 (0.87 to 2.74)	
Validation cohort										
C/C	74	63.9	0.52	0.35	1	1	0.005	1	1	0.98

Continued

Table 1 Continued

Genotype	N	TR			PFS			OS					
		ORR (%)	P value*	P value†	Univariate analysis		Multivariate analysis		Univariate analysis			Multivariate analysis	
					HR (95% CI)	P value‡	HR (95% CI)	P value‡	HR (95% CI)	P value‡	HR (95% CI)	P value‡	
Any T	24	71.4			1.30 (0.75 to 2.25)		2.71 (1.37 to 5.37)		0.99 (0.58 to 1.71)		1.30 (0.70 to 2.40)		
Control cohort 1													
C/C	126	65.9	0.28		0.35	1	0.79	1	1	0.68	1	0.38	
Any T	33	75.8			1.23 (0.79 to 1.92)		0.93 (0.52 to 1.66)		1.13 (0.63 to 2.02)		0.71 (0.33 to 1.53)		
Control cohort 2													
C/C	162	58.9	0.45		0.34	1	0.59	1	1	0.57	1	0.66	
Any T	44	52.4			0.83 (0.57 to 1.22)		0.90 (0.60 to 1.34)		0.89 (0.61 to 1.31)		1.10 (0.73 to 1.66)		

Significant values are indicated in bold characters.

*P values were based on χ^2 test.

†P values were based on log-rank test for PFS and OS in the univariate analysis.

‡P values were based on Wald test in the multivariate Cox proportional hazards regression model.

ICD, immunogenic cell death; ORR, overall response rate; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism; TR, tumor response.

minor allele frequency in Caucasians of at least 10% in the Ensemble Genome Browser (<https://www.ensembl.org>), (2) having potential biological functions based on public databases (<https://snpinfo.niehs.nih.gov>; <https://www.ncbi.nlm.nih.gov>), and (3) tag SNPs chosen from HapMap genotype data with an r^2 threshold of 0.8 (<https://snpinfo.niehs.nih.gov>). In total, 10 SNPs were selected, as presented in online supplementary table S2.

Statistical analysis

Selected SNPs were evaluated for their associations with tumor response, progression-free survival (PFS), and overall survival (OS). The overall response rate (ORR) was calculated as the percentage of patients with either a complete or partial response using the Response Evaluation Criteria in Solid Tumors version 1.1. PFS was defined as the time from randomization to disease progression or death from any cause. OS was defined as the time from randomization to death from any cause. Patients who experienced no events were censored at the last follow-up date. The correlation between each SNP and ORR was examined using the χ^2 test. To test the association between each SNP and PFS or OS, univariate and multivariate analyzes using the Cox proportional hazards regression model and the log-rank test were performed. In the multivariate analyzes, the following study-specific adjusted covariates were used: the TRIBE trial included sex, age, Eastern Cooperative Oncology Group (ECOG) performance status (PS), primary tumor site, liver-limited disease, adjuvant chemotherapy, *BRAF* status, and *RAS* status; and the MAVERICC trial included ethnicity, sex, age, ECOG PS, primary tumor site, primary tumor resected, number of metastases, and *KRAS* status. SNPs were coded using a dominant genetic model for the number of variant alleles. To formally assess the predictive value, the treatment-by-SNP interaction was tested within each trial using the HR calculated in the multivariate analyzes. All analyzes were two-sided at a significance level of 0.05 and were performed using SAS V.9.4 software.

RESULTS

Patient characteristics

In total, 648 patients were included in this study (figure 1). Some characteristics were unbalanced between the trial cohorts. In particular ECOG PS 0, left-sided primary tumor, metastasis in ≤ 2 organs, and primary tumor resection were more prevalent in the TRIBE trial arms than in the MAVERICC trial arms (online supplementary table S3). The median follow-up and survival time in each cohort are summarized in online supplementary table S4.

Predictive value of ICD-related SNPs

All tested associations between each SNP and clinical outcomes are presented in online supplemental tables S5–8. Although several significant associations were observed in the discovery or validation cohort, few associations were confirmed in control cohort 1 or 2.




Figure 2 OS of patients with ANXA1 rs1050305 variants. (A) Discovery cohort (FOLFOX plus bevacizumab arm in MAVERICC), (B) validation cohort (FOLFOXIRI plus bevacizumab arm in TRIBE), (C) control cohort 1 (FOLFIRI plus bevacizumab arm in MAVERICC), and (D) control cohort 2 (FOLFIRI plus bevacizumab arm in TRIBE). OS, overall survival.

In particular, three SNPs displayed significant associations with outcome in the univariate analysis in the discovery cohort (table 1, figure 2). First, patients carrying any A allele of *CALR* rs1010222 had a better PFS than those with a G/G genotype (HR=0.61, 95% CI 0.43–0.88, $p=0.008$). Second, patients carrying any G allele of *ANXA1* rs1050305 had worse OS than those with an A/A genotype (HR=1.87, 95% CI 1.04–3.35, $p=0.03$). Finally, patients carrying any T allele of *LRP1* rs1799986 had worse OS than those with a C/C genotype (HR=1.69, 95% CI 1.07–2.70, $p=0.03$). Multivariate analysis confirmed these trends, but none of the associations reached statistical significance. In the validation cohort, two SNPs were confirmed to have significant associations with clinical outcome (table 1, figure 2). Namely, patients carrying any G allele of *ANXA1* rs1050305 had worse ORR, PFS (revealed in univariate analysis), and OS (revealed in both univariate and multivariate analyses) than those with an A/A genotype; and patients carrying any T allele of *LRP1* rs1799986 had

worse PFS (revealed in multivariate analysis) than those with a C/C genotype, even though this clinical endpoint was not same as that in the discovery cohort. Of note, in the two control cohorts, no significant associations were observed between these SNPs (*CALR* rs1010222, *ANXA1* rs1050305, and *LRP1* rs1799986) and clinical outcomes (table 1, figure 2).

The results of the treatment-by-SNP interaction test are presented in table 2. In total, six SNPs exhibited significant interactions with treatment: *HMGB1* rs1360485 (PFS in TRIBE), *ANXA1* rs1050305 (OS in TRIBE), *LRP1* rs1799986 (PFS in TRIBE), *LRP1* rs11172113 (both PFS and OS in TRIBE), *P2RX7* rs208294 (OS in MAVERICC and OS in TRIBE), and *P2RX7* rs1718119 (OS in MAVERICC).

DISCUSSION

This is the first study to identify genetic variants in ICD-related pathways that may predict the efficacy of oxaliplatin-based first-line chemotherapy in patients with

Table 2	Treatment-by-single nucleotide polymorphism interaction test											
	MAVERICC						TRIBE					
	Hazard ratio on PFS			Hazard ratio on OS			Hazard ratio on PFS			Hazard ratio on OS		
	FOLFOLFOX+ BEV	FOLFIRI+ BEV	Interaction p value	FOLFOLFOX+ BEV	FOLFIRI+ BEV	Interaction p value	FOLFOLFOX+ BEV	FOLFIRI+ BEV	Interaction p value	FOLFOLFOX+ BEV	FOLFIRI+ BEV	Interaction p value
CALR rs1049481												
T/T	1	1	0.87	1	1	0.98	1	1	0.11	1	1	0.42
Any G	0.85 (0.54– 1.36)	0.91 (0.54– 1.51)		1.03 (0.60– 1.76)	0.92 (0.47– 1.78)		0.69 (0.39– 1.21)	1.14 (0.81– 1.62)		0.74 (0.45– 1.24)	0.97 (0.69– 1.36)	
CALR rs1010222												
G/G	1	1	0.41	1	1	0.41	1	1	0.34	1	1	0.57
Any A	0.67 (0.42– 1.07)	0.96 (0.58– 1.60)		0.78 (0.46– 1.35)	0.97 (0.50– 1.89)		0.87 (0.52– 1.46)	1.11 (0.79– 1.56)		0.79 (0.49– 1.29)	0.96 (1.69– 1.33)	
HMGB1 rs1045411												
C/C	1	1	0.35	1	1	0.58	1	1	0.07	1	1	0.50
Any T	1.43 (0.85– 2.41)	1.09 (0.63– 1.89)		1.46 (0.79– 2.69)	1.20 (0.58– 2.48)		1.93 (1.06– 3.53)	1.11 (0.74– 1.65)		1.03 (0.59– 1.80)	0.95 (0.65– 1.38)	
HMGB1 rs1412125												
T/T	1	1	0.87	1	1	0.51	1	1	0.30	1	1	0.11
Any C	0.74 (0.43– 1.28)	0.78 (0.45– 1.35)		0.80 (0.43– 1.48)	1.06 (0.52– 2.16)		1.22 (0.64– 2.30)	0.98 (0.69– 1.40)		1.14 (0.64– 2.01)	0.83 (0.58– 1.17)	
HMGB1 rs1360485												
T/T	1	1	0.28	1	1	0.73	1	1	0.04	1	1	0.42
Any C	1.45 (0.89– 2.38)	1.03 (0.62– 1.69)		1.38 (0.78– 2.43)	1.19 (0.62– 2.25)		1.70 (0.95– 3.06)	1.06 (0.74– 1.50)		0.93 (0.54– 1.60)	0.98 (0.70– 1.37)	
ANXA1 rs1050305												
A/A	1	1	0.82	1	1	0.15	1	1	0.16	1	1	0.002
Any G	1.29 (0.66– 2.51)	1.07 (0.54– 2.11)		1.96 (0.93– 4.12)	0.81 (0.33– 1.98)		1.62 (0.83– 3.16)	1.09 (0.68– 1.74)		2.00 (1.07– 3.72)	0.67 (0.42– 1.09)	
LRP1 rs1799986												
C/C	1	1	0.66	1	1	0.23	1	1	0.03	1	1	0.55
Any T	1.18 (0.71– 1.94)	0.93 (0.52– 1.66)		1.55 (0.87– 2.74)	0.71 (0.33– 1.53)		2.71 (1.37– 5.37)	0.90 (0.60– 1.34)		1.30 (0.70– 2.40)	1.10 (0.73– 1.66)	

Continued

Table 2 Continued

	MAVERICC						TRIBE					
	Hazard ratio on PFS			Hazard ratio on OS			Hazard ratio on PFS			Hazard ratio on OS		
	FOLFOX+ BEV	FOLFIRI+ BEV	Interaction p value	FOLFOX+ BEV	FOLFIRI+ BEV	Interaction p value	FOLFOXIRI+ BEV	FOLFIRI+ BEV	Interaction p value	FOLFOXIRI+ BEV	FOLFIRI+ BEV	Interaction p value
<i>LRP1</i> rs11172113												
T/T	1	1	0.31	1	1	0.73	1	1	0.004	1	1	0.004
Any C	1.05 (0.63– 1.74)	1.22 (0.72– 2.06)		0.97 (0.55– 1.69)	0.66 (0.34– 1.26)		0.55 (0.31– 0.98)	1.35 (0.96– 1.89)		0.58 (0.34– 0.96)	1.20 (0.87– 1.67)	
<i>P2RX7</i> rs208294												
C/C	1	1	0.33	1	1	0.046	1	1	0.62	1	1	0.02
Any T	0.93 (0.55– 1.58)	1.26 (0.72– 2.22)		0.62 (0.33– 1.16)	1.84 (0.84– 4.06)		0.92 (0.54– 1.58)	1.02 (0.71– 1.47)		0.54 (0.33– 0.89)	0.92 (0.64– 1.32)	
<i>P2RX7</i> rs1718119												
G/G	1	1	0.66	1	1	0.03	1	1	0.85	1	1	0.77
Any A	1.02 (0.62– 1.70)	1.26 (0.74– 2.14)		0.97 (0.53– 1.78)	2.57 (1.16– 5.67)		0.77 (0.45– 1.31)	0.87 (0.62– 1.21)		0.69 (0.42– 1.15)	0.83 (0.60– 1.15)	

Significant values are indicated in bold characters. Each HR is presented with a point estimate and a 95% CI (in parentheses) calculated in the multivariate analyses. BEV, bevacizumab; OS, overall survival; PFS, progression-free survival.

mCRC. No predictive biomarkers for the personalized use of cytotoxic agents in mCRC have been identified so far. DNA damage response genes and the transcriptome-based consensus molecular subtype have been examined in biomarker research, but their predictive value has not been validated.^{16–20} Gray *et al* reported that genetic variants of three PRR genes (*FPR1*, *TLR3*, and *TLR4*) were not related to the efficacy of oxaliplatin in patients with CRC in the adjuvant setting (using data from the SCOT trials) and the first-line setting for patients with metastasis (using data from the COIN/COIN-B trials).¹⁹ However, DAMPs are also essential tumor-sided factors for ICD induction, whereas PRRs are host-side factors. Therefore, in our study, we extensively assessed the functional variants of both DAMPs and PRRs. According to our hypothesis, ICD-related SNPs were significantly associated with the efficacy of oxaliplatin-based combination. These findings imply the linkage between ICD pathway and oxaliplatin efficacy, which can be coupled with the potential synergy of oxaliplatin and immunotherapy.^{21–22}

Our findings identified *ANXA1* rs1050305 and *LRP1* rs1799986 as SNPs exhibiting significant associations with the efficacy of oxaliplatin but not irinotecan as confirmed in the discovery and validation cohorts. *ANXA1* is one of the DAMPs released from the cytoplasm of dying cells, and it interacts with *FPR1*, which is expressed on dendritic cells.²³ The extracellular accumulation of *ANXA1* constitutes a hallmark of ICD.¹² *ANXA1* rs1050305 is a synonymous SNP located at an exonic splicing enhancer/silencer site that disrupts splicing activity and causes alternative splicing (<https://snpinfo.niehs.nih.gov/snpinfo/snpfunc.html>). *LRP1* is a PRR activated via docking of cell surface-exposed CALR, which is another hallmark of chemotherapy-induced ICD.^{12–24} *LRP1* rs1799986 is a missense SNP that may participate in splicing regulation. This SNP is reported to be associated with susceptibility to Alzheimer's disease, supporting its functional significance.²⁵ Of note, while the predictive value of *ANXA1* rs1050305 was consistently validated on OS, that of *LRP1* rs1799986 was observed on inconsistent clinical outcomes between the discovery and validation cohorts: the discovery cohort showed a significant association with OS, and the validation cohort showed that with PFS. This discrepancy might be due to some unidentified factors such as different contribution of second-line treatment between two cohorts, which might affect the associations between biomarker and clinical outcome.

Our study had great strengths in its design, featuring discovery, validation, and control cohorts obtained from two randomized clinical trials. The validated effects of the SNPs on clinical outcomes in the discovery and validation cohorts, lacking of that in the two control cohorts, strongly supports the predictive value of *ANXA1* rs1050305 and *LRP1* rs1799986 specifically for oxaliplatin efficacy. The multiple significant associations observed in the discovery and validation cohorts but not in the control cohorts further indicate that the ICD pathway plays an important role in the efficacy of oxaliplatin but

not irinotecan. Another strength was that the treatment-by-SNP interaction test confirmed the predictive value of *ANXA1* rs1050305 and *LRP1* rs1799986 and highlighted the predictive utility of 6 out of 10 tested SNPs.²⁶

This study had two key limitations. First, this was a prospective-retrospective study; thus, the results need to be validated in prospective clinical trials. According to published guidelines, one or more validation studies are required to reach a sufficient level of evidence for medical utility.^{27–28} Second, the validation cohort consisted of patients treated with FOLFOXIRI plus bevacizumab, which is both an oxaliplatin-containing and irinotecan-containing regimen. To validate the different effects of SNPs on oxaliplatin and irinotecan efficacy, it will be better to use a treatment arm containing only oxaliplatin-based regimens as the validation cohort. However, our study provides reliable evidence describing the specific connection of ICD-related SNPs with the efficacy of “oxaliplatin-containing” treatment based on the use of “oxaliplatin-non-containing” cohorts as controls.

In conclusion, our study provides the first evidence that genetic variants in the ICD pathway are significantly associated with the efficacy of oxaliplatin-based, but not irinotecan-based, first-line chemotherapy in patients with mCRC. Further translational studies are warranted to validate our novel findings and realize the personalized use of conventional cytotoxic chemotherapeutic agents.

Author affiliations

¹Division of Medical Oncology, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

²Department of Preventive Medicine, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

³Clinical and Experimental Oncology Department, Medical Oncology Unit 1, Veneto Institute of Oncology IOV-IRCCS, Padua, Italy

⁴Oncology Biomarker Development, Genentech Inc, Basel, Switzerland

⁵Department of Translational Genomics, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁶Lawrence J. Ellison Institute for Transformative Medicine, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁷Department of Biochemistry and Molecular Medicine, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁸Department of Urology, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁹Department of Translational Medicine, Division of Medical Oncology, University of Pisa, Pisa, Italy

Acknowledgements We thank all patients who contributed to this study.

Contributors HA and H-JL primarily planned, designed, and drafted the manuscript. YX and JM did statistical analyses. All authors made substantial contributions to data collection and drafting the manuscript. All authors read and approved the final manuscript.

Funding This work was supported by the National Cancer Institute [P30CA 014089 to HJL], Gloria Borges WunderGlo Foundation, Dhont Family Foundation, Victoria and Philip Wilson Research Fund, San Pedro Peninsula Cancer Guild, and Daniel Butler Research Fund.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval All patients provided informed consent for molecular research prior to study enrollment. The study protocol was approved by the Institutional

Review Board of each participating institution and was conducted in accordance with the tenets of the Declaration of Helsinki as well as the Good Clinical Practice and REMARK guidelines.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. De-identified datasets analyzed in the current study are available from the corresponding author on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Hiroyuki Arai <http://orcid.org/0000-0002-1886-7070>

Yi Xiao <http://orcid.org/0000-0003-3816-461X>

REFERENCES

- 1 NCCN clinical practice guidelines in oncology: colorectal cancer. version 4, 2020. Available: https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf [Accessed 11 July 2020].
- 2 Van Cutsem E, Cervantes A, Adam R, *et al*. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol* 2016;27:1386–422.
- 3 Sepulveda AR, Hamilton SR, Allegra CJ, *et al*. Molecular biomarkers for the evaluation of colorectal cancer: guideline from the American Society for clinical pathology, College of American pathologists, association for molecular pathology, and the American Society of clinical oncology. *J Clin Oncol* 2017;35:1453–86.
- 4 Arnold D, Lueza B, Douillard J-Y, *et al*. Prognostic and predictive value of primary tumour side in patients with Ras wild-type metastatic colorectal cancer treated with chemotherapy and EGFR directed antibodies in six randomized trials. *Ann Oncol* 2017;28:1713–29.
- 5 Tejpar S, Stintzing S, Ciardiello F, *et al*. Prognostic and predictive relevance of primary tumor location in patients with Ras wild-type metastatic colorectal cancer: retrospective analyses of the crystal and FIRE-3 trials. *JAMA Oncol* 2017;3:194–201.
- 6 Loupakis F, Yang D, Yau L, *et al*. Primary tumor location as a prognostic factor in metastatic colorectal cancer. *J Natl Cancer Inst* 2015;107.
- 7 Kroemer G, Galluzzi L, Kepp O, *et al*. Immunogenic cell death in cancer therapy. *Annu Rev Immunol* 2013;31:51–72.
- 8 Galluzzi L, Buqué A, Kepp O, *et al*. Immunological effects of conventional chemotherapy and targeted anticancer agents. *Cancer Cell* 2015;28:690–714.
- 9 Galluzzi L, Senovilla L, Zitvogel L, *et al*. The secret ally: immunostimulation by anticancer drugs. *Nat Rev Drug Discov* 2012;11:215–33.
- 10 Obeid M, Tesniere A, Ghiringhelli F, *et al*. Calreticulin exposure dictates the immunogenicity of cancer cell death. *Nat Med* 2007;13:54–61.
- 11 Garg AD, More S, Rufo N, *et al*. Trial Watch: immunogenic cell death induction by anticancer chemotherapeutics. *Oncoimmunology* 2017;6:e1386829.
- 12 Galluzzi L, Buqué A, Kepp O, *et al*. Immunogenic cell death in cancer and infectious disease. *Nat Rev Immunol* 2017;17:97–111.
- 13 Ruan H, Leibowitz BJ, Zhang L, *et al*. Immunogenic cell death in colon cancer prevention and therapy. *Mol Carcinog* 2020;59:783–93.
- 14 Tesniere A, Schlemmer F, Boige V, *et al*. Immunogenic death of colon cancer cells treated with oxaliplatin. *Oncogene* 2010;29:482–91.
- 15 Pozzi C, Cuomo A, Spadoni I, *et al*. The EGFR-specific antibody cetuximab combined with chemotherapy triggers immunogenic cell death. *Nat Med* 2016;22:624–31.
- 16 Parikh AR, Lee F-C, Yau L, *et al*. MAVERICC, a randomized, Biomarker-stratified, phase II study of mFOLFOX6-Bevacizumab versus FOLFIRI-Bevacizumab as first-line chemotherapy in metastatic colorectal cancer. *Clin Cancer Res* 2019;25:2988–95.
- 17 Loupakis F, Cremolini C, Masi G, *et al*. Initial therapy with FOLFOXIRI and bevacizumab for metastatic colorectal cancer. *N Engl J Med* 2014;371:1609–18.
- 18 Amos CI, Dennis J, Wang Z, *et al*. The OncoArray Consortium: a network for understanding the genetic architecture of common cancers. *Cancer Epidemiol Biomarkers Prev* 2017;26:126–35.
- 19 Gray V, Briggs S, Palles C, *et al*. Pattern recognition receptor polymorphisms as predictors of oxaliplatin benefit in colorectal cancer. *J Natl Cancer Inst* 2019;111:828–36.
- 20 Aderka D, Stintzing S, Heinemann V. Explaining the unexplainable: discrepancies in results from the CALGB/SWOG 80405 and FIRE-3 studies. *Lancet Oncol* 2019;20:e274–83.
- 21 Boku N, Ryu MH, Oh D-Y, *et al*. LBA7_PR nivolumab plus chemotherapy versus chemotherapy alone in patients with previously untreated advanced or recurrent gastric/gastroesophageal junction (G/GEJ) cancer: ATTRACTION-4 (ONO-4538-37) study. *Annals of Oncology* 2020;31:S1192.
- 22 Moehler M, Shitara K, Garrido M, *et al*. LBA6_PR Nivolumab (nivo) plus chemotherapy (chemo) versus chemo as first-line (1L) treatment for advanced gastric cancer/gastroesophageal junction cancer (GC/GEJC)/esophageal adenocarcinoma (EAC): First results of the CheckMate 649 study. *Annals of Oncology* 2020;31:S1191.
- 23 Baracco EE, Petrazzuolo A, Kroemer G. Assessment of annexin A1 release during immunogenic cell death. *Methods Enzymol* 2019;629:71–9.
- 24 Garg AD, Krysko DV, Verfaillie T, *et al*. A novel pathway combining calreticulin exposure and ATP secretion in immunogenic cancer cell death. *Embo J* 2012;31:1062–79.
- 25 Wang Y, Liu S, Wang J, *et al*. Association between LRP1 C766T polymorphism and Alzheimer's disease susceptibility: a meta-analysis. *Sci Rep* 2017;7:8435.
- 26 Ballman KV. Biomarker: predictive or prognostic? *J Clin Oncol* 2015;33:3968–71.
- 27 Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst* 2009;101:1446–52.
- 28 Polley M-YC, Freidlin B, Korn EL, *et al*. Statistical and practical considerations for clinical evaluation of predictive biomarkers. *J Natl Cancer Inst* 2013;105:1677–83.

Table S1. DAMPs and PRRs related to chemotherapy-induced ICD

DAMPs	PRRs
ANXA1	FPR1
ATP	P2RX7 P2RY2
CALR	LRP1
Cellular RNA	TLR3
HMGB1	TLR4

Abbreviations: DAMPs, damage-associated molecular patterns; ICD, immunogenic cell death; PRRs, pattern recognition receptors.

Table S2. Characteristics of selected SNPs in ICD pathway

ICD-related gene	Chromosome	SNP (rs number)	Location	MAF	Base exchange	Function
CALR	19	rs1049481	3'UTR	0.37	G>T	Tag SNP, transcription factor binding site
		rs1010222	5'near	0.31	G>A	Transcription factor binding site
HMGB1	13	rs1045411	3'UTR	0.27	C>T	Tag SNP, miRNA binding site Associated with outcome in lung cancer patients treated with platinum agent Changing the mRNA levels
		rs1412125	5'near	0.48	T>C	Transcription factor binding site Associated with platinum-based chemotherapy response in lung cancer
		rs1360485	3'near	0.31	C>T	Inversely associated with EGFR mutations among NSCLC patients who smoked
ANXA1	9	rs1050305	exon	0.10	A>G	Synonymous SNP, splicing regulation
LRP1	12	rs1799986	exon	0.15	C>T	Tag SNP, missense SNP, splicing regulation Associated with susceptibility to Alzheimer's disease
		rs11172113	intron	0.38	C>G	Tag SNP Associated with susceptibility to migraine, syncope, and stroke
P2RX7	12	rs208294	exon	0.44	T>C	Tag SNP, missense SNP, splicing regulation Associated with prognosis of prostate cancer patients
		rs1718119	exon	0.38	A>G	Missense SNP, splicing regulation Associated with risk of obesity, pulmonary tuberculosis, and systemic lupus erythematosus

MAF presents minor allele frequency in Caucasians in the Ensemble Genome Browser (<http://www.ensembl.org>).

Abbreviations: ICD, immunogenic cell death; MAF, minor allele frequency; miRNA, micro RNA; mRNA, messenger RNA; NSCLC, non-small-cell lung cancer; SNP, single nucleotide polymorphism; UTR, untranslated region.

Table S3. Comparison of patient characteristics between cohorts

Characteristics		MAVERICC				TRIBE		P-value
		Total N = 648	N = 324		N = 324			
			FOLFOX+BEV	FOLFIRI+BEV	FOLFOXIRI+BEV	FOLFIRI+BEV		
			(discovery cohort)	(control cohort 1)	(validation cohort)	(control cohort 2)		
		N = 161	N = 163	N = 109	N = 215			
Sex							0.97	
	Male	402	101 (62.7%)	103 (63.2%)	66 (60.6%)	132 (61.4%)		
	Female	246	60 (37.3%)	60 (36.8%)	43 (39.4%)	83 (38.6%)		
Age							0.10	
	≤65	452	117 (72.7%)	101 (62.0%)	78 (71.6%)	156 (72.6%)		
	>65	196	44 (27.3%)	62 (38.0%)	31 (28.4%)	59 (27.4%)		
Performance status							<0.001	
	ECOG 0	450	81 (50.3%)	97 (59.5%)	95 (87.2%)	177 (82.3%)		
	ECOG 1	196	79 (49.1%)	66 (40.5%)	14 (12.8%)	37 (17.2%)		
	Unknown*	2	1 (0.6%)	0 (0%)	0 (0%)	1 (0.5%)		
Primary tumor site							0.007	
	Right-sided	214	64 (39.8%)	67 (41.1%)	30 (27.5%)	53 (24.7%)		
	Left-sided	413	97 (60.2%)	96 (58.9%)	73 (67.0%)	147 (68.4%)		
	Unknown*	21	0 (0%)	0 (0%)	6 (5.5%)	15 (7.0%)		
Number of metastases							<0.001	
	≤2	474	101 (62.7%)	106 (65.0%)	89 (81.7%)	178 (82.8%)		
	>2	174	60 (37.3%)	57 (35.0%)	20 (18.3%)	37 (17.2%)		

	Unknown*	0	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
<hr/>							
Liver limited disease							0.31
	No	220	NA	NA	70 (64.2%)	150 (69.8%)	
	Yes	104	NA	NA	39 (35.8%)	65 (30.2%)	
<hr/>							
Primary tumor resected							<0.001
	No	412	148 (91.9%)	153 (93.9%)	31 (28.4%)	80 (37.2%)	
	Yes	236	13 (8.1%)	10 (6.1%)	78 (71.6%)	135 (62.8%)	
<hr/>							
Adjuvant chemotherapy							0.68
	No	571	146 (90.7%)	143 (87.7%)	94 (86.2%)	188 (87.4%)	
	Yes	77	15 (9.3%)	20 (12.3%)	15 (13.8%)	27 (12.6%)	
	Unknown*	0	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
<hr/>							
RAS status							0.32
	Wildtype	84	NA	NA	34 (31.2%)	50 (23.3%)	
	Mutant	167	NA	NA	57 (52.3%)	110 (51.2%)	
	Unknown*	73	NA	NA	18 (16.5%)	55 (25.6%)	
<hr/>							
BRAF status							0.25
	Wildtype	256	NA	NA	88 (80.7%)	168 (78.1%)	
	Mutant	19	NA	NA	9 (8.3%)	10 (4.7%)	
	Unknown*	49	NA	NA	12 (11.0%)	37 (17.2%)	

*Unknown groups were not included in the analysis.

P-values was estimated by Chi-square test.

Abbreviations: BEV, bevacizumab; ECOG, Eastern Cooperative Oncology Group; NA, not assessed.

Table S4. Follow-up and survival time in each cohort

Time (months)	MAVERICC		TRIBE	
	<i>N</i> = 324		<i>N</i> = 324	
	FOLFOX+BEV <i>N</i> = 161	FOLFIRI+BEV <i>N</i> = 163	FOLFOXIRI+BEV <i>N</i> = 109	FOLFIRI+BEV <i>N</i> = 215
Follow-up				
Median	26.8	23.3	54.5	48.9
PFS				
Median	10.1	12.5	10.8	9.7
OS				
Median	24.7	27.4	26.0	26.2

Abbreviations: BEV, bevacizumab; OS, overall survival; PFS, progression-free survival.

Table S5. Associations between ICD-related SNPs and clinical outcomes in MAVERICC FOLFOX+Bevacizumab arm (discovery cohort)

Genotype	N	TR			PFS					OS				
		Yes	No	P ^a	mPFS (months)	Univariate analysis		Multivariate analysis		mOS (months)	Univariate analysis		Multivariate analysis	
						HR (95% CI)	P ^b	HR (95% CI)	P ^c		HR (95% CI)	P ^b	HR (95% CI)	P ^c
CALR rs1049481														
T/T	59	39 (67.2%)	19 (33.8%)	0.45	10.2	1	0.12	1	0.51	22.7	1	0.27	1	0.93
Any G	101	60 (61.2%)	38 (38.8%)		10.1	0.75 (0.52–1.08)		0.85 (0.54–1.36)		26.1	0.78 (0.50–1.21)		1.03 (0.60–1.76)	
CALR rs1010222														
G/G	72	47 (66.2%)	24 (33.8%)	0.48	9.0	1	0.008	1	0.10	22.5	1	0.07	1	0.38
Any A	87	51 (60.7%)	33 (39.3%)		11.0	0.61 (0.43–0.88)		0.67 (0.42–1.07)		28.8	0.67 (0.44–1.04)		0.78 (0.46–1.35)	
HMGB1 rs1045411														
C/C	89	54 (63.5%)	31 (36.5%)	0.84	11.2	1	0.18	1	0.18	25.5	1	0.56	1	0.23
Any T	63	39 (61.9%)	24 (38.1%)		8.9	1.29 (0.89–1.87)		1.43 (0.85–2.41)		22.8	1.14 (0.73–1.80)		1.46 (0.79–2.69)	
HMGB1 rs1412125														
T/T	33	19 (61.3%)	12 (38.7%)	0.76	11.8	1	0.78	1	0.29	23.2	1	0.92	1	0.48
Any C	128	81 (64.3%)	45 (35.7%)		10.1	1.07 (0.67–1.70)		0.74 (0.43–1.28)		25.5	0.97 (0.58–1.63)		0.80 (0.43–1.48)	
HMGB1 rs1360485														
T/T	75	45 (62.5%)	27 (37.5%)	0.77	12.3	1	0.09	1	0.14	25.8	1	0.23	1	0.26
Any C	86	55 (64.7%)	30 (35.3%)		8.8	1.36 (0.95–1.95)		1.45 (0.89–2.38)		23.9	1.31 (0.84–2.03)		1.38 (0.78–2.43)	
ANXA1 rs1050305														
A/A	138	85 (63.4%)	49 (36.6%)	0.99	11.0	1	0.26	1	0.47	25.5	1	0.03	1	0.10
Any G	22	14 (63.6%)	8 (36.4%)		8.8	1.33 (0.81–2.21)		1.29 (0.66–2.51)		18.7	1.87 (1.04–3.35)		1.96 (0.93–4.12)	
LRP1 rs1799986														
C/C	111	70 (64.2%)	39 (35.8%)	0.99	11.0	1	0.22	1	0.53	25.8	1	0.03	1	0.14
Any T	44	27 (64.3%)	15 (35.7%)		8.5	1.30 (0.85–1.97)		1.18 (0.71–1.94)		16.9	1.69 (1.06–2.70)		1.55 (0.87–2.74)	
LRP1 rs11172113														
T/T	51	35 (71.4%)	14 (28.6%)	0.17	10.2	1	0.50	1	0.85	23.9	1	0.78	1	0.91
Any C	110	65 (60.2%)	43 (39.8%)		10.9	1.14 (0.78–1.68)		1.05 (0.63–1.74)		25.5	0.94 (0.60–1.47)		0.97 (0.55–1.69)	
P2RX7 rs208294														
C/C	41	29 (72.5%)	11 (27.5%)	0.18	11.0	1	0.69	1	0.80	26.1	1	0.47	1	0.14
Any T	120	71 (60.7%)	46 (39.3%)		10.2	1.09 (0.71–1.66)		0.93 (0.55–1.58)		24.8	0.83 (0.51–1.37)		0.62 (0.33–1.16)	
P2RX7 rs1718119														
G/G	60	40 (67.8%)	19 (32.2%)	0.41	11.0	1	0.67	1	0.93	25.5	1	0.67	1	0.92
Any A	101	60 (61.2%)	38 (38.8%)		10.0	1.08 (0.75–1.57)		1.02 (0.62–1.70)		24.3	1.10 (0.71–1.72)		0.97 (0.53–1.78)	

Significant values are indicated in bold characters.

Abbreviations: CI, confidence interval; HR, hazard ratio; ICD, immunogenic cell death; mOS, median overall survival; mPFS, median progression-free survival; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism; TR, tumor response.

^a *P*-values were based on Chi-square test.

^b *P*-values were based on log-rank test for PFS and OS in the univariate analysis.

^c *P*-values were bases on Wald test in the multivariate Cox proportional hazards regression model.

Table S6. Associations between ICD-related SNPs and clinical outcomes in TRIBE FOLFOXIRI+Bevacizumab arm (validation cohort)

Genotype	N	TR			PFS					OS				
		Yes	No	P ^a	mPFS (months)	Univariate analysis		Multivariate analysis		mOS (months)	Univariate analysis		Multivariate analysis	
						HR (95% CI)	P ^b	HR (95% CI)	P ^c		HR (95% CI)	P ^b	HR (95% CI)	P ^c
CALR rs1049481														
T/T	40	27 (71.1%)	11 (28.9%)	0.41	10.3	1	0.73	1	0.19	26.7	1	0.82	1	0.26
Any G	69	41 (63.1%)	24 (36.9%)		11.0	0.92 (0.58–1.47)		0.69 (0.39–1.21)		23.6	0.95 (0.61–1.48)		0.74 (0.45–1.24)	
CALR rs1010222														
G/G	44	30 (71.4%)	12 (28.6%)	0.33	10.3	1	0.63	1	0.61	26.2	1	0.76	1	0.35
Any A	65	38 (62.3%)	23 (37.7%)		11.1	0.90 (0.57–1.41)		0.87 (0.52–1.46)		25.1	0.93 (0.60–1.44)		0.79 (0.49–1.29)	
HMGB1 rs1045411														
C/C	69	41 (62.1%)	25 (37.9%)	0.39	11.9	1	0.06	1	0.03	28.4	1	0.99	1	0.93
Any T	34	22 (71.0%)	9 (29.0%)		9.7	1.57 (0.97–2.53)		1.93 (1.06–3.53)		23.2	1.00 (0.62–1.62)		1.03 (0.59–1.80)	
HMGB1 rs1412125														
T/T	31	19 (65.5%)	10 (34.5%)	0.95	12.1	1	0.69	1	0.54	28.1	1	0.48	1	0.65
Any C	78	49 (66.2%)	25 (33.8%)		10.1	1.11 (0.68–1.80)		1.22 (0.64–2.30)		25.1	1.19 (0.73–1.92)		1.14 (0.64–2.01)	
HMGB1 rs1360485														
T/T	64	39 (63.9%)	22 (36.1%)	0.59	11.9	1	0.05	1	0.07	27.4	1	0.89	1	0.80
Any C	45	29 (69.0%)	13 (31.0%)		9.6	1.56 (0.99–2.46)		1.70 (0.95–3.06)		26.1	0.97 (0.62–1.51)		0.93 (0.54–1.60)	
ANXA1 rs1050305														
A/A	91	60 (70.6%)	25 (29.4%)	0.03	11.3	1	0.009	1	0.17	31.0	1	<0.001	1	0.04
Any G	18	8 (44.4%)	10 (55.6%)		9.5	2.17 (1.20–3.92)		1.62 (0.83–3.16)		17.1	2.69 (1.56–4.61)		2.00 (1.07–3.72)	
LRP1 rs1799986														
C/C	74	46 (63.9%)	26 (36.1%)	0.52	11.3	1	0.35	1	0.005	26.7	1	0.98	1	0.42
Any T	24	15 (71.4%)	6 (28.6%)		8.9	1.30 (0.75–2.25)		2.71 (1.37–5.37)		23.7	0.99 (0.58–1.71)		1.30 (0.70–2.40)	
LRP1 rs11172113														
T/T	42	24 (60.0%)	16 (40.0%)	0.30	10.1	1	0.22	1	0.04	21.6	1	0.18	1	0.03
Any C	67	44 (69.8%)	19 (30.2%)		11.3	0.75 (0.48–1.19)		0.55 (0.31–0.98)		28.4	0.74 (0.48–1.15)		0.58 (0.34–0.96)	
P2RX7 rs208294														
C/C	35	20 (66.7%)	10 (33.3%)	0.93	10.3	1		1	0.76	19.0	1	0.004	1	0.02
Any T	74	48 (65.8%)	25 (34.2%)		11.0	0.78 (0.49–1.25)		0.92 (0.54–1.58)		31.0	0.52 (0.33–0.82)		0.54 (0.33–0.89)	
P2RX7 rs1718119														
G/G	44	29 (67.4%)	14 (32.6%)	0.80	10.3	1	0.67	1	0.33	26.1	1	0.34	1	0.16
Any A	65	39 (65.0%)	21 (35.0%)		10.9	0.91 (0.57–1.43)		0.77 (0.45–1.31)		26.2	0.81 (0.52–1.25)		0.69 (0.42–1.15)	

Significant values are indicated in bold characters.

Abbreviations: CI, confidence interval; HR, hazard ratio; ICD, immunogenic cell death; mOS, median overall survival; mPFS, median progression-free survival; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism; TR, tumor response.

^a *P*-values were based on Chi-square test.

^b *P*-values were based on log-rank test for PFS and OS in the univariate analysis.

^c *P*-values were bases on Wald test in the multivariate Cox proportional hazards regression model.

Table S7. Associations between ICD-related SNPs and clinical outcomes in MAVERICC FOLFIRI+Bevacizumab arm (control cohort 1)

Genotype	N	TR			PFS					OS				
		Yes	No	P ^a	mPFS (months)	Univariate analysis		Multivariate analysis		mOS (months)	Univariate analysis		Multivariate analysis	
						HR (95% CI)	P ^b	HR (95% CI)	P ^c		HR (95% CI)	P ^b	HR (95% CI)	P ^c
CALR rs1049481														
T/T	69	46 (67.6%)	22 (32.4%)	0.97	11.2	1	0.46	1	0.70	26.5	1	0.13	1	0.80
Any G	94	62 (67.4%)	30 (32.6%)		12.7	0.87 (0.59–1.27)		0.91 (0.54–1.51)		31.3	0.68 (0.42–1.12)		0.92 (0.47–1.78)	
CALR rs1010222														
G/G	81	54 (67.5%)	26 (32.5%)	1.00	12.9	1	0.98	1	0.88	26.9	1	0.41	1	0.94
Any A	82	54 (67.5%)	26 (32.5%)		12.6	0.99 (0.69–1.44)		0.96 (0.58–1.60)		27.9	0.81 (0.49–1.34)		0.97 (0.50–1.89)	
HMGB1 rs1045411														
C/C	109	71 (67.0%)	35 (33.0%)	0.81	11.8	1	0.39	1	0.76	27.9	1	0.97	1	0.62
Any T	42	29 (69.0%)	13 (31.0%)		12.8	0.83 (0.54–1.27)		1.09 (0.63–1.89)		32.3	0.99 (0.56–1.75)		1.20 (0.58–2.48)	
HMGB1 rs1412125														
T/T	50	30 (61.2%)	19 (38.8%)	0.26	12.3	1	0.63	1	0.38	32.3	1	0.32	1	0.87
Any C	113	78 (70.3%)	33 (29.7%)		12.6	0.91 (0.61–1.34)		0.78 (0.45–1.35)		27.4	1.32 (0.76–2.30)		1.06 (0.52–2.16)	
HMGB1 rs1360485														
T/T	98	62 (65.3%)	33 (34.7%)	0.47	11.6	1	0.13	1	0.92	27.9	1	0.75	1	0.60
Any C	65	46 (70.8%)	19 (29.2%)		14.8	0.75 (0.51–1.09)		1.03 (0.62–1.69)		27.5	0.92 (0.56–1.53)		1.19 (0.62–2.25)	
ANXA1 rs1050305														
A/A	128	85 (66.9%)	42 (33.1%)	0.67	12.3	1	0.72	1	0.85	27.9	1	0.82	1	0.63
Any G	33	22 (71.0%)	9 (29.0%)		13.8	1.08 (0.70–1.67)		1.07 (0.54–2.11)		27.5	1.07 (0.60–1.92)		0.81 (0.33–1.98)	
LRP1 rs1799986														
C/C	126	81 (65.9%)	42 (34.1%)	0.28	12.8	1	0.35	1	0.79	27.9	1	0.68	1	0.38
Any T	33	25 (75.8%)	8 (24.2%)		12.3	1.23 (0.79–1.92)		0.93 (0.52–1.66)		25.0	1.13 (0.63–2.02)		0.71 (0.33–1.53)	
LRP1 rs11172113														
T/T	60	36 (61.0%)	23 (39.0%)	0.18	12.9	1	0.75	1	0.45	26.9	1	0.60	1	0.21
Any C	103	72 (71.3%)	29 (28.7%)		12.3	1.07 (0.72–1.58)		1.22 (0.72–2.06)		31.3	0.87 (0.52–1.46)		0.66 (0.34–1.26)	
P2RX7 rs208294														
C/C	42	27 (65.9%)	14 (34.1%)	0.79	15.0	1	0.41	1	0.41	-	1	0.03	1	0.11
Any T	121	81 (68.1%)	38 (31.9%)		12.3	1.20 (0.78–1.85)		1.26 (0.72–2.22)		27.5	2.07 (1.05–4.08)		1.84 (0.84–4.06)	
P2RX7 rs1718119														
G/G	72	48 (69.6%)	21 (30.4%)	0.63	14.0	1	0.60	1	0.38	32.3	1	0.05	1	0.01
Any A	91	60 (65.9%)	31 (34.1%)		11.2	1.11 (0.76–1.61)		1.26 (0.74–2.14)		23.8	1.70 (1.01–2.86)		2.57 (1.16–5.67)	

Significant values are indicated in bold characters.

Abbreviations: CI, confidence interval; HR, hazard ratio; ICD, immunogenic cell death; mOS, median overall survival; mPFS, median progression-free survival; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism; TR, tumor response.

^a *P*-values were based on Chi-square test.

^b *P*-values were based on log-rank test for PFS and OS in the univariate analysis.

^c *P*-values were bases on Wald test in the multivariate Cox proportional hazards regression model.

Table S8. Associations between ICD-related SNPs and clinical outcomes in TRIBE FOLFIRI+Bevacizumab arm (control cohort 2)

Genotype	N	TR			PFS					OS				
		Yes	No	P ^a	mPFS (months)	Univariate analysis		Multivariate analysis		mOS (months)	Univariate analysis		Multivariate analysis	
						HR (95% CI)	P ^b	HR (95% CI)	P ^c		HR (95% CI)	P ^b	HR (95% CI)	P ^c
CALR rs1049481														
T/T	81	49 (62.0%)	30 (38.0%)	0.39	10.8	1	0.40	1	0.45	28.0	1	0.94	1	0.85
Any G	131	71 (55.9%)	56 (44.1%)		9.5	1.15 (0.83–1.57)		1.14 (0.81–1.62)		26.1	1.01 (0.74–1.39)		0.97 (0.69–1.36)	
CALR rs1010222														
G/G	90	56 (63.6%)	32 (36.4%)	0.16	10.8	1	0.37	1	0.56	27.4	1	0.75	1	0.79
Any A	119	62 (53.9%)	53 (46.1%)		9.5	1.16 (0.84–1.58)		1.11 (0.79–1.56)		26.2	1.05 (0.77–1.44)		0.96 (1.69–1.33)	
HMGB1 rs1045411														
C/C	138	78 (58.2%)	56 (41.8%)	0.58	9.9	1	0.87	1	0.62	25.5	1	0.51	1	0.79
Any T	59	35 (62.5%)	21 (37.5%)		9.5	0.97 (0.68–1.38)		1.11 (0.74–1.65)		29.8	0.89 (0.63–1.26)		0.95 (0.65–1.38)	
HMGB1 rs1412125														
T/T	63	37 (60.7%)	24 (39.3%)	0.64	9.7	1	0.77	1	0.92	22.5	1	0.43	1	0.29
Any C	152	84 (57.1%)	63 (42.9%)		9.7	1.05 (0.76–1.46)		0.98 (0.69–1.40)		28.0	0.88 (0.63–1.22)		0.83 (0.58–1.17)	
HMGB1 rs1360485														
T/T	134	76 (58.5%)	54 (41.5%)	0.91	9.9	1	0.78	1	0.76	25.8	1	0.41	1	0.89
Any C	81	45 (57.7%)	33 (42.3%)		9.7	0.95 (0.69–1.32)		1.06 (0.74–1.50)		29.8	0.88 (0.64–1.20)		0.98 (0.70–1.37)	
ANXA1 rs1050305														
A/A	184	102 (57.6%)	75 (42.4%)	0.70	9.9	1	0.92	1	0.72	26.2	1	0.23	1	0.09
Any G	31	19 (61.3%)	12 (38.7%)		9.2	1.02 (0.66–1.58)		1.09 (0.68–1.74)		31.6	0.76 (0.48–1.19)		0.67 (0.42–1.09)	
LRP1 rs1799986														
C/C	162	93 (58.9%)	65 (41.1%)	0.45	9.5	1	0.34	1	0.59	26.2	1	0.57	1	0.66
Any T	44	22 (52.4%)	20 (47.6%)		11.5	0.83 (0.57–1.22)		0.90 (0.60–1.34)		27.1	0.89 (0.61–1.31)		1.10 (0.73–1.66)	
LRP1 rs11172113														
T/T	98	56 (58.9%)	39 (41.1%)	0.84	10.4	1	0.32	1	0.09	26.9	1	0.87	1	0.27
Any C	117	65 (57.5%)	48 (42.5%)		9.5	1.17 (0.86–1.59)		1.35 (0.96–1.89)		25.5	1.03 (0.76–1.39)		1.20 (0.87–1.67)	
P2RX7 rs208294														
C/C	64	37 (59.7%)	25 (40.3%)	0.77	11.0	1	0.38	1	0.92	27.1	1	0.89	1	0.66
Any T	151	84 (57.5%)	62 (42.5%)		9.5	1.16 (0.83–1.61)		1.02 (0.71–1.47)		26.3	0.98 (0.70–1.36)		0.92 (0.64–1.32)	
P2RX7 rs1718119														
G/G	90	48 (55.8%)	38 (44.2%)	0.56	9.5	1	0.91	1	0.41	27.1	1	0.74	1	0.26
Any A	125	73 (59.8%)	49 (40.2%)		10.6	0.98 (0.72–1.34)		0.87 (0.62–1.21)		26.3	0.95 (0.70–1.29)		0.83 (0.60–1.15)	

Significant values are indicated in bold characters.

Abbreviations: CI, confidence interval; HR, hazard ratio; ICD, immunogenic cell death; mOS, median overall survival; mPFS, median progression-free survival; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism; TR, tumor response.

^a *P*-values were based on Chi-square test.

^b *P*-values were based on log-rank test for PFS and OS in the univariate analysis.

^c *P*-values were bases on Wald test in the multivariate Cox proportional hazards regression model.