models. AMP immunogens are efficiently delivered to lymph nodes, where innate and adaptive immune responses are generated.

Methods Female, 6 to 8-week-old C57BL/6J and BALB/c mice and 37-week-old C57BL/6J mice received two or more doses of benchmark (alum or CpG) or AMP-modified vaccines, comprised of Spike RBD protein and AMP-CpG adjuvant, subcutaneously injected into the tail base in two-week intervals. Antigen was dose spared to determine if AMP-CpG would maintain the immune response. Cellular immune responses were determined via ELISpot analysis of IFN γ production by splenocytes, intracellular cytokine staining of peripheral blood and lung-resident T-cells, and flowcytometric bead array analysis of Th1/2/17 cytokines. Humoral immune responses were determined via blood serum ELISAs to determine sera antibody binding titers, and pseudoviral neutralization assays for comparison to human convalescent serum.

Results Compared to alum, AMP immunization induced 29-fold higher antigen-specific T cells which produced multiple Th1 cytokines and trafficked into lung parenchyma. Antibody responses favored Th1 isotypes (IgG2bc, IgG3) and potently neutralized Spike-2-ACE2 receptor binding, with titers >100-fold higher than the natural immune response from convalescent COVID-19 patients; responses were maintained despite 10-fold dose-reduction in Spike antigen. Both cellular and humoral immune responses were preserved in aged mice.

Conclusions ELI-005 exhibits the qualities of an optimal SARS-CoV-2 vaccine, which should (1) induce robust and durable CD8+ and CD4+ T cell responses, (2) elicit high magnitude neutralizing antibodies, (3) produce Th1 bias in the elicited antibody and T cell responses, (4) potentially expand pre-existing cross-reactive T cells, (5) enable dose-sparing of required immunogens to improve the speed and cost of broad vaccination campaigns, and (6) be efficacious in elderly populations. These advantages merit clinical translation to SARS-CoV-2 and other protein subunit vaccines.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0479>

480

PRELIMINARY EVALUATION OF A NOVEL CORONAVIRUS VACCINE (CORVAX) USING ELECTROPORATION OF PLASMID DNA ENCODING A STABILIZED PREFUSION SARS-COV-2 SPIKE PROTEIN ALONE OR WITH TRANSFECTION OF PLASMID IL-12

¹Shawn Jensen*, ²Christopher Twitty, ³Christopher Paustian, ³Madelein Laws, ³Glenna McDonnell, ¹Keith Wegmann, ¹Tarsem Moudgil, ¹Michael Afentoulis, ²Mia Han, ²Kellie Malloy Foerter, ²David Canton, ²Jack Lee, ²Bianca Nguyen, ²John Rodriguez, ²Kim Jaffe, ¹Brian Piening, ¹Carlo Bifulco, ²Daniel O'Connor, ¹Walter Urba, ¹Rom Leidner, ³Traci Hilton, ¹Hong-Ming Hu, ¹Bernard Fox. ¹Earle A. Chiles Research Institute, Prov, Portland, OR, USA; ²OncoSec Medical Incorporated, San Diego, San Diego, CA, USA; ³UbiVac, Portland, Oregon, Portland, OR, USA

Background SARS-CoV-2 (CoV2) has precipitated a global pandemic and the effectiveness of standard vaccine strategies to induce potent and persistent immunity to CoV2 is in question, particularly for the elderly. This problem is not dissimilar to what we have struggled with in our quest to induce immunity to cancer antigens, where vaccine-induced anti-cancer immune responses can be weak. Here, we describe a novel vaccine approach which leverages electroporation (EP) of a plasmid encoding a prefusion stabilized CoV2 spike protein

(CORVax). As IL-12 has been shown to augment the efficacy of immunotherapy in aged mice,¹ we have initiated studies to evaluate if plasmid IL-12 (TAVO™) can similarly augment anti-CoV2 immune responses in young mice and have planned studies in aged animals.

Methods A prefusion stabilized CoV2 spike plasmid expression vector was constructed, a master cell bank generated and clinical-grade plasmid manufactured. C57BL/6 and BALB/c were vaccinated via intramuscular (IM) and/or intradermal (ID) injection followed immediately by EP of plasmids encoding the CoV2 spike protein with or without plasmid-encoded murine IL-12 on days 1 and 14 or 21. Mice were followed for >120 days to assess safety. Splenocytes and serum were harvested at different time points to interrogate virus-specific cellular responses as well anti-spike IgG1/IgG2 antibody titers. A surrogate viral neutralization test (sVNT) assessed serum blockade of soluble hACE2R binding to immobilized CoV2 spike.

Results Preliminary data shows that EP of CORVax alone or combined with IL-12 was safe. EP of CORVax was able to elicit anti-Spike IgG antibodies (IC50 = 1/2112), as well as IgG antibodies targeting the receptor binding domain of the Spike protein (IC50 = 1/965) approximately 40 days after the booster vaccination. In 2 of 2 experiments, CORVax combined with IL-12 significantly (P<0.0001) increased the sVNT titers at 2 months, but this benefit was lost by 3 months.

Conclusions Early preclinical data shows that EP of CORVax can induce IgG responses to CoV2 Spike and the receptor binding domain (RBD) as well as apparent viral neutralizing activity. The addition of IL-12, at least transiently, increased sVNT titer. We plan to investigate alternate vaccine boosting strategies while extending these studies into aged animals and initiate a clinical trial in the near future.

REFERENCES

1. Ruby CE, Weinberg AD. OX40-Enhanced tumor rejection and effector T cell differentiation decreases with age. *J Immunol* 2009;**182**:1481–9. <https://doi.org/10.4049/jimmunol.182.3.1481>.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0480>

481

IMPACT OF COVID-19 ON CANCER PATIENTS RECEIVING IMMUNE CHECKPOINT INHIBITORS

¹Kevin Tyan*, ¹Ai-Tram Bui, ²Anita Giobbie-Hurder, ²Isaac Klein, ²Michael Manos, ³Leyre Zubiri, ³Kerry Reynolds, ⁴Shilpa Grover, ⁴Gerald Weinhouse, ²Patrick Ott, ⁴Nicole LeBoeuf, ²Osama Rahma. ¹Harvard Medical School, Somerville, MA, USA; ²Dana-Farber Cancer Institute, Boston, MA, USA; ³Massachusetts General Hospital, Boston, MA, USA; ⁴Brigham and Women's Hospital, Boston, MA, USA

Background There are conflicting data regarding the vulnerability of cancer patients receiving immune checkpoint inhibitors (ICIs) to COVID-19 infection.^{1–3} In addition, immune-related adverse events (irAEs) driven in part by cytokine dysregulation could parallel the cytokine storm implicated in COVID-19. We sought to evaluate the impact of COVID-19 infection on irAEs and mortality in cancer patients receiving ICIs.

Methods We performed a retrospective matched cohort study of 25 patients receiving ICIs within one year of a confirmed COVID-19 diagnosis between March 20, 2020 and June 3, 2020 at the Dana-Farber Cancer Institute/Mass General Brigham network. Cases were matched 1:1 with controls without ICI use based on age, sex, and use of non-ICI anti-cancer