Immune cell biology

492

INTEGRATION OF HIGH DIMENSIONAL DATASETS IN AN IMMUNOCOMPETENT MAMMARY MOUSE MODEL REVEALS PATHWAYS OF TOLERANCE AND RESISTANCE TO IMMUNE CHECKPOINT BLOCKADE

¹Lin Ma, ²Jian-Hua Mao, ¹Mary Helen Barcellos-Hoff, ¹Jade Moore*. ¹University of California, San Francisco, San Francisco, CA, USA; ²Lawrence Berkeley National Laboratory, Berkeley, CA, USA

Background Checkpoint inhibitors can induce robust and durable responses in a subset of patients. Extending this benefit to more patients could be facilitated by better understanding of how interacts with immune cells with the tumor microenvironment, which is a critical barrier to control both local and systemic disease. The composition and pattern of the immune infiltrate associates with the likelihood of response to immunotherapy. Inflamed tumors that exhibit a brisk immune cell infiltrate are responsive, while those in which immune cells are completely or partially excluded are not. Transforming growth factor β (TGF β) is immunosuppressive and associated with the immune excluded phenotype.

Methods Using an immune competent mammary tumor derived transplant (mTDT) model recently developed in our lab, exhibits inflamed, excluded or deserts immune infiltrate phenotypes based on localization of CD8 lymphocytes. Using whole transcriptome deep sequencing, cytof, and PET-CT imaging, we evaluated the tumor, microenvironment, and immune pathway activation among immune infiltrate phenotypes.

Results Three distinct inflamed tumors phenotypes were identified: 'classically' inflamed characterized by pathway evidence of increased CD8+ T cells and decreased PD-L1 expression, inflamed tumors with pathways indicative of neovascularization and STAT3 signaling and reduced T cell mobilization, and an inflamed tumor with increased immunosuppressive myeloid phenotypes. Excluded tumors were characterized by TGF β gene expression and pro-inflammatory cytokine signaling (e.g. TNF α , IL1 β), associated with decreased leukocytes homing and increased immune cell death of cells. We visualized and quantified TGF β activity using PET-CT imaging of ⁸⁹Zr-fresolimumab, a TGF β neutralizing antibody. TGF β activity was significantly increased in excluded tumors compared to inflamed or desert tumors, which was supported by quantitative pathology (Perkin Elmer) of its canonical signaling target, phosphorylated SMAD2 (pSMAD2). pSMAD2 was positively correlated with PD-L1 expression in the stroma of excluded tumors. In contrast, in inflamed tumors, TGF β activity positively correlated with increased F4/80 positive macrophages and negatively correlated with expression of PD-L1. CyTOF analysis of tumor and spleen immune phenotypes revealed increased trafficking of myeloid cells in mice bearing inflamed tumors compared to excluded and deserts.

Conclusions The immunocompetent mTDT provides a model that bridges the gap between the immune landscape and tumor microenvironment. Integration of these high-dimensional data with further studies of response to immunotherapies will help to identify tumor features that favor response to treatment or the means to convert those that are unresponsive.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0492>

493

TIRED AND HUNGRY: A POTENTIAL ROLE FOR CD47 IN T CELL EXHAUSTION

¹Levi Mangarin, ¹Caillan Liu, ²Roberta Zappasodi, ³Pamela Holland, ¹Jedd Wolchok, ¹Taha Merghoub, ¹Chien-Huan Weng*, ¹Chien-Huan Weng, ¹David Schroder. ¹Memorial Sloan-Kettering Cancer Center, New York, NY, USA; ²Weill Cornell Medical College, New York, NY, USA; ³Surface Oncology, Inc., Cambridge, MA, USA

Background Multiple suppressive mechanisms within the tumor microenvironment are capable of blunting anti-tumor T cell responses, including the engagement of inhibitory receptors expressed in tumor-associated, exhausted CD8+ T cells, such as programmed cell death protein 1 (PD-1), T-cell immunoglobulin and mucin-domain containing-3 (TIM-3), lymphocyte-activation gene 3 (LAG-3), 2B4 (also known as CD244), and T cell immunoreceptor with Ig and ITIM domains (TIGIT).¹⁻² While immune checkpoint blockade therapies aimed at reinvigorating T cell effector function have demonstrated their clinical effectiveness,³⁻⁴ not all patients demonstrate long-term disease control.⁵ The refractory nature of terminally differentiated, exhausted CD8+ T cells to be reinvigorated by PD-1 blockade is one potential cause.⁶⁻⁸ This limitation warrants the need to explore modulatory pathways that potentially program T cells toward exhaustion.

Methods Single cell-RNA sequencing (scRNA-seq) data derived from the tumor-infiltrating lymphocytes (TILs) of melanoma patients⁹ were used for transcriptomic analysis and flow cytometry results were used to quantify protein levels in TILs. Murine B16-F10 (B16) melanoma model was used for both in vitro and in vivo studies. TCR-transgenic Pmel-1 and OT-1 transgenic mice, as well as CD47-/- (knockout, KO) mice were purchased from the Jackson Laboratory to generate CD47+/+ (wild-type, WT), CD47 $\pm$ (heterozygote, HET) mice with Pmel-1 or OT-1 background. For T cell co-transfer studies, Rag-deficient mice or C57BL/6j mice with sub-lethal irradiation (600cGy) were used as recipients. Naïve TCR-transgenic CD47-WT and CD47-HET CD8+ T cells were labelled, mixed in a 1:1 ratio for co-transfer experiments.

Results Flow cytometry analysis of human melanoma TILs found a strong upregulation of CD47 expression in tumor-associated, exhausted CD8+ T cells. We confirmed that CD47 transcription is significantly elevated among CD8+ T cells with a phenotype consistent with exhaustion using scRNA-seq results of TILs derived from melanoma patients.⁹ Our study in murine B16 melanoma model confirms our finding in melanoma patients. To specifically address the role of CD47 in anti-tumor CD8 effector function, we conducted T cell co-transfer studies and found that CD8+ T cells with lower copy number of CD47 (CD47-HET) significantly outnumber the co-transferred CD47-WT CD8+ T cells within the tumor, exhibiting an enhanced effector function and less exhausted phenotype. Our study demonstrates a potentially novel role for CD47 in mediating CD8+ T cell exhaustion.

Conclusions CD47 expression in CD8+ T cells programs T cells toward exhaustion.

Ethics Approval All mice were maintained in microisolator cages and treated in accordance with the NIH and American Association of Laboratory Animal Care regulations. All mouse procedures and experiments for this study were approved by the MSKCC Institutional Animal Care and Use Committee (IACUC).

REFERENCES

- Wherry EJ and M Kurachi. Molecular and cellular insights into T cell exhaustion. *Nat Rev Immunol* 2015;**15**(8): p. 486–99.