



Abstract 606 Figure 3 Correlation between IFN- γ and CXCL10 expression stratified by EGF expression. A. Cutaneous squamous cell carcinoma cell lines (GSE98767, n=44). B. Cutaneous squamous and basal cell carcinoma tumor samples (GSE125285, n=35)

role of EGF in modulating inflammation, and to understand this process in the pathogenesis of EGF receptor inhibitor-induced cutaneous toxicities and skin cancers.

Acknowledgements Emory Integrated Genomics Core

REFERENCES

- Annunziata MC, De Stefano A, Fabbrocini G, Leo S, Marchetti P, Romano MC and Romano I. 'Current Recommendations and novel strategies for the management of skin toxicities related to Anti-EGFR therapies in patients with metastatic colorectal cancer.' *Clin Drug Investig* 2019; **39**(9): 825–834.
- Hu JC, Sadeghi P, Pinter-Brown LC, Yashar S and Chiu MW. 'Cutaneous side effects of epidermal growth factor receptor inhibitors: clinical presentation, pathogenesis, and management.' *J Am Acad Dermatol* 2007; **56**(2): 317–326.
- Lacouture ME (2006). 'Mechanisms of cutaneous toxicities to EGFR inhibitors.' *Nat Rev Cancer* **6**(10):803–812.
- Liao Y, Wang J, Jaehnig EJ, Shi Z and Zhang B (2019). 'WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs.' *Nucleic Acids Res* **47**(W1): W199–W205.
- Lichtenberger BM, Gerber PA, Holcman M, Buhren BA, Amberg N, Smolle V, Schrupf H, Boelke E, Ansari P, Mackenzie C, Wollenberg A, Kislak A, Fischer JW, Rock K, Harder J, Schroder JM, Homey B and Sibilia M. 'Epidermal EGFR controls cutaneous host defense and prevents inflammation.' *Sci Transl Med* 2013; **5**(199):199ra111.
- Love MI, Huber W and Anders S. 'Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.' *Genome Biol* 2014; **15**(12):550.
- Ommori R, Park K, Miyagawa F, Azukizawa H, Kanno M and Asada H. 'Epidermal growth factor receptor (EGFR) inhibitory monoclonal antibodies and EGFR tyrosine kinase inhibitors have distinct effects on the keratinocyte innate immune response.' *Br J Dermatol* 2018; **178**(3): 796–797.
- Tan EH and Chan A. 'Evidence-based treatment options for the management of skin toxicities associated with epidermal growth factor receptor inhibitors.' *Ann Pharmacother* 2009; **43**(10): 1658–1666.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0607>

607 TJ210 (MOR210), A DIFFERENTIATED ANTI-C5AR ANTIBODY FOR ANTI-CANCER THERAPY

¹Jane Meng*, ¹Zhengyi Wang, ¹Wei Cao, ¹Chan Chen, ¹Joan Huaqiong Shen, ²Christian Augsburg, ²Julia Neugebauer, ²Stefan Haertle. ¹I-Mab BiPharma, Beijing, China; ²MorphoSys AG, Munich, Germany

Background Extensive investigations into the tumor microenvironment (TME) have uncovered molecular mechanisms linking aberrant complement activation and cancer progression. Specifically, C5a, as a highly potent chemoattractant, recruits immune suppressive myeloid derived suppressive cells (MDSCs), neutrophils and M2 macrophages into the tumor site and accelerates tumor progression. Blockade of C5a/C5aR (CD88) pathway has been identified as a promising target to control MDSCs and restore tumor-killing ability of T and NK

cells. TJ210, in licensed from MorphoSys as MOR210, is a differentiated anti-C5aR monoclonal antibody with a unique binding epitope.

Methods Interaction of TJ210 with C5aR was assessed through binding of the recombinant antigen, Flp-In CHO cells expressing C5aR and primary neutrophils. In vitro blockade of C5a/C5aR pathway was tested by inhibition of CD11b upregulation on granulocytes and monocytes induced by C5a, as well as neutrophil migration towards C5a. The in vitro synergistic effect of TJ210 with anti-PD-1 antibody was assessed in a T cell and differentiated MDSC co-culture system. The in vivo anti-tumor effect was tested in the MC38 syngeneic mouse model, in which mice were treated with a TJ210 mouse surrogate antibody either alone or in combination with an anti-PD-1 antibody.

Results TJ210 bound to C5aR with high affinity and did not cross-react with other GPCR members including C5L2, ChemR23, FPR1 and C3aR. Unlike the reference antibody, TJ210 specifically interacted with the N-terminus of C5aR but not extracellular loops. TJ210 effectively inhibited CD11b upregulation on granulocytes and monocytes as well as neutrophil migration mediated by C5a. When compared with the reference antibody, TJ210 maintained potent antagonism at high ligand concentrations and over longer duration, properties that might translate into beneficial in vivo effects at pathophysiological conditions. In the in vitro co-culture system, presence of TJ210 and anti-PD-1 antibody enhanced IFN- γ release compared to either single agent, indicating a synergistic effect on T cells. In the in vivo syngeneic mouse model, combination treatment effectively inhibited tumor growth. Immune cell population analysis revealed significant elevation of CD8+ T cells and M1 macrophages compared to mono-treatment.

Conclusions This series of in vitro and in vivo data demonstrate that TJ210 is a differentiated anti-C5aR antibody with unique binding epitope exhibiting superior anti-tumor potential especially in combination with an anti-PD-1 antibody. These data support further clinical studies of TJ210 in patients with solid tumors.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0607>

608 IMMUNODOMINANT LISTERIA EPITOPES COMPETE WITH VACCINE-DIRECTED CD8+ T-CELL RESPONSES RESCUED BY PEPTIDE-MHC STABILIZING MODIFICATIONS

John Flickinger*, Jagmohan Singh, Yanki Yarman, Robert Carlson, Scott Waldman, Adam Snook. *Thomas Jefferson University, Philadelphia, PA, USA*

Background The Gram-positive bacterium *Listeria monocytogenes* (Lm) is a promising vector for cancer immunotherapy due to its ability to directly infect antigen-presenting cells, induce potent CD8+ T-cell immunity, and remodel immunosuppressive tumor microenvironments.¹ Recent clinical trials have demonstrated safety and immunogenicity of Lm-based cancer vaccines in lung, cervical, pancreatic, and other cancers. In colorectal cancer, the transmembrane receptor guanylyl cyclase C (GUCY2C) is an emerging target for immunotherapy.² Here, we examined the immunogenicity of a recombinant strain of *Listeria monocytogenes* secreting GUCY2C (Lm-GUCY2C). Surprisingly, Lm-GUCY2C vaccination induced