REFERENCES

1. Gentles AJ, Newman AM, Liu CL, *et al.* The prognostic landscape of genes and infiltrating immune cells across human cancers. *Nature Medicine* 2015;**21**(8):938–945.
2. Tosolini M, Pont F, Poupot M, *et al.* Assessment of tumor-infiltrating TCRV γ 9V δ 2 $\gamma\delta$ lymphocyte abundance by deconvolution of human cancers microarrays. *Oncoimmunology*. 2017;**6**(3):e1284723.
3. Harly C, Guillaume Y, Nedellec S, *et al.* Key implication of CD277/butyrophilin-3 (BTN3A) in cellular stress sensing by a major human $\gamma\delta$ T-cell subset. *Blood* 2012;**120**(11):2269–2279.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0687>

688

IN VIVO EXPANSION OF GAMMA DELTA T CELLS BY A CD19-TARGETED BUTYROPHILIN HETERODIMER LEADS TO ELIMINATION OF PERIPHERAL B CELLS

Suresh De Silva*, George Fromm, Louis Gonzalez, Arpita Patel, Kyung Yoon, Zachery Opheim, Robert Farmer, Dean Chamberlain, Taylor Schreiber. *Shattuck Labs, Durham, NC, USA*

Background A primary mechanism of cancer immunotherapy resistance involves downregulation of specific antigens or major histocompatibility complex based antigen presentation, which renders tumor cells invisible to alpha-beta T cells, but not gamma-delta T cells. Recently, a two-step model of gamma-delta T cell activation has emerged, wherein one butyrophilin (BTN, ie. BTN2A1) directly binds the gamma-delta TCR but is only activated if certain molecular patterns (eg. phosphoantigens) facilitate recruitment of a second BTN (ie. BTN3A1) into a complex to form a BTN2A1/3A1 heterodimer. The BTN2A1/3A1 complex specifically activates the predominant gamma-delta T cell population in the peripheral blood, comprising the Vg9d2 T cell receptor (TCR), but does not activate the primary gamma-delta T cell population in mucosal tissues, comprising the Vg4 TCR. The unique mechanism of action and specificity of gamma-delta TCR/BTN interactions suggests that therapeutic proteins comprising specific BTN heterodimers could be used to target specific gamma-delta T cell populations, with a lower risk of off-target activation common with CD3-directed T cell engagers.

Methods Human BTN2A1/3A1-Fc-CD19scFv and mouse BTNL1/6-Fc-CD19scFv heterodimeric fusion proteins were purified and binding to CD19 or the respective gamma-delta TCRs was assessed by ELISA, Octet and flow cytometry using gd T-cells isolated from human peripheral blood and mouse intestinal tissue. The functionality of the constructs to activate gamma-delta T cells and mediate killing of tumor cells was assessed using live cell imaging in vitro as well as a murine B-cell lymphoma model in vivo.

Results The CD19-targeting scFv domains of the BTN heterodimer fusion proteins bound to human and mouse CD19 with low nanomolar affinity. The BTN2A1/3A1-Fc-CD19scFv compound specifically bound to the Vg9d2 TCR on human gd T cells while the mouse BTNL1/6-Fc-CD19scFv bound to Vg7d4 TCR on mouse gd T cells. Both compounds were able to activate gd T cells in a co-culture assay resulting in degranulation and increased surface expression of CD107a and also increased apoptosis of CD19+ tumor cells. Intraperitoneal administration of the mouse BTNL1/6-Fc-CD19scFv led to anti-tumor effects in A20 tumor bearing BALB/c mice. Phenotyping from BTNL1/6-Fc-CD19scFv treated mice revealed profound and rapid expansion of the endogenous gamma-delta T cells in the circulation and tumor, with concomitant depletion of peripheral CD19+ B-cells, confirming the mechanism of

action of the heterodimer as a gamma-delta T cell specific engager.

Conclusions These results provide proof of mechanism for in vivo manipulation of gamma-delta T cells using antigen-targeted butyrophilin heterodimeric fusion proteins for the treatment of cancer.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0688>

689

ATRC-101 DRIVES POTENT SINGLE-AGENT ACTIVITY IN MOUSE SYNGENEIC TUMOR MODELS VIA A NOVEL CELLULAR MECHANISM OF ACTION

Yvonne Leung, Nikhil Vad, Anne Ye, Daniel Santos, Wei Cao, Cathrin Czupalla, John Vivian, Carlene Williams, Judevin Sapugay, Michael Harbell, Shaun Lippow, Chantia Carroll, Lance Kates, Benjamin Haugen, Gary Bolton, Mark Armanini, Iraz Aydin, Mark Whidden, Mauricio Velasco-Delgado, Felix Chu, Erin Wechsler, Ngan Nguyen, William Robinson, Tito Serafini, Daniel Emerling, Norman Greenberg, Amy Manning-Bog*, Alexander Scholz. *Atreca, Inc, South San Francisco, CA, USA*

Background We have previously demonstrated adaptive antibody responses targeting public tumor antigens in cancer patients. ATRC-101, a clinical stage, engineered version of an antibody identified in such a patient, displays robust single-agent activity in syngeneic tumor models requiring Fc receptors (FcRs) expressed by innate immune cells and the presence of CD8+ T cells. The novel target of ATRC-101 was found to be a tumor-restricted ribonucleoprotein (RNP) complex, and because RNP complexes drive T cell responses in infectious and autoimmune disease via innate immune cells, we further characterized the mechanism-of-action of ATRC-101. Here we describe changes in immune cell populations in a tumor model proximal to treatment initiation with ATRC-101. **Methods** Mice bearing EMT6 tumors received ATRC-101 beginning on day 7 post-tumor inoculation. Tissues were harvested between days 7 and 14 and analyzed by flow cytometry and immunohistochemistry. Transcriptome analysis was performed using RNA sequencing on whole tumors taken on days 7, 9, and 12.

Results The earliest significant changes induced by ATRC-101, relative to vehicle, were noted just 24 hours after dosing: increased numbers of cDC1 cells in blood, and decreased numbers of cDC2 cells in blood and M-MDSCs in tumor. A significant increase of CD8+ T cells was observed in blood 48 hours after dosing and in tumor 96 hours after dosing. Increased numbers of NK cells were also observed in blood and tumor at this later time. Multiplex analysis of circulating cytokines demonstrated a very early increase in myeloid chemo-attractants, such as MCP1 and MIP1a. Whole exome sequencing of tumor samples showed that ATRC-101 dosing drives a significant increase, relative to vehicle, in the expression of interferon-stimulated genes. Co-culturing experiments demonstrated that induced, bone marrow-derived dendritic cells are activated by ATRC-101 and its target in a dose-dependent fashion.

Conclusions Dosing with ATRC-101 in the EMT6 syngeneic tumor model, in which ATRC-101 displays notable single-agent activity, leads to changes in immune cell composition in the blood and tumor, with the earliest changes observed in myeloid or myeloid-derived cell populations, and to the early appearance of myeloid chemo-attractants. We believe these data indicate that ATRC-101 acts proximally on the myeloid cell populations in the tumor, leading to a remodeling of the tumor environment and an adaptive immune response that

Abstracts

includes CD8⁺ T cells driving tumor regression. Our data demonstrate that ATRC-101, bound to its target which is an RNP complex, can activate myeloid cells and are consistent with this activation occurring via FcR and Toll-like receptor (TLR) pathways.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0689>

690

CD122-SELECTIVE IL-2 COMPLEXES TREAT OVARIAN CARCINOMAS, INDUCE TREG FRAGILITY AND PROMOTE T CELL STEM CELLS

¹Yilun Deng*, ²Justin Drerup, ¹Xinyue Zhang, ¹Ryan Reyes, ¹Jenny Mendez, ¹Aravind Kancharla, ¹Myrna Garcia, ¹Haiyan Bai, ¹Alvaro Padron, ¹Harshita Gupta, ¹Tyler Curiel. ¹UT Health San Antonio, San Antonio, TX, USA; ²UT Southwestern, Dallas, TX, USA

Background Ovarian cancer (OC) responds poorly to immunotherapies. Regulatory T cells (Treg) engage IL-2 by high-affinity CD25 for differentiation and function,¹ and anti-tumor effector T cells (Teff) use intermediate affinity CD122. We studied IL-2 complexes (IL-2c) that selectively activate CD122 (Teff) over CD25 (Tregs).

Methods Orthotopic ID8agg-luc mouse OC burden was measured by in vivo imaging. Tumor, ascites and draining lymph nodes (TDLN) were analyzed by flow and tSNE. IL-2c was complexed using 1.5 µg/mouse IL-2 and 7.5 µg/mouse aIL-2 (clone JES6-5H4) before i.p. injection every other day x 4 starting at day 7. antiPD-L1 was given at 100ug/mouse every 3 days x 4 starting from Day 11. FIR mice² were used to sort live Tregs.

Results IL-2c but not antiPD-L1 potently inhibits ID8agg (figure 1). IL-2c decreased ascites Treg functional markers (e.g., CD25, granzymeB) while upregulating the same markers on Teffs (figure 2). IL-2c inhibited Treg suppression in ascites while TDLN Tregs were unaffected (figure 3). tSNE showed great similarity of TDLN Tregs treated with isotype and IL-2c while ascites Tregs after IL-2c showed a fragile phenotype (e.g., increased PD-1, T-bet, and IFNγ with maintained FoxP3 expression [figure 4]) which is known to contribute to better response to cancer immunotherapy.³⁻⁴ We observed a complete reduction of tumor bioluminescence with IL-2c and antiPD-L1 combo treatment in nearly all subjects significantly exceeding effect of IL-2c alone (figure 5). A CD8⁺CXCR5⁺

Abstract 690 Figure 1 IL-2c but not aPD-L1 treats ID8agg Luciferase signal of ID8agg-luc tumors treated with isotype, αPD-L1, or IL-2c measured by in vivo imaging. Arrows indicate treatments.

Abstract 690 Figure 2 IL-2c inhibits functional markers on tregs and promotes teff

Expression of CD25 and granzymeB were measured by flow cytometry in indicated population from isotype or IL-2c treated ascites 3 weeks after final IL-2c dose.

Abstract 690 Figure 3 IL-2c reduces ascites Treg suppressive function

+TCF-1⁺ T cell stem cell (TCSC) population reportedly improves immune checkpoint blockade efficacy.⁵⁻⁶ Since CD122 is regulated by TCF-1,⁷ we explored the effect of IL-2c on these TCSC. IL-2c significantly induced a CD8⁺TCF-1⁺ + TCSC population in ID8agg tumors (figure 6), possibly through a positive feedback loop by further enhancing CD122 expression on TCF-1⁺, but not TCF-1⁻ cells (figure 7). tSNE analysis of detailed immune phenotype of IL-2c induced TCSC revealed that these TCSC differed from those induced by antiPD-L1. In ID8agg, antiPD-L1-induced TCSC are mostly CXCR5⁺ and PD1⁺, consistent with previous reports in other cancers³⁻⁴ while IL-2c-induced TCSC were PD1⁻ (figure 8), expressed CCR2 and CXCR3, and produced TNFα (figure 9).

Abstract 690 Figure 4 IL-2c induces Treg fragility in ascites but not TDLN tSNE analysis on CD45⁺CD3⁺CD4⁺FoxP3⁺ cells from ascites and TDLN of isotype and IL-2c treated ID8agg-luc challenged mice. Right, representative bar graphs of flow data.