patients ($p = 0.04$). Kaplan-Meier survival analysis showed that higher density of CD163+ and CD79a+ cells were independently associated with shorter recurrence free survival (RFS). Notably, these differences in RFS remained in BCG immunotherapy-naïve patients ($n=170$).

Conclusions These findings are the first evidence of sexual dimorphism in the TIME of NMIBC and may help to partially explain the worse clinical outcomes experienced by female patients. This study also provides the first evidence of the negative prognostic impact of B cells in NMIBC. Overall, this study provides insight into more rational implementation of immune-based therapies in female NMIBC patients.

Ethics Approval This study was approved by the Ethics Review Board at Queen's University, Kingston, ON, Canada.

REFERENCES

1. Saginala K, Barsouk A, Aluru JS, Rawla P, Padala SA, Barsouk A. Epidemiology of bladder cancer. *Med Sci* 2020;**8**.
2. Uhlig A, Strauss A, Seif Amir Hosseini A, Lotz J, Trojan L, Schmid M, *et al*. Gender-specific differences in recurrence of non-muscle-invasive bladder cancer: a systematic review and meta-analysis. *Eur Urol Focus* 2018;**4**:924–36.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0870>

871

IN VITRO IMMUNOREGULATORY EFFECT FROM CERVICAL CANCER DERIVED MESENCHYMAL STROMAL CELLS OVER MOLECULES EXPRESSION IN MONOCYTE DERIVED MACROPHAGE POLARIZATION

¹Víctor Cortés-Morales*, ²Juan Montesinos, ²Luis Chávez-Sánchez, ²Sandra Espíndola-Garibay, ²Alberto Monroy-García, ²Teresa Apresa-García, ²Héctor Mayani. ¹Universidad Nacional Autónoma de México, Mexico City, Mexico; ²Instituto Mexicano del Seguro Social, Ciudad de México, Mexico

Background Macrophages are immunological cells that sense microenvironmental signals that may result in the polarized expression of either proinflammatory (M1) or anti-inflammatory (M2) phenotype.¹ Macrophages M2 are present in tumoral microenvironment and their presence in patients with cervical cancer (CeCa) is related with less survival.² Mesenchymal Stromal Cells (MSCs) are also present in tumor microenvironment of cervical cancer (CeCa-MSC), which have shown immunoregulatory effects over CD8 T cells, decreasing their cytotoxic effect against tumoral cells.³ Interestingly, MSCs from bone marrow (BM-MSC) decrease M1 and increase M2 macrophage polarization in an in vitro coculture system.⁴ Macrophages and MSCs are present in microenvironment of cervical cancer, however it is unknown if MSCs play a role in macrophage polarization. In the present study, we have evaluated the immunoregulatory capacity of CeCa-MSCs to induce macrophage polarization.

Methods CD14 monocytes were isolated from peripheral blood and cultivated in the absence or presence of MSCs from BM, normal cervix (NCx) and CeCa. Two culture conditions were included, in the presence of induction medium to favors M1 (GM-CSF, LPS and IFN γ) or M2 (M-CSF, IL-4 and IL-13) macrophage polarization. M1 (HLA-DR, CD80, CD86 and IFN γ) or M2 (CD14, CD163, CD206, IDO and IL-10) macrophage molecular markers were evaluated by flow cytometry. Finally, we evaluated concentration of IL-10 and TNF α in conditioned medium from all coculture conditions.

Results We observed that CeCa-MSCs and BM-MSCs in presence of M1 induction medium, decreased M1 macrophage

markers (HLA-II, CD80, CD86 and IFN γ), and increase the expression of CD14 (M2 macrophage marker). Interestingly, in presence of M2 induction medium, BM-MSCs and CeCa-MSCs but not CxN-MSC increased CD163, CD206, IDO and IL-10 (M2 macrophage markers). We observed a decreased concentration of TNF α in the supernatant medium from all cocultures with MSCs, but only in presence of CeCa-MSCs, increased IL-10 concentration was detected in such cocultures.

Conclusions In contrast to NCx-MSCs, CeCa-MSCs similarly to BM-MSCs have in vitro capacity to decrease M1 and increase M2 macrophage phenotype.

Acknowledgements Acknowledgments The authors are indebted to gratefully acknowledge to CONACYT (Grant No. 272793) and IMSS (Grant no. 1731) for support to Juan J. Montesinos research.

REFERENCES

1. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep* 2014;**6**:13.
2. Petrillo M, Zannoni GF, Martinelli E, *et al*. Polarization of tumor-associated macrophages toward M2 phenotype correlates with poor response to chemoradiation and reduced survival in patients with locally advanced cervical cancer. *PLoS One* 2015;**10**: e0136654.
3. Montesinos JJ, Mora-García Mde L, *et al*. In vitro evidence of the presence of mesenchymal stromal cells in cervical cancer and their role in protecting cancer cells from cytotoxic T cell activity. *Stem Cells Dev* 2013;**22**:2508-2519.
4. Vasandan AB, Jahnvi S, Shashank C. Human mesenchymal stem cells program macrophage plasticity by altering their metabolic status via a PGE 2-dependent mechanism. *Sci Rep* 2016;**6**:38308.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0871>

872

NEOADJUVANT CHEMORADIOTHERAPY ENHANCES T CELL INFILTRATION IN PANCREATIC DUCTAL ADENOCARCINOMA BUT HIGH PERCENTAGE OF REGULATORY T CELLS ASSOCIATES WITH POOR SURVIVAL

¹Benjamin Fullerton*, ²Robyn Gartrell, ³ThomasENZler, ⁴Pan Kim, ⁵Ladan Fazlollahi, ⁶Andrew Chen, ⁷Subha Perni, ⁸Stuart Weisberg, ⁹Emanuelle Rizk, ⁸Eun Jeong Oh, ²Xinzheng Guo, ²Codruta Chiuzan, ²Raul Rabadán, ²Donna Farber, ⁵Helen Remotti, ⁵David Horowitz, ⁵Yvonne Saenger. ¹Columbia University Medical Center, New York, NY, USA; ²Columbia University Irving Medical Center, Edgewater, NJ, USA; ³University of Michigan Medicine, Ann Arbor, MI, USA; ⁴University of Pittsburgh Medical Center, Pittsburgh, USA; ⁵Columbia University Irving Medical Center/New York Presbyterian, New York, NY, USA; ⁶Vagelos College of Physicians and Surgeons, New York, NY, USA; ⁷Massachusetts General Hospital and Brigham and Women's Hospital/Dana-Farber Cancer Institute, Boston, USA; ⁸Columbia University, Mailman School of Public Health, New York, USA

Background Currently, diagnosis with pancreatic ductal adenocarcinoma (PDAC) renders an almost intrinsically poor patient prognosis. Despite complete surgical resection and intense neoadjuvant and/or adjuvant treatment the great majority of patients will ultimately relapse and die from the disease. Further, PDAC has been characterized as highly immune resistant. It is speculated that radiation, chemotherapy, or chemoradiation cause the release of tumor antigens and inflammatory cytokines eventually leading to increased immunogenicity of PDAC.

Methods We used computational quantitative multiplex immune fluorescence (qmIF) ($n=31$) and the NanoString assay ($n=34$) to quantitatively analyze the effect of neoadjuvant