expansion. We found that shortening ex vivo expansion of either TCR-specific murine Th17 cells or human CAR Th17 cells licenses the cell product to eradicate large tumors in low doses and generate long-lived memory against tumor.¹ Therapeutic Th17 cells induce the systemic release of IL-6, IL-17, GM-CSF, and MCP-1 among other cytokines in tumor-bearing hosts, reminiscent of clinical cytokine release syndrome. As the toxicity of cytokine release is managed in patients through IL-6 blockade, we addressed the impact of IL-6 on efficacy and durability of Th17 cell therapy. We hypothesized that IL-6, induced by Th17 cells, was fueling the durable memory properties of this cell product.

Methods Th17 cells were expanded ex vivo using the TRP-1 transgenic mouse model in which CD4+ T cells express a TCR that recognizes tyrosinase-related protein 1 on melanoma. Naïve CD4+ T cells were polarized to the Th17 phenotype and infused into mice with B16F10 melanoma after a nonmyeloablative total body irradiation (5 Gy) preparative regimen.

Results IL-6 blockade, targeting either IL-6R or neutralization of the cytokine, did not significantly impact the primary immune response of adoptively transferred Th17 cells against tumor. However, administering IL-6 blockade acutely after Th17 transfer resulted in a higher incidence of tumor relapse upon secondary tumor challenge, thereby compromising long-lived antitumor immunity.¹ Mounting a secondary response to tumor was dependent on CD4+ T cells, but not CD8+ T cells, persisting in the host. Mechanistically, IL-6 blockade reduced pSTAT3 and Bcl2 in transferred T cells but did not greatly impact the concentration of other systemic cytokines. As a small fraction of Tregs remain in the Th17 cell product ex vivo, we examined the engraftment of those Tregs after transfer. IL-6 was critical to suppress engraftment of FoxP3+ donor T cells from the CD4+ T cell product. Thus, IL-6 promoted robust tumor infiltration by donor effector over regulatory cells for early Th17 cells relative to cell products expanded longer durations ex vivo.¹

Conclusions Overall, short-term expanded Th17 cells uniquely induced IL-6 unlike Th17 cells expanded longer ex vivo. IL-6 promoted Th17 survival, reduced engraftment of tumor-specific Tregs, and was critical to durable memory. This work may suggest that the universal strategy to inhibit IL-6 during cytokine release syndrome may come at the expense of long-term efficacy for specific cell therapy approaches.

REFERENCE

1. Knochelmann HM, Dwyer CJ, Smith AS, Bowers JS, Wyatt MM, Nelson MH, Rangel Rivera GO, Horton JD, Krieg C, Armeson K, Lesinski GB, Rubinstein MP, Li Z, Paulos CM. IL-6 fuels durable memory for Th17 cell-mediated responses to tumors. *Cancer Res.* 2020. Epub ahead of print. PMID: 32561531.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0119>

120

P-MUC1C-ALLO1: AN ALLOGENEIC CAR-T FOR MULTIPLE SOLID TUMOR INDICATIONS

Anna Kozłowska*, Yan Zhang, Jacqueline Fritz, Steven Wang, Rebecca Codde, Elvira Argus, Samad Ibitokou, Vanitra Richardson, Sumiti Jain, Maximilian Richter, Deepak Patil, Yening Tan, Min Tong, Lu Yao, Majid Ghoddsu, Eric Ostertag, Julia Coronella, Devon Shedlock. *Poseida, San Diego, CA, USA*

Background MUC1 is a highly glycosylated protein that is expressed at the apical border of mucosal epithelium where it plays a protective role. MUC1 is comprised of an N-terminal subunit (MUC1N) tethered to a C-terminal subunit (MUC1C),

forming a stable complex on the cell surface. A proteolytic ‘stump’ of MUC1C that may be aberrantly glycosylated is over-represented in cancer, making it an attractive therapeutic target. Here we report generation of allogeneic MUC1C-specific CAR T cells, P-MUC1C-ALLO1, that are designed to leverage the learnings of our P-BCMA-ALLO1 program. P-MUC1C-ALLO1 targets a MUC1C epitope and has the potential for efficacy against a wide range of solid tumors, without targeting normal epithelial cells.

Methods mRNA-generated MUC1C CAR-T cells were evaluated for specificity and function by degranulation assay against various solid tumor and normal cells and cell lines. Autologous and allogeneic MUC1C CAR-T cells were produced using the piggyBac® DNA Modification System, a non-viral CAR-T manufacturing method that produces CAR-T products with an exceptionally high percentage of T stem cell memory (T_{SCM}) cells. To produce allogeneic cells, multiplex editing of both TRBC and B2M was performed with the Cas-CLOVER™ Site-Specific Gene Editing System to reduce or eliminate GvHD and host versus graft alloreactivity, respectively. To determine in vivo antitumor efficacy of MUC1C CAR-T cells, we employed the MDA.MB.468 triple negative breast cancer model and the OVCAR3 disseminated ovarian cancer model.

Results Specific degranulation of transiently-expressing CAR+ T cells was observed against multiple tumor cells, with no observable activity against normal human primary cells. In assay of stable P-MUC1C-101 CAR-T cells, more than 95% expressed CAR, and were comprised of an exceptionally high percentage of T_{SCM} cells (CD45RA+CD62L+CD45RO-). In vitro, P-MUC1C-ALLO1 cells specifically proliferated, lysed, and secreted IFN-γ against MUC1C+ breast and ovarian tumor cell lines. In breast cancer in vivo xenograft model, both unedited (MUC1C CAR-T) and edited (P-MUC1C-ALLO1) MUC1C CAR-T eliminated established, triple negative MDA.MB.468 tumor cells to undetectable levels, demonstrating the efficacy of the MUC1C CAR-T and the robustness of the allogeneic platform. In the OVCAR3 xenograft model, intraperitoneally administered MUC1C CAR-T eliminated established tumor cells to levels below the limit of detection.

Conclusions P-MUC1C-ALLO1 is Poseida’s allogeneic CAR T_{SCM} product that has a potential to treat multiple MUC1C-expressing indications. P-MUC1C-ALLO1 displayed in vitro specificity for tumor vs normal cells, and in vivo efficacy against xenograft models of breast and ovarian cancer. We anticipate an IND filing and initiation of a Phase 1 clinical trial in 2021.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0120>

122

GUANYLYL CYCLASE C AS A TARGET FOR CAR-T CELL THERAPY IN A METASTATIC GASTRIC CANCER MODEL

Amanda Lisby*, Trevor Baybutt, Megan Weindorfer, Robert Carlson, Alicja Zalewski, Scott Waldman, Adam Snook. *Thomas Jefferson University, Philadelphia, PA, USA*

Background Gastric cancer is the sixth most common cancer and second-leading cause of cancer-related mortality worldwide.¹ The heterogenous and genetically complex nature of this disease underlies the challenges in developing effective therapies for metastatic gastric cancer. In the majority of cases, stomach tumors evolve from intestinal metaplasia resulting in ectopic expression of the enterocyte differentiation antigen guanylyl cyclase C (GUCY2C) by ~50% of primary and

metastatic gastric cancers.^{2–4} In the context of the efficacy of GUCY2C-directed chimeric antigen receptor (CAR)-T cells against metastatic colorectal cancer in animal models,^{5,6} we hypothesized that this adoptive cell therapy may be effective against metastatic gastric cancer.

Methods Here, we explored the efficacy of GUCY2C-directed CAR-T cells for gastric cancer in a patient derived xenograft (PDX) tumor model. Also, we interrogated translational GUCY2C biomarker assays using RT-qPCR, immunoblot analysis, and immunohistochemistry (IHC) for the intended purpose of identifying patients whose tumors express GUCY2C and could benefit from GUCY2C-directed CAR-T cell therapy.

Results GUCY2C-directed CAR-T cells significantly reduced subcutaneous T84 colorectal tumor growth, producing a 5-fold reduction in tumor volume, compared to control treated tumors. GUCY2C-directed CAR-T cells produced no response in tumors produced from the GUCY2C-deficient colorectal cancer cell line, SW480. Importantly, GUCY2C-directed CAR-T cells controlled gastric cancer PDX growth, maintaining a >12-fold reduction in tumor volume compared to control and in some cases produced complete tumor elimination. Furthermore, IHC based assays, indicate that antibodies developed in our laboratory may be suitable for development of a companion diagnostic for GUCY2C-directed CAR-T cells. Indeed, the commercial polyclonal antibody demonstrated robust, non-specific staining regardless of tissue type or GUCY2C mRNA profile, while novel monoclonal antibodies produced in our laboratory primarily detected protein localized to the membrane of glandular epithelial cells, demonstrating antigen specificity, and indicating their potential for further development in diagnostic companion assays to identify gastric cancer patients who may benefit from GUCY2C-directed CAR-T cell therapy.

Conclusions GUCY2C-directed CAR-T cells prevented the growth of, and at times eliminated, a subcutaneous gastric cancer PDX model. In the context of previously established safety in mouse models, additional studies defining the efficacy of GUCY2C-directed CAR-T cells in gastric cancer models may allow future translation of this therapy to patients with advanced gastric cancers. Concurrent development of a novel companion diagnostic IHC assay would permit identification of the ~50% of gastric cancer patients whose tumors express GUCY2C and could benefit from this therapy.

Acknowledgements This work was supported by a DeGregorio Family Foundation Award and by the Department of Defense Congressionally Directed Medical Research Programs (W81XWH-17-1-0299, W81XWH-19-1-0263, and W81XWH-19-1-0067) to AES. SAW is supported by the National Institutes of Health (NIH) (R01 CA204881, R01 CA206026, and P30 CA56036), the Defense Congressionally Directed Medical Research Program W81XWH-17-PRCRP-TTSA, and Targeted Diagnostic & Therapeutics. SAW and AES were also supported by a grant from The Courtney Ann Diacont Memorial Foundation. SAW is the Samuel M.V. Hamilton Professor of Thomas Jefferson University. AZ was supported by NIH institutional award T32 GM008562 for Postdoctoral Training in Clinical Pharmacology. The authors thank the NCI Patient-Derived Models Repository for their support and resources to make this research possible. The authors also thank the Sidney Kimmel Cancer Center Translational Research & Pathology Core Facility, and the Office of Animal Resources at Thomas Jefferson University for their continued technical assistance and support in this research.

Ethics Approval This study was approved by the Thomas Jefferson University Institutional Review Board (#14.0204) and the Institutional Animal Care and Use Committee (Protocol #01529).

REFERENCES

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424. doi:10.3322/caac.21492.
- Park J, Schulz S, Haaf J, Kairys JC, Waldman SA. Ectopic expression of guanylyl cyclase C in adenocarcinomas of the esophagus and stomach. *Cancer Epidemiol Biomarkers Prev* 2002;11:739–44.
- Birbe R, Palazzo JP, Walters R, Weinberg D, Schulz S, Waldman SA. Guanylyl cyclase C is a marker of intestinal metaplasia, dysplasia, and adenocarcinoma of the gastrointestinal tract. *Hum Pathol*. 2005;36:170–9. doi:10.1016/j.humpath.2004.12.002.
- Mathur D, Root AR, Bugaj-Gaweda B, Bisulco S, Tan X, Fang W, et al. A Novel GUCY2C-CD3 T-Cell Engaging Bispecific Construct (PF-07062119) for the Treatment of Gastrointestinal Cancers. *Clin Cancer Res* 2020;26:2188–202. doi:10.1158/1078-0432.CCR-19-3275.
- Magee MS, Kraft CL, Abraham TS, Baybutt TR, Marszalowicz GP, Li P, et al. GUCY2C-directed CAR-T cells oppose colorectal cancer metastases without autoimmunity. *Oncoimmunology* 2016;5:e1227897. doi:10.1080/2162402X.2016.1227897.
- Magee MS, Abraham TS, Baybutt TR, Flickinger JC, Ridge NA, Marszalowicz GP, et al. Human GUCY2C-Targeted Chimeric Antigen Receptor (CAR)-Expressing T Cells Eliminate Colorectal Cancer Metastases. *Cancer Immunol Res* 2018;6:509–16. doi:10.1158/2326-6066.CIR-16-0362.

<http://dx.doi.org/10.1136/jitc-2020-SITC2020.0122>

123

NATURAL KILLER CELLS ENGINEERED WITH AN INDUCIBLE, RESPONSIVE GENETIC CONSTRUCT TARGETING TIGIT AND CD73 TO RELIEVE IMMUNOSUPPRESSION WITHIN THE GBM MICROENVIRONMENT

Kyle Lupo *, Sandro Matosevic. *Purdue University, Lafayette, USA*

Background Solid tumors such as GBM are particularly difficult to treat, being largely resistant to traditional treatments, fueling interest in alternative treatment approaches, including cell-based immunotherapy. Natural killer (NK) cells have emerged as promising effectors to target GBM through genetic modifications and ex vivo manipulation. However, immunosuppressive conditions within the tumor microenvironment (TME) further complicate NK cell-based treatments. Specifically, within the TME tumor cells release of high levels of ATP extracellularly. While intracellular ATP is necessary for cell metabolism, extracellular ATP is converted into adenosine (ADO) by ectonucleotidases CD39 and CD73, both overexpressed on GBM.¹ Extracellular ADO induces immunometabolic suppression of NK cells through binding with A2A adenosine receptors (A2ARs) on NK cells, suppressing cytokine secretion, proliferation, and other functional activities.^{2–4} Adding to the suppression of NK cells is the interaction between CD155, expressed highly on GBM and other solid tumors, and T cell immunoreceptor with Ig and ITIM domains (TIGIT) expressed on NK cells. This interaction signals inhibition of NK cell cytolytic function, allowing for cancer cell immune-evasion.^{5,6}

Methods To restore impaired NK cell anti-tumor activity, we have engineered NK cells to concomitantly target CD155 and CD73-induced immunosuppression on GBM using a tumor-responsive genetic construct. The construct is capable of blocking the immunosuppressive CD155/TIGIT interaction,