cancer vaccination opportunity when combined in heterologous prime-boost regimen.

Materials and Methods Mice were vaccinated with subcutaneous (s.c.) injection of KISIMA-TAA vaccine and/or with intravenous injection of VSV-GP-TAA in different settings. Immunogenicity was assessed by measuring the peripheral antigen-specific response. Anti-tumoral efficacy as well as in depth monitoring of TILs and tumor microenvironment modulation were assessed following therapeutic vaccination in different tumor models. Additionally, transcriptome and immunohistochemistry analyses of the TC-1 tumor have been performed. Combination of heterologous prime-boost with checkpoint blockade PD-1 therapy has been assessed.

Results Priming with KISIMA-TAA followed by VSV-GP-TAA boost induced a large pool of polyfunctional and persistent antigen-specific cytotoxic T cells in the periphery as well as within the tumor in several tumor models. Frequencies of antigen specific T cells are significantly higher than the respective homologous vaccinations. Additionally, transcriptome analysis of a cold tumor model revealed profound changes in the tumor microenvironment upon heterologous vaccination, including a strong upregulation of gene signatures of several pro-inflammatory cytokines and chemokines required for anti-tumor immunity along with dendritic and T cell trafficking and activation. This was corroborated by flow-cytometric analysis of tumor-infiltrating leukocytes showing massive CD8⁺ and CD4⁺ T cell infiltration as well as repolarization of M2-like macrophages towards M1-phenotype. The presence of the CD8⁺ T cells within the tumor core was confirmed by immunohistochemistry analysis. Moreover, combining heterologous vaccination with checkpoint blockade further improved its therapeutic efficacy and the number of long-term survivors.

Conclusions The KISIMA/VSV-GP heterologous prime-boost approach holds great promise for patients with primary or acquired resistance to checkpoint blockade due to its ability to induce tumor-specific T cell, improve T cell infiltration and increase tumor inflammation, even in tumors with limited permissivity for the oncolytic virus.

Disclosure Information E. Belnoue: A. Employment (full or part-time); Significant; AMAL Therapeutics SA. K. Das: None. M. Rossi: A. Employment (full or part-time); Significant; AMAL Therapeutics SA. T. Hofer: None. S. Danklmaier: None. T. Nolden: A. Employment (full or part-time); Significant; Viratherapeutics GmbH. L. Schreiber: None. K. Angerer: None. J. Kimpel: None. S. Hoegler: None. L. Kenner: None. D. von Laer: None. K. Elbers: A. Employment (full or part-time); Significant; Viratherapeutics GmbH. G. Wollmann: None. M. Derouazi: A. Employment (full or part-time); Significant; AMAL Therapeutics SA.

P03.03

NOVEL MACHINE-LEARNING TOOLS IMPROVE COST-EFFECTIVE DEVELOPMENT OF PERSONALISED IMMUNOTHERAPIES: LOWERING FALSE POSITIVE RATES IN THE SEARCH FOR ACTIONABLE IMMUNOGENIC NEOANTIGENS

¹C Bogaert, ¹L Van Oudenhove*, ¹L Pfitzer, ²N Mill, ¹B Fant. ¹myNEO, Ghent, Belgium; ²University of Zurich, Zurich, Switzerland

10.1136/jitc-2021-ITOC8.26

Background Personalised immunotherapy approaches rely on the ability of tumour-derived neoantigens to elicit a T-cell

immune reaction able to recognise and kill the tumour cells expressing them. Clinical attempts to leverage the power of neoantigens have however yielded mixed results. This can mostly be attributed to the difficulty of finding truly immunogenic peptides from the set of novel peptides generated by mutations in a given cancer patient. In silico approaches can help alleviate this heavy cost by reducing the neoantigen search space, prioritising epitopes based on various parameters such as epitope expression or MHC binding likelihood. Here we present a suite of tools aimed at further assisting clinicians in selecting the most actionable peptides from a set of potential candidates.

Materials and Methods We developed neoMS, a neural network algorithm able to predict epitope presentation at the cell surface with unparalleled performance. Furthermore, the neoIM algorithm is able to discriminate, in an HLA-agnostic fashion, which of the presented peptides will elicit a T-cell immune reaction. This first-in-class algorithm is a random forest classifier specifically trained to classify short peptides of length 9-11 amino acids as immunogenic or non-immunogenic.

Results The neoMS model achieves up to 0.61 precision at recall 0.4 on its test set, vastly outperforming the current industry standards. In addition, due to its sequence-based comparison method, neoMS exhibits extrapolation capabilities, achieving non-zero predictive power when evaluated on ground truth ligandome data derived from an HLA allele completely absent from the training set. In some cancer indications we showed, moreover, that the neoMS-predicted rate of neoantigen presentation can be used in combination with tumour mutational burden as a high-specificity predictor of response to immune checkpoint inhibitor treatment. neoIM vastly outperforms the currently available methods and can predict peptide immunogenicity with high accuracy (AUC=0.88). Interestingly, neoIM confirmed ELISPOT data obtained by Dillon et al. (2017) showing a response in 4 out of 11 breast cancer patients to a vaccine consisting of 9 MHC class-I restricted breast cancer-associated peptides. The 2 antigens that resulted in a CD8⁺ T-cell specific response were predicted by neoIM as the highest scoring showing its potential in finding the truly immunogenic neoantigens.

Conclusions Taken together, these tools decrease false positive rates significantly as they enable improved identification of immunogenic peptides and the predictions correlate with intensity of immune response and clinical benefits. As such, these tools represent a cost-efficient preliminary step in the search for actionable, immunogenic neoantigens.

REFERENCE

1. Dillon PM, Petroni GR, Smolkin ME, Brenin DR, Chianese-Bullock KA, Smith KT, Olson WC, Fanous IS, Nail CJ, Brenin CM, Hall EH, Slingluff CL Jr. A pilot study of the immunogenicity of a 9-peptide breast cancer vaccine plus poly-ICLC in early stage breast cancer. *J Immunother Cancer* 2017 Nov 21;5(1):92.

Disclosure Information C. Bogaert: None. L. Van Oudenhove: None. L. Pfitzer: None. N. Mill: None. B. Fant: None.

P03.04

NOVEL ALGORITHM DISCOVERS UP TO 35 PERCENT MORE EPITOPES TRANSLATED FROM NON-CODING REGIONS IN COLD TUMOURS

C Bogaert, L Van Oudenhove*, B Fant. *mNEO, Ghent, Belgium*

10.1136/jitc-2021-ITOC8.27