MULTIOMIC BIOMARKER SIGNATURES IDENTIFY SUBSETS OF PATIENTS WHO MAY BENEFIT FROM EITHER NIVOLUMAB OR SOTIGALIMAB IN COMBINATION WITH CHEMOTHERAPY IN METASTATIC PANCREATIC CANCER

¹Deena Maurer*, ¹Jia Xin Yu, ²Kamil Sklodowski, ²Marco Tognetti, ²Lukas Reiter, ²Roland Bruderer, ²Jakob Vowinkel, ¹Shannon Pfeiffer, ³Mark O'Hara, ⁴Eileen O'Reilly, ⁵Robert Wolff, ⁶Zev Wainberg, ⁷Andrew Ko, ⁸Osama Rahm, ⁹George Fisher, ¹Jadlyn Lyman, ¹Christopher Cabanski, ¹Pier Federico Gherardini, ¹⁰Jill O'Donnell-Tormey, ¹Theresa LaVallee, ³Robert Vonderheide, ¹Lacey Kitch. ¹Parker Institute for Cancer Immunotherapy, San Francisco, CA, USA; ²Biognosys, Schlieren, Switzerland; ³Abramson Cancer Center at University of Pennsylvania, Philadelphia, PA, USA; ⁴Memorial Sloan Kettering Cancer Center, New York, NY, USA; ⁵University of Texas MD Anderson Cancer, Houston, TX, USA; ⁶University of California, Los Angeles, Santa Monica, CA, USA; ⁷University of California, San Francisco, San Francisco, CA, USA; ⁸Dana-Farber Cancer Institute, Boston, MA, USA; ⁹Stanford University School of Medicine, Stanford, CA, USA; ¹⁰Cancer Research Institute, New York, NY, USA

Background Gemcitabine/nab-Paclitaxel (GnP) is a standard of care regimen for first-line metastatic pancreatic ductal adenocarcinoma (PDAC) and has a 1-year overall survival (OS) rate of approximately 35%. There is an urgent need for novel therapeutics and precision medicine approaches in PDAC. PRINCE, a randomized phase 2 trial, reported an increased 1-year OS relative to historical data, for patients treated with nivolumab (nivo)/GnP (57.3%, $p = 0.007$, $n=34$) and sotigalimab (sotiga) (APX005M; CD40 agonist)/GnP (48.1%, $p = 0.062$, $n= 36$).

Methods To investigate immune modulatory and pharmacodynamic (PD) effects of nivo or sotiga in combination with GnP we used several orthogonal minimally invasive, blood-based biomarker technologies. Immune population profiles were evaluated by CyTOF and features of T cell phenotype and function by multicolor flow cytometry. Soluble proteins were evaluated with predefined panels using the Olink platform (Immuno-oncology (IO) and Immune Response) along with an unbiased mass spectrometry proteomic approach (Biognosys) that identified circulating soluble proteins of significance.

Results Relative to baseline, patients who received nivo/GnP had numerically increased frequencies of proliferating, activated CD8+ and CD4+ effector memory T cells in circulation across multiple timepoints. These patients also had significantly increased levels of soluble proteins associated with type II interferon responses and immune cell migration and T cell activation, as well as significantly decreased levels of immunomodulatory proteins. Patients who received sotiga/GnP had increased expression of the co-stimulatory molecule CD86 on conventional dendritic cells. These patients also had significantly increased concentrations of soluble proteins associated with mature antigen presenting cells, and the activation of helper CD4+ T cells, B cells, and monocytes. Significant increases in soluble proteins associated with type-1 cell-mediated effector immunity and decreases in immunosuppressive factors were observed in both arms. Significant proteins were defined as $p \leq 0.05$, $\log_2$ expression fold change ≥ 0.5 (Olink) and Sparse PLS discriminant analysis was used with zero as a threshold (Biognosys).

Conclusions This study is a first to use multiomic minimally invasive biomarker approaches in PDAC to demonstrate PD effects and immune modulation with immunotherapy/chemotherapy combinations. Orthogonal assays demonstrate that nivo/GnP and sotiga/GnP elicit unique immune responses and the observed effects are consistent with their distinct mechanisms of action. These data suggest that multiomic biomarker

signatures may identify subsets of patients who may benefit from an immunotherapy/chemo approach in PDAC. Moreover, results from these analyses will support early phase clinical study development decisions.

Acknowledgements We extend our gratitude to the patients, their families, the clinical investigators, and their site staff members who are making this trial possible. We would also like to thank Sultan Nawabi at Parker Institute for Cancer Immunotherapy (PICI) for operations leadership of the trial. We thank Bristol Myers Squibb (BMS) and Apexigen for collaboration and study drugs. The study was funded by Cancer Research Institute, BMS and PICI.

Trial Registration NCT03214250

Ethics Approval This study was approved by University of Pennsylvania Institutional Review Board; Federalwide assurance #00004028.

<http://dx.doi.org/10.1136/jitc-2021-SITC2021.343>