AN INVESTIGATION INTO THE ROLE OF CD4+ TUMOR-INFILTRATING LYMPHOCYTES (TIL) IN METASTATIC MELANOMA PATIENTS WITH A COMPLETE RESPONSE TO ADOPTIVE CELL THERAPY

¹MacLean Hall*, ¹Holly Branthoover, ¹Patrick Innamarato, ¹Amy Hall, ¹Alex Alfaro, ¹Allison Richards, ¹Jeani Rich, ¹Jonathan Hensel, ²Jim Bender, ²Jake Ceccarelli, ²TJ Langer, ¹Matthew Beatty, ¹John Mullinax, ¹Jamie Teer, ¹Amod Sarnaik, ¹Shari Pilon-Thomas. ¹Moffitt Cancer Center, Tampa, FL, USA; ²Turnstone Biologics, New York, NY, USA

Background Immunotherapy for cancer has long been focused on the generation of CD8+ cytotoxic T lymphocyte responses, independent of their dynamic CD4+ T cell counterpart. One promising approach, adoptive cell transfer (ACT) of tumor-infiltrating lymphocytes (TIL), has yielded response rates ranging from 28–55%.^{1–2} Investigation into the role of CD4+ TIL in this setting remains critically underexplored as an opportunity to improve upon these successes.

Methods Two metastatic melanoma patients (PT1 and PT2) were treated with TIL on a completed clinical trial at Moffitt Cancer Center (NCT01005745). Tumor recognition by TIL was assessed via co-culture with tumor. Whole exome (WES) and RNA Sequencing were performed on cryopreserved tumor sections and mutant peptide-MHC binding was predicted. TIL were stimulated with antigen presenting cells (APCs) loaded with neoantigen-derived 25mer peptides and sorted based on 41BB/OX40 upregulation, followed by functional immunologic assays. TCR sequencing was conducted on patient peripheral blood as well as isolated neoantigen-specific TIL clones to determine persistence in vivo and cognate peptide-MHC targets were determined empirically.

Results PT1, infused with predominantly CD4+ TIL (88%), achieved a complete response (CR) despite lack of IFN γ detection with conventional in vitro tumor co-culture methods. Infusion product TIL were sorted by upregulation of OX40 and 41BB upon stimulation with APCs loaded with the mutant peptide pool. Neoantigen reactivity arose from a single peptide sequence, which conferred recognition by a CD4+ TIL clone, which comprised 17% of the infusion product and enriched to greater than 80% after sorting via FACS. These CD4+ TIL produced IFN γ , TNF α , and granzyme B in response to peptide-loaded APCs in an HLA-DR dependent manner. TCR β overlap revealed this CD4+ clone peaked at two weeks post-infusion (40%) and persisted after infusion for at least six weeks. PT2 was infused with highly reactive, primarily CD8+ (88%) TIL and also achieved a CR. Isolated CD4+ TIL were also responsive to tumor antigens in the context of MHC Class II in vitro. Tumor-reactive CD4+ TIL were enriched by IFN γ capture and delayed xenograft growth in vivo ($p < 0.01$). Neoantigen peptides stimulated predominantly CD4+ TIL to upregulate OX40/41BB and produce IFN γ , TNF α , and granzyme B.

Conclusions Investigation of these case studies demonstrated evidence of CD4+ TIL involvement in complete clinical responses after ACT. Ongoing studies will define the precise role of tumor-reactive CD4+ T cells in the anti-tumor immune response and provide the framework for future investigation into their function and therapeutic efficacy.

Trial Registration NCT01005745

REFERENCES

- 1.. Bailey SR, *et al.* Human CD26high T cells elicit tumor immunity against multiple malignancies via enhanced migration and persistence. *Nat Commun* 2017;**8** (1):1961.

- 2.. Tran E, *et al.* Cancer immunotherapy based on mutation-specific CD4+ T cells in a patient with epithelial cancer. *Science* 2014;**344**(6184):641–5.

Ethics Approval Approved by USF IRB approval number Ame5_107905. All participants gave informed consent before taking part.

<http://dx.doi.org/10.1136/jitc-2021-SITC2021.384>