





**A** Radiographic patterns between steroid-refractory and response CIP


Patterns	Refractory N (%)	Response N (%)
COP	10 (37.0%)	11 (29.7%)
NSIP	6 (22.2%)	11 (29.7%)
AIP/ARDS	7 (25.9%)	9 (24.3%)
Others	4 (14.8%)	6 (16.2%)

**B** Lesion distribution between steroid-refractory and response CIP

Distribution	Refractory N (%)	Response N (%)
Peripheral	9 (33.3%)	15 (40.5%)
Mixed/multifocal	8 (29.6%)	12 (32.4%)
Diffuse	10 (37.0%)	10 (27.0%)

**C** Ichikado CT score between steroid-refractory and response CIP**D** Radiographic patterns between initial and late steroid-refractory CIP

Patterns	Initial N (%)	Late N (%)
COP	7 (36.8%)	3 (37.5%)
NSIP	3 (15.8%)	3 (37.5%)
AIP/ARDS	7 (36.8%)	0 (0.0%)
Others	2 (10.5%)	2 (25.0%)

**E** Lesion distribution between initial and late steroid-refractory CIP

Distribution	Initial N (%)	Late N (%)
Peripheral	7 (36.8%)	2 (25.0%)
Mixed/multifocal	3 (15.8%)	5 (62.5%)
Diffuse	9 (47.4%)	1 (12.5%)

**F** Ichikado CT score between initial and late steroid-refractory CIP