$\begin{array}{lll} \textbf{Supplementary Table 1.} & \textbf{Clinicopathologic features of patients with HER2-positive and } \\ \textbf{HER2-negative GC} \end{array}$

Characteristics	HER2-negative N=40	HER2-positive n=33	P value
Gender			
Male	30	30	0.077
Female	10	3	
Age			
<60	6	8	0.318
≥60	34	25	
Tumor location			
Upper third	5	5	0.870
Middle third	11	9	
Lower third	15	14	
More than 2/3 stomach	9	5	
Tumor size			
<4cm	14	7	0.195
≥4cm	26	26	
pT stage			
pT1	6	3	0.767^{a}
pT2	6	3	
pT3	4	3	
pT4a	20	18	
pT4b	4	6	
pN stge			
pN0	10	10	0.398
pN1	8	10	
pN2	8	8	
pN3a	9	4	
pN3b	5	1	
pTNM stage			
I	8	4	0.467
II	5	7	
III	27	22	
Histopathological classification			
High-moderately			
differentiated	24	16	0.325
adenocarcinoma			
Low-undifferentiated	16	17	
adenocarcinoma	10	1 /	
Borrman type			
I	2	1	0.554^{a}
II	9	4	
III	10	7	

IV 19 21

Note: a, fisher exact test; *, P<0.05;